


Some aspects of shock propagation through Supernova envelope and ambient interstellar medium (91)

- deep in star where energy from outward propagating material comes from radioactivity, thermalization occurs with temp in opt-UV range.
- when outflow becomes optically thin, effective "temperature" goes up (that is, γ and X-ray photons are not down scattered efficiently so we see high energy non-thermal emission)
- source of energy eventually changes from radioactive decay, to conversion of bulk flow energy at shock (remember shocks are sites of bulk flow dissipation)
- Forward shock and Reverse shocks are present:



Shocks propagate away from the highest density regions. Because of rapid cooling by Bremsstrahlung in the compressed regions, the high density region also supersonically migrates "backward" into the free ejecta in rest frame \rightarrow

- note that the ejecta, contact discontinuity and reverse shock are all moving outward in the lab frame, but in the frame of the contact discontinuity there are shocks propagating both outward = forward and inward toward the explosion point = reverse shock.
 - forward & reverse shocks are important concepts throughout supersonic astrophysics (jets, GRB, etc...)
 - The supernova Remnant SNR (scales $\geq 1000 \text{ AU}$) emits by conversion of bulk flow energy at shock: ejecta has kinetic energy

$$\approx 10^{51} \text{ erg} = \frac{1}{2} M_{ej} v_{ej}^2$$


$$M_{ej} \approx 2 M_{\odot} \Rightarrow v_{ej}^2 \geq 10^{18} \frac{\text{cm}^2}{\text{s}} \Rightarrow v_{ej} \approx 10^9 \frac{\text{cm}}{\text{s}}$$
- \Rightarrow Initial "temperatures" as high as $10^8 - 10^9 \text{ K}$
 (using $v = \left(\frac{kT}{m_p}\right)^{1/2}$).
- but there is an important subtlety as the shock reaches these scales $\geq 1000 \text{ AU}$
 Lets look a bit at the shock physics



• Recall from our brief discussion of shocks form as waves steepen non-linearly



waves are calculated as linear perturbations of the hydro equations. They move at speed $\approx c_s$ for μ -magnetized plasma. Because pressure disturbance from ejecta moves at $v_{eject} \gg \left(\frac{T_{ISM} k_B}{m_p}\right)^{1/2}$ waves pile up:



The role of "nonlinearity" arises in the Navier-Stokes equation (fluid momentum)

$$\frac{\partial \mathbf{v}}{\partial t} = \underbrace{-\mathbf{v} \cdot \nabla \mathbf{v}}_{\text{non-linear term}} - \frac{\nabla P}{\rho} + \underbrace{\nu \nabla^2 \mathbf{v}}_{\substack{\text{dissipation term, increases} \\ \text{entropy}}} \quad (91)$$

Important when there are large gradients even for small viscosity ν .

Viscosity is always approximately \propto speed \cdot length:
Typically, for ambient ISM into which shock propagates $\nu \approx c_s \lambda_{mfp}$.
 \uparrow sound speed \uparrow mean free path
Because "non-linear" effects induce dissipation we know shocks are important where $|\mathbf{v} \cdot \nabla \mathbf{v}| \approx \nu \nabla^2 \mathbf{v}$ (92)



$\therefore q2 \Rightarrow |v \cdot \nabla v| = v \frac{dv}{dx}$
 $v^2/l = \frac{v_{eff}^2}{l_{eff}}$

some "effective" mean free path.

or $l = \frac{v_{eff}}{v_1} \approx \frac{c_{s2} l_{eff}}{v_1}$ (93)

In vicinity of shock, the velocity transits from $v_1 \gg c_{s1}$ to $v_2 \approx c_{s2}$. Thus

eqn (93) $\Rightarrow l \approx l_{eff}$ should be the scale over which the flow changes from upstream to "downstream".

Typically, therefore we expect the shock thickness to be $\approx l_{eff}$. (In reality, instabilities broaden the shock somewhat, but put that aside for the moment).

Now let us estimate this for Supernova Remnants: At ejection velocity

$v_{ej} \approx 10^9 \frac{cm}{s}$ kinetic energy per proton in the ejecta

is about 2 MeV. As these protons hit an H atom of the ISM, the latter will ionize.

largely neutral

Cross section of interaction is $\sigma_{ion} \approx 10^{-17} cm^2 \left(\approx \frac{k^2}{m_p^2 v_{ej}^2} \right)$

Energy lost per ionization is $\approx 50 eV$, (which represents the inelastic part of the collision).

The stopping distance of the impinging protons is therefore

$l_{eff} = \frac{E}{\frac{dE}{dx}} \approx \frac{E}{\Delta E} \frac{1}{n \sigma_{ion}} = \frac{2 MeV}{50 eV} \frac{1}{n \sigma_{ion}}$ (94)

$\Rightarrow l_{eff} \approx 4 \times 10^4 10^{-17} \approx 4 \times 10^{21} cm \approx 10^3 pc!$

But shock thicknesses observed are MUCH smaller than $10^3 pc$. In fact the entire remnants become invisible (merged with ambient medium) on scales of $50 pc$.

Thus, how can thin shock form if the scale ℓ_{eff} were actually $10^3 pc$??


Here the answer is magnetic fields!

Calculate the Larmor radius for microgauss field:

$$r_L \equiv \frac{m c v_{th}}{e B} = \frac{(10^{-24} g) (3 \times 10^{10} \frac{cm}{s}) (10^9 \frac{cm}{s})}{(4 \times 10^{-10}) (3 \times 10^{-6} G)}$$

$$\approx 2.5 \times 10^{10} cm \approx 10^{-8} pc !!!$$

note very small strength field of ISM



B-fields are fundamental

for "collisionless shocks" in astrophysics

They make the "effective mean-free path" equal to the Larmor radius which is much smaller than the collisional mfp even for extremely weak magnetic fields.


and replaces the ℓ_{mfp} in (94) with much smaller value so ℓ_{eff} is then also much smaller

~~Shock~~ on shock jump conditions And Application to Supernova Blastwave

Assume that the shock represents a thin discontinuity. (this was justified in part last lecture):

Conservation of mass, energy & momentum can all be written $\partial_t Q + \nabla \cdot \vec{F}_Q = 0$

If we integrate such a conservation law across the thin discontinuity using the "pill box" as shown:



In steady state:

$$\partial_t Q + \nabla \cdot \vec{F}_Q = 0$$

$$0 \Rightarrow \nabla \cdot \vec{F}_Q = 0$$

but volume is arbitrary

$$\text{so that } \int \nabla \cdot \vec{F}_Q d^3x = 0 \Rightarrow \int_{\partial V} \vec{F}_Q \cdot d\vec{S} = 0 \quad (95)$$

by Gauss' theorem

For mass continuity:

$$\partial_t \rho + \nabla \cdot (\rho \vec{u}) = 0 \Rightarrow \oint \rho \vec{u} \cdot d\vec{S} = 0$$

$$\rho_1 u_1 ds_1 - \rho_2 u_2 ds_2$$

$ds_1 = ds_2$
for pill box

$$\Rightarrow \boxed{\rho_1 u_1 = \rho_2 u_2}$$

(96)

Similarly: for flows in which β -field is energetically negligible: (97)

$$w_1 + \frac{1}{2} v_1^2 = w_2 + \frac{1}{2} v_2^2 \quad \text{energy conservation (97)}$$

$$p_1 + \rho_1 v_1^2 = p_2 + \rho_2 v_2^2 \quad \text{momentum flux conservation (98)}$$

$$(w = \text{enthalpy density} = \frac{\Gamma}{\Gamma-1} \frac{p}{\rho} = \frac{c_s^2}{\Gamma-1})$$

96-98 are the Rankine-Hugoniot jump conditions

for a shock. Define $M_1^2 \equiv v_1^2 / c_{s1}^2$

Solving (96-98) (I leave as exercise)

$$\frac{\rho_2}{\rho_1} = \frac{(\Gamma+1) M_1^2}{(\Gamma+1) + (\Gamma-1)(M_1^2-1)} = \frac{v_1}{v_2} \quad (99)$$

$$\frac{p_2}{p_1} = \frac{(\Gamma+1) + 2\Gamma(M_1^2-1)}{\Gamma+1} \quad (100)$$

$$\frac{c_{s2}^2}{c_{s1}^2} = \frac{T_2}{T_1} = \frac{[(\Gamma+1) + 2\Gamma(M_1^2-1)][(\Gamma+1) + (\Gamma-1)(M_1^2-1)]}{[(\Gamma+1)^2 M_1^2]} \quad (101)$$



Assume flow is supersonic on side 1

(98)

$$\text{so } M_1 = \frac{V_1}{c_{1s}} > 1.$$

Then

$$\frac{P_2}{P_1} > 1, \quad \frac{\rho_2}{\rho_1} > 1, \quad \frac{V_2}{V_1} < 1, \quad \frac{T_2}{T_1} > 1.$$

strongest shock $\Rightarrow M_1^2 \gg 1$

$$\Rightarrow \frac{\rho_2}{\rho_1} = \frac{\Gamma + 1}{\Gamma - 1}; \quad \frac{P_2}{P_1} \gg 1, \quad \frac{T_2}{T_1} \gg 1 \quad (102)$$

(limiting relation as $M_1^2 \rightarrow \infty$!) \Rightarrow for $\Gamma = 5/3 \Rightarrow \frac{\rho_2}{\rho_1} = 4$

Note: momentum conservation and mass conservation are usually satisfied as in 96 & 98, but energy conservation can have important radiative terms, chemical reaction terms, thermal conduction..., we ignore these for the moment.


The above treatment assumes that the viscous terms operate only in the thin layer of the shock itself; this gets back to our notion from the previous lecture that the shock thickness can be estimated by comparing dissipative & bulk velocity terms:

In momentum equation, compare $v \cdot \nabla v$ term to $\nabla \nabla^2 v$ term: (see page 93)

$\Rightarrow \frac{v^2}{L} \approx \frac{\nu_{eff}}{L^2} \Rightarrow v = \frac{\nu_{eff}}{L}$, where ν_{eff} is the effective "viscosity" at the shock.

Now across the shock, the bulk energy of the flow in v_1 gets converted to random thermal energy such that $C_s \approx v_1$. As discussed on p. 95 of this lecture the previous lecture, ν_{eff} is determined by multiples of Debye radius rather than collisional mean free path.

The shock is actually a "current sheet": when B-field is included in jump conditions. This is because Maxwell's equations require that tangential component of E is conserved across the shock:

Consider "pill surface" crossing shock \rightarrow 
from Maxwell's equations:

$\frac{1}{c} \frac{\partial B}{\partial t} + \nabla \times E = 0 \Rightarrow \int (\nabla \times E) \cdot d\vec{S} = 0$
0 in steady state Stokes theorem surface is arbitrary

$\Rightarrow \oint E \cdot dl = 0$
for arbitrarily thin pill surface only: sides contribute:

$\Rightarrow \oint E \cdot dl = 0 = E_{1,T} l - E_{2,T} l = 0 \Rightarrow E_{1,T} = E_{2,T}$

Since Ohms law implies

$$E = -\frac{v \times B}{c} + \eta J \quad \text{then}$$

$$E_{1,T} = E_{2,T} \quad \text{'''}$$

$$\Rightarrow \left(-\frac{v \times B}{c} + \eta J\right)_{1,T} = \left(-\frac{v \times B}{c} + \eta J\right)_{2,T}$$

but $J = \frac{c}{4\pi} \nabla \times B$ and away from shock, $\frac{c}{4\pi} \nabla \times B$ can be considered small;

η is the resistivity and most astro-plasmas have low resistivity. However, near the


shock $\nabla \times B \approx \frac{B}{l_{\text{eff}}} \approx \frac{B}{r_L}$

(see eqn 94 and page 95 of this lecture)

The gradient scale is small and near the shock ηJ is important. This is why

a shock is a "current sheet." Magnetic

Reconnection provides another example of a current sheet based on same principle:




← magnetic field annihilation at dotted intertacle: **Exercise**: show that

intertacle is a current sheet if intertacle is thin!

Now back to the evolution of the expanding SN shock: Transition to Sedov phase

During the early stages of the propagation of the optically thin phase of the shock's progress through the envelope and into ISM, the ejecta material has much more inertia than the ISM with which it interacts. The ejecta speed V_e is thus constant.

BUT: There exists a critical radius r_c at which the ejecta mass $M_{ejecta} = \frac{4}{3}\pi \rho_{ISM} r_c^3$. At this point the blast enters the Sedov phase. Now the mass is piling up behind the shock and this mass starts to dominate the total mass of the ejecta. The mass piles up behind the shock, but ahead of the contact discontinuity:



Once the Sedov phase is underway the speed of the blast wave is no longer constant: In the Sedov phase mass is dominated by that

accumulated from ISM. Also the Energy is

$$E \approx \frac{1}{2} \frac{4\pi}{3} (\rho_{ISM} r^3) v_1^2 = \text{constant} \quad (103)$$

radius of expanding shell \uparrow just the initial explosion energy

constant $\rho_{ISM} \Rightarrow$

$$E \propto r^3 v_1^2 \Rightarrow r^3 \left(\frac{dr}{dt}\right)^2 = \text{constant}$$

$$\Rightarrow r^{3/2} dr = dt$$

$$\Rightarrow r = (\text{constant}) t^{2/5} \quad (104)$$

Another way to arrive at this is to note that ρ_{ISM} and E are constant and

$$E \approx \frac{1}{2} M \left(\frac{r}{t}\right)^2 = \text{const} \quad (105)$$

$$\rho_{ISM} = \frac{M}{\frac{4\pi}{3} r^3} = \text{const.} \quad (106)$$

$$\Rightarrow \frac{E}{\rho_{ISM}} = \text{const} = \frac{2\pi}{3} \frac{r^5}{t^2} \Rightarrow r = \left(\frac{E t^2}{\rho_{ISM}}\right)^{1/5} \quad (107)$$

\rightarrow

$$\Rightarrow r = \left(\frac{E}{\rho_{ISM}} \right)^{1/5} t^{2/5} = 3 \text{ pc} \left(\frac{E}{10^{51} \text{ erg}} \right)^{1/5} \eta_{ISM}^{-2/5} \left(\frac{t}{300 \text{ yr}} \right)^{2/5} \quad (108)$$

↑ applies only for $r > r_{\text{crit}} \approx \left(\frac{3 M_{\text{ejecta}}}{4 \pi \rho} \right)^{1/3}$

$$\Rightarrow V_1 \approx \frac{r}{t} \approx 3 \times 10^3 \frac{\text{km}}{\text{s}} \left(\frac{E}{10^{51} \text{ erg}} \right)^{1/5} \eta_{ISM}^{-1/5} \left(\frac{t}{300 \text{ yr}} \right)^{-3/5} \quad (109)$$

Using $V_1 \approx C_{s2} \Rightarrow$

$$T \approx \frac{m_p}{k} C_{s2}^2 \approx \frac{m_p}{k} V_1^2 \approx 9 \times 10^8 \text{ K} \left(\frac{E}{10^{51} \text{ erg}} \right)^{2/5} \eta_{ISM}^{-2/5} \left(\frac{t}{300 \text{ yr}} \right)^{-6/5} \quad (110)$$

\Rightarrow at $t = 3.5 \times 10^4 \text{ yr}$, $T \approx 3 \times 10^6 \text{ K}$


Thus if SNR is observed with $T \approx 3 \times 10^6 \text{ K}$ (as is Cygnus Loop), the time in sedov phase to reach that stage is, from (110)

$$t_{\text{sedov}} = 3.5 \times 10^4 \text{ yr} \left(\frac{T}{3 \times 10^6} \right)^{-5/6} \left(\frac{E}{10^{51} \text{ erg}} \right)^{1/3} \eta_{ISM}^{-1/3} \quad (111)$$

\Rightarrow for given V or T and r observed (to determine if $r > r_{\text{crit}}$)
age can be determined

\rightarrow

Now, as deceleration becomes significant outer shells of expanding sphere decelerate first => material in the inner region catches up with material in the outer regions




Region (A) is supersonic with respect to (B) => reverse shock moves "backward" in frame of contact discontinuity, in lab frame everything is moving outward.


At the reverse shock, kinetic energy of ejecta is re-heated by reverse shock dissipation as it passes through -> implies some of the bulk energy of the ejecta goes back into heat of ejected material. The forward shock converts some of the bulk energy into heating ambient ISM material. (X-ray emission is visible from both shocked regions)

We mentioned, and will discuss later, the Rayleigh Taylor instability, which takes place during the Sedov phase. The

Rayleigh - Taylor fingers :



low density ISM
=>
expansion direction



(Both forward and reverse directions have equivalent of "heavy" fluid falling on "light" fluid)

Radiative phase of SNR blast wave

once radiative cooling time becomes short compared to Sedov age we have radiative phase. Sedov age is given by (111).

For cooling time, note that for $T < 10^6$ K

C, N, O gain e^- and become atomic; cooling by atomic cascade of e^- falling to lower levels dominates:

$$n_H^2 \Lambda(T) = 10^{-22} \text{ erg/cm}^3\text{-s} \quad n_H^2 \left(\frac{T}{10^6 \text{ K}} \right)^{-1/2} \quad (112)$$

$$t_{\text{cool}} \approx \frac{n k T}{n^2 \Lambda(T)} \approx 2 \times 10^5 \left(\frac{T}{3 \times 10^6} \right)^{3/2} n_H^{-1} \text{ yr} \quad (106) \quad (113)$$

↑
number density
for compressed
region

$t_{\text{cool}} < t_{\text{setov}}$ when
from (111) and (113)

$$T^{7/3} < \frac{2 \times 10^5}{3.5 \times 10^4} (3 \times 10^6)^{7/3} \frac{n_H}{n_{\text{ISM}}^{1/3}} \left(\frac{E}{10^{51} \text{ erg}} \right)^{1/3}$$

$$\text{or } T < \left(\frac{3.5 \times 10^4}{2 \times 10^5} \right)^{3/7} (3 \times 10^6)^{7/3} (4) (n_H^{2/3})^{3/7} \left(\frac{E}{10^{51}} \right)^{1/7}$$

↓
Compression
ratio across
shock just before
cooling becomes
important (see eqn. 102)

$$T < 5.7 \times 10^6 \text{ K } (n_H^{2/3})^{3/7} \left(\frac{E}{10^{51} \text{ erg}} \right)^{1/7} \quad (114)$$


$$\text{or } v \approx \left(\frac{k T}{m} \right)^{1/2} \leq 240 \frac{\text{km}}{\text{s}} (E_{51} n_H^2)^{1/14}$$

notice the weak dependence
on E and n_H !

In radiative phase

Shock becomes isothermal as it evolves.

Hot interior region but a cooled, isothermal interior shell: In frame of contact discontinuity:



cooling takes away most of the shock energy but momentum is conserved because

radiation is essentially isotropic. Thus

$$\frac{d}{dt} \left(\underbrace{\frac{4\pi \rho_{ISM}}{3} r^3 \dot{r}}_{\text{momentum}} \right) \approx 0 \quad \text{in radiative phase} \quad (115)$$

$$\Rightarrow r^3 \dot{r} = \text{constant} \quad \text{for} \quad \frac{d\rho_{ISM}}{dt} \approx 0.$$

$$\Rightarrow r^3 dr = dt$$

$$\Rightarrow r \propto t^{1/4}$$

and

$$\dot{r} \propto t^{-3/4}, \quad \left[\dot{r} = 240 \frac{\text{km}}{\text{s}} \left(E_{51} n_H^2 \right)^{1/4} \left(\frac{t}{5 \cdot 2 \times 10^4 \text{ yr}} \right)^{-3/4} \right]$$

using (114) & (112)

(116)




Fig. 4.6. The radius of the supernova shell as a function of time during the different phases.

This integrates to give

$$R = R_0 \left[1 + 4 \frac{v_0}{R_0} (t - t_0) \right]^{1/4}, \quad \dot{R} = v_0 \left[1 + 4 \frac{v_0}{R_0} (t - t_0) \right]^{-3/4}. \quad (4.107)$$

For large t , $R \propto t^{1/4}$ and

$$\dot{R} \propto t^{-3/4} \simeq 200 \text{ km s}^{-1} (t/3 \times 10^4 \text{ yr})^{-3/4}. \quad (4.108)$$

The time constant in relation (4.108) is fixed by equating the Sedov phase velocity of Eq. (4.101) to 200 km s^{-1} .

In the final phase, the speed of the shell drops below the sound velocity of the ISM, which is approximately $(10-100) \text{ km s}^{-1}$ in a time scale of $t \approx (1-5) \times 10^5 \text{ yr}$. Around this time scale, the remnant loses its identity, and it is dispersed by random motions in the ISM. The evolution is shown schematically in Fig. 4.6.

It should be noted that supernova explosions and their eventual dispersion of ejected material have the effect of enriching the ISM with the material processed in stellar interiors. In particular, the heavy elements synthesised inside a star reach the ISM through this process. Because massive stars evolve at shorter time scales and also are more likely to end up as supernovas, the evolution of the first generation of massive stars changes the character of the ISM. Second and later generations of stars condense out of this enriched ISM and will have a higher proportion of heavier elements.

A supernova explosion creates an expanding luminous region from the centre of the explosion. The heat and ionise such a region from OIII was detected approximately 1 yr after the supernovas also lead to light phenomena discussed in Volume two light echos were detected

approximately 1 yr after the supernova emits x rays from the plasma at a temperature are formed during phase 3, and in the material with a temperature of the radiating atoms. In addition remnants are also strong sources spiraling in the magnetic field Vol. I, Chap. 6, Section 6.11, electrons per unit volume is 1

then the total flux of an optical be expressed as

$$S_\nu = \frac{G}{d^2} V K B^{(1+p)/2} \nu^{-}$$

where V is the volume of the is a numerical factor. In the case is strongly ionised during the frozen to the plasma fluid. It f

If the energy of individual relativistic expansion of the volume, the energy the pressure of relativistic electrons to give $\epsilon \propto r^{-4}$. The total energy