Exam 2 (April 22, 2009)

Please read the problems carefully and answer them in the space provided. Write on the back of the page, if necessary. Show your work where requested in order to be considered for partial credit. In problems where you are requested to show your work, no credit will be given unless your work is shown.

Problem 1 (true or false, each part is worth 2 points):

- a) _____ In typical multi-electron atoms, the electrons all reside in the atom's lowest energy quantum state.
- b) _____ If a sample of uranium-235 is subcritical, a nuclear explosion is imminent.
- c) _____ According to quantum field theory, the gluon is the virtual particle (gauge boson) responsible for conveying the strong nuclear force.
- d) _____ The force of gravity is many times stronger than the weak nuclear force.
- e) _____ A typical chemical reaction involves changes in energy of millions of electron-volts.
- f) _____ Water is a chemical compound.
- g) _____ Carbon dioxide (CO₂) is an isotope of carbon monoxide (CO).
- h) _____ According to quantum field theory, the Z particle is the virtual particle (gauge boson) responsible for conveying the strong nuclear force.
- i) _____ According to quantum theory, the more precisely the position of an electron is determined the better known is the electrons velocity.
- j) _____ The Higgs particle was discovered (first seen) in 2006.
- k) _____ A chain reaction refers to the chemical processes that occur when hydrogen is mixed with oxygen and a match.
- 1) _____ The strong nuclear force is stronger than the electromagnetic force.
- m) _____ Nuclear fission is the energy source that powers stars.
- n) _____ In quantum mechanics, the wave function specifies the exact position of a particle.
- o) _____ Young stars are formed mostly of hydrogen.

Problem 2 (4 points):

A quantum state is

- a) a term that refers to atoms that are able to emit photons.
- b) a state that is smaller than most other states for example, Rhode Island.
- c) a phrase describing the frame of mind of a physicist who is contemplating the complicated aspects of matter.
- d) a reference to the potential spatial regions and energies allowed for particles according to quantum mechanics.
- e) the place where a particle is seen to be located.

You walk into a room that is contaminated with radioactive material. What factors are important in determining the potential danger to you of the radioactive material?

- a) the location of the material relative to you.
- b) the activity of the sample.
- c) the time you spend in the proximity of the sample.
- d) the type of radiation emitted by the sample.
- e) All of the above.

Problem 4 (4 points):

Which of the four fundamental forces binds the nucleus together, and which binds the atom [electron orbits] together?

- a) Strong nuclear force binds the nucleus, weak nuclear force binds the atom.
- b) Gravity binds the nucleus, electromagnetic force binds the atom.
- c) Weak nuclear force binds the nucleus, strong nuclear force binds the atom.
- d) Electromagnetic force binds the nucleus, strong nuclear force binds the atom.
- e) Strong nuclear force binds the nucleus, electromagnetic force binds the atom.

Problem 5 (4 points):

Quarks interact with other particles in nature via

- a) the strong nuclear force.
- b) the weak nuclear force.
- c) the electromagnetic force.
- d) gravitation.
- e) all of the above.

Problem 6 (4 points):

The source of energy for the creation of most gold atoms (a gold atom is heavier than an iron atom) used in jewelry is

- a) the gravitational collapse of a star.
- b) a goldsmith's smelter.
- c) The fission of a heavier element such as radium or hafnium.
- d) the shock wave of a supernova explosion at the end of the stellar life cycle for a large star.
- e) the heat at the center of the earth.

Scores							
1.	/30						
2.	/4						
3.	/4						
4.	/4						
5.	/4						
6.	/4						
7.	/4						
8.	/5						
9.	/6						
10.	/5						
11.	/10						
12.	/10						
13.	/10						
Total							
/100							

P100University of RochesterS. ManlySpring 2009

NAME _

Problem 7 (4 points):

Carbon is

- a) an element.
- b) a chemical compound.
- c) made up entirely of protons.
- d) chemically inert (unreactive).
- e) never found in stars.

Problem 8 (5 points):

A nuclear fusion process occurs and is described by the equation below. Determine and provide the unknown nucleus (symbolized by X) in the equation.



Problem 9 (6 points):

Suppose the nucleus below undergoes alpha decay, what is the nucleus left behind?



Would you expect this process to release energy or absorb energy?

Problem 10 (5 points):

In the quark model of particle physics, is it possible to have a baryon with an electric charge of +2? If so, give a possible example. If not, why?

Problem 11 (10 points):

Briefly describe the relationship between quantum uncertainty and radioactive decay.

Problem 12 (10 points):

The iodine isotope ¹³¹I is a naturally radioactive nucleus that emits a beta (e-) with a half-life of 8 days. Because iodine is absorbed by the thyroid gland, people with malfunctioning thyroids can be treated medically by drinking a solution laced with ¹³¹I. The radioactive iodine decays and causes tissue damage preferentially in the thyroid.

- a) What nuclear isotope does the ¹³¹I become when it decays?
- b) If a sample of ¹³¹I sitting on the hospital shelf has an activity of 200 decays per second today, how much time will pass before that activity drops to 25 decays per second?

Problem 13 (10 points):

In 1929, one of the big players in physics at the time, P.A.M. Dirac, said of quantum mechanics, "The underlying physical laws necessary for the mathematical theory of a large part of physics and the whole of chemistry are thus completely known …" Briefly defend or dispute Dirac's assertion that quantum mechanics explains chemistry.

	University of Rochester	NAME	
nly	Spring 2009		

hellum 2 4.0026	10 10	Ne	20.180 argon	Ar	39.948 krvnton	36	Y	83.80 Xenon	Xe	131.29	radon 86	Rh	222							
1	fluorine 9	ш	18.998 chlorine	บ	35.453 hromine	35	Ы	/9.904 iodine 53		126.90	astatine 85	At	[210]							
E	oxygen 8	0	15.999 sulfur 16	ŝ	32.065 selenium	34	Se	18.96 tellurium 53	Te	127.60	polonium 84	Ро	209			ytterbium 70	Υb	173.04 nobelium	201	NO [259]
i.	nitrogen 7	z	14.007 phosphorus	<u>۽</u> ح	30.974 arsenic	33	As	antimony	Sb	121.76	bismuth 83	ï	208.98			thulium 69	Tm	168.93 mendelevium		
1	carbon 6	ပ	12.011 silicon 1.1	Si	28.086 dermanium	32	Ge	12.61 tin 50	Sn	118.71	ead 82	РЬ	207.2 ununquadium 114	Duq	602	erbium 68	Ш	167.26 fermium		
1	boron 5	В	10.811 aluminium 13	AI	26.982 dallium	31	Ga	69.723 indium 49	ln	114.82	81	F	204.38			holmium 67	Ч	164.93 einsteinium	⁶⁶	
ļ					zinc	30	Zn	65.39 cadmium 4 8	Cd	112.41	mercury 80	Hg	200.59 ununbium 112	Uub		dysprosium 66	D	162.50 californium	98	5 ^{251]}
					CODDAL	29	Cu	63.546 silver 47	Ag	107.87	000 79	Au	196.9/ unununium 111	Uuu	4	terbium 65	Tb	158.93 berkelium		
1					nickel	28	Z	palladium	Pd	106.42	platinum 78	Ł	135.08 ununnilium 110	Uun	- a	gadolinium 64	Qd	157.25 curium	96 (C ²⁴¹
6					cohalt	27	ပိ	thodium	Rh	102.91	77	<u> </u>	192.22 meitnerium 109	Mt		europium 63	Eu	151.96 americium	с <u>6</u>	
E2					imn	26	Ъ С	55.845 ruthenium	Ru	101.07	osmum 76	Os	190.23 hassium 108	HS Deel		samarium 62	Sm	150.36 plutonium	⁸	L ^[244]
					asenabnam	25	Mn	54.938 technetium 43	Lc	[86]	rhenlum 75	Re	186.21 bohrium 107	Bh		promethium 61	Pm	[145] neptunium	۶۶ •	ND
C.					chromium	24	ບັ	51.996 molybdenum	Mo	95.94	tungsten 74	3	183.84 seaborgium 106	Sg		n neodymium 60	Νd	144.24 I uranium	76	238.03
ę.					vanadium	23	>	50.942 niobium 41	N b	92.906	tantalum 73	Ta	180.95 105	Db		praseodymiun 59	Ъ	140.91 protactinium	ۍ ۲	231.04
					titaninm	22	Ē	47.867 zirconium	Zr	91.224	72	Ηŧ	1/8.49 rutherfordiun 104	Rf	22	cerium 58	Ce	140.12 thorium	0 ⁶	1 232.04
ŝ					scandium	21	Sc	44.956 yttrium 30	~	88.906	71	Lu	1/4.9/ lawrencium 103		7.72	lanthanum 57	La	138.91 actinium	68	AC
											57-70	*	89-102	*		cariac	200		eries	
1	beryllium 4	Be	9.0122 magnesium	Mg	24.305 caldium	20	Ca	40.078 strontium 38	s S	87.62	56 56	Ba	137.33 radium 88	Ra	0.27	apida		-	cinide s	
hydrogen 1 0079	Bithium	[]	6.941 sodium	Na	22.990 notassium	19	Y	39.098 rubidium 37	Rb	85.468	caesium 55	Cs	132.91 francium 87	L L	0.44	* 201	L		ACI	

Some potentially Useful formulas

$$F = G M, M_{2} \left[M, and M_{2} in k_{9} \right] \rightarrow G = 6.7 \times 10^{-11}$$

$$F = \frac{M}{r^{2}} \left[9, 92 M Coulombs \right] \rightarrow k = 9 \times 10^{9}$$

$$F = Ma \left[F in Newtons \right]$$

$$F = Ma \left[F in Newtons \right]$$

$$F = Ma \left[F in Newtons \right]$$

$$I Joule = 1.6 \times 10^{-19} eV$$

$$(distance) = (Speed) (time)$$

$$V = \frac{\Delta x}{\Delta t}$$

$$Q = \frac{\Delta V}{\Delta t}$$

$$Morrk = force \times distance$$

$$Momentum = P = MV$$

$$\Delta x' = 8 \Delta x, \Delta x Ionsect$$

$$in Propen Frame$$

$$V = \frac{\Delta x}{\Delta t}$$

$$V = \frac{1}{\sqrt{1 - (\frac{x}{2})^{2}}}$$

$$\frac{\Delta N}{\Delta t} = \lambda N \quad t_{1/2} = 0.643/\Lambda$$