Astronomy 102: Black Holes, Time Warps, and the Large-Scale Structure of the Universe

What do black holes, wormholes, time warps, spacetime curvature, hyperspace and the Big Bang have in common?

☐ Explanations with their origins in Einstein's theories: the **special theory of relativity** (1905) and the **general theory of relativity** (1915).

This semester we will discuss all of these exotic phenomena, mostly qualititatively, in the context of Einstein's theories.



Artist's conception of the quasar 3C273, from Thorne, Black holes and time warps.


1 September 2009

Astronomy 102, Fall 2009

Our primary goals in teaching Astronomy 102

- □ to demystify black holes, the Big Bang, and relativity, so you can evaluate critically the things you find about them in the media;
- ☐ to show you how scientific theories are conceived and advanced in general.

In doing so we aim primarily at **non-science majors.**




This semester's AST 102 class, plotted by concentration.

1 September 2009

Astronomy 102, Fall 2009

Our primary goals in teaching Astronomy 102 (continued)

We hope that by the end of the course you will understand and retain enough to be able to offer correct explanations of black holes and such to your friends and family, and that you will retain a permanent, basic understanding of how science works.





This semester's AST 102 class, plotted by cohort.

1 September 2009

Astronomy 102, Fall 2009

Human and printed features of Astronomy 102 People: Dan Watson, professor Brian DeCesare, teaching assistant Jae Song, teaching assistant Textbooks (one required, three recommended): Kip S. Thorne, Black holes and time warps (1994). Michael A. Seeds, Foundations of astronomy (2008). Also used in AST 104, 105, and 106. Stephen Hawking, A brief history of time (1988). Joseph Silk, The Big Bang (2005).





90% of success is showing up.

All members of the class are expected to attend all of the lectures, and encouraged to attend one recitation per week.

☐ This is for your own good.

You will very probably
get a better grade if you
go to class, as is demonstrated
by these average test score

90 80 80 70 80 90 100 Attendance (% of class)

and average attendance data from past AST 102 classes. ☐ You may attend any recitation you like, whether you're registered for it or not.

1 September 2009

Astronomy 102, Fall 2009

Mid-Lecture Break

This will be a regular feature of Astronomy 102 lectures.



Image: wide-field view of the Orion Nebula, by David Malin (Anglo-Australian Observatory).

1 September 2009

Astronomy 102, Fall 2009

Today in Astronomy 102: How big is that? Before discussing black holes, the Big Bang, and other celestial objects and phenomena, we need to become ☐ familiar with distances, time scales, masses, luminosities and speeds of astronomical importance, and ☐ proficient at unit conversion. 1 2 3 4 5 Million light years 1 September 2009 Astronomy 102. Fall 2009 9

Sizes and distances in astronom	m	om	ı
---------------------------------	---	----	---

	centimeters	kilometers	miles	light years
Diameter of a hydrogen atom	1.1×10 ⁻⁹			
Diameter of a human hair	8.0×10 ⁻³			
Diameter of a penny	1.9			
Diameter of Rochester	2.0×106	20	12	
Diameter of the Earth	1.3×109	1.3×104	7.9×10^{3}	
Diameter of the Moon	3.5×10^{8}	3.5×10^{3}	2.1×10^3	
Diameter of Jupiter	1.4×10^{10}	1.4×10^{5}	8.8×10^{4}	
Diameter of the Sun	1.4×10^{11}	1.4×10^{6}	8.6×10 ⁵	
Diameter of the Milky Way galaxy	1.6×10^{23}			1.7×10 ⁵
Distance to Buffalo	1.0×10 ⁷	100	62	
Distance to the Moon	3.8×10^{10}	3.8×10 ⁵	2.4×10 ⁵	
Distance to the Sun	1.5×10^{13}	1.5×10^{8}	9.2×107	
Distance to the next nearest star, α	3.8×10^{18}			4
Centauri				
Distance to the center of the Milky Way				2.7×10 ⁴
Distance to the nearest galaxy	1.6×10 ²³			1.7×10 ⁵
1 September 2009 Ast	ronomy 102, Fa	11 2009		1

Typical lengths and important conversions

- ☐ Diameter of normal stars: millions of *kilometers* (km)
- $f \square$ Distance between stars in a galaxy: a few light-years (ly)
- ☐ Diameter of normal galaxies: tens of kilo-light-years (kLy)
- ☐ Distances between galaxies: a million light-years (Mly)
- \square 1 ly = 9.46052961×10¹⁷ cm = 9.46052961×10¹² km
- \square 1 km = 10⁵ cm; 1 kly = 10³ ly; 1 Mly = 10³ kly = 10⁶ ly.

Example: The Andromeda nebula (a galaxy a lot like our Milky Way) lies at a distance D = 2.5 Mly. How many centimeters is that?

$$D = 2.5 \text{ Mly} \times \frac{10^6 \text{ ly}}{1 \text{ Mly}} \times \frac{9.46 \times 10^{17} \text{ cm}}{1 \text{ ly}} = 2.4 \times 10^{24} \text{ cm}$$

1 September 2009

Astronomy 102, Fall 2009

More detail on numerical answers

Note that the last answer was written as 2.4×10^{24} cm.

- \square Not just 2.4×10^{24} . Numerical answers in the physical sciences and engineering are incomplete without units.
- ☐ And not 2.36513240×10²⁴ cm, even though that's how your calculator would put it. Numerical answers should be rounded off: display no more than one more significant figure than the least precise input number.
 - If we had been told that the distance to the Andromeda galaxy is 2.5000000 Mly, then the conversion factor would have to have been put in with more significant figures, and the right answer would have been $2.36513240 \times 10^{24}$ cm.

1 September 2009

Astronomy 102, Fall 2009

12

11

More detail on Unit Conversion

Previous example: repeated multiplication by 1. One may always multiply anything by 1 without changing its real

The unit conversions always give a couple of useful forms of 1. Take, for example, the conversion 1 ly = 9.46×10^{17} cm: $\frac{9.46 \times 10^{17} \text{ cm}}{1 \text{ cm}} = 1 = \frac{1 \text{ ly}}{17}$

$$\frac{9.46 \times 10^{17} \text{ cm}}{1 \text{ ly}} = 1 = \frac{1 \text{ ly}}{9.46 \times 10^{17} \text{ cm}}$$

Choose forms of 1 that cancel out the units you want to get rid of, and that insert the units to which you wish to convert. This sometimes takes repeated multiplication by 1, as in the previous example:

$$D = 2.5 \text{ MHy} \times \frac{10^6 \text{ ky}}{1 \text{ MHy}} \times \frac{9.46 \times 10^{17} \text{ cm}}{1 \text{ ky}} = 2.4 \times 10^{24} \text{ cm}$$

1 September 2009

Astronomy 102, Fall 2009

There is nothing sacred about centimeters, grams and seconds.

Units are generally chosen to be convenient amounts of whatever is being measured. Examples:

☐ The **light-year** (ly) is far more convenient than the centimeter for expression of length in astronomy; on large scales we even use millions of ly (Mly).



14

☐ The convenient unit of mass in astronomy is the solar mass: the mass of the Sun.

Values of physical quantities are ratios to the values of the unit quantities.

1 September 2009

Astronomy 102, Fall 2009

Masses in astronomy Grams Pounds Solar masses (M_{\odot}) Hydrogen atom 1.67×10^{-24} Penny (uncirculated) 3.2 0.0071 1.02×10^{6} 2240 Ton Earth 6.0×10^{27} 1.3×10^{25} 3.0×10^{-6} 7.4×10^{25} 3.7×10^{-8} Moon 1.9×10^{30} Jupiter 1.0×10^{-3} Sun 2.0×10^{33} 1 Milky Way galaxy 3×10^{12} 6×10^{45} 1 September 2009 Astronomy 102, Fall 2009 15

Typical masses and important conversions

- \blacksquare Smallest stars: 0.08 solar masses (M_{\odot})
- \blacksquare Normal stars: around one M_\odot
- \blacksquare Giant stars: tens of M_{\odot}
- \Box Normal galaxies: 10^{11} $10^{12}\,M_\odot$
- \blacksquare Clusters of galaxies: 10^{14} $10^{15}\,M_{\odot}$
- \square 1 M_{\odot} = 2.0×10³³ grams = solar mass = mass of the Sun
- ☐ 1 pound = 454 grams

Example: Vega, the brightest star in the Northern sky, has a mass of about 2.5 $\rm M_{\odot}$. What is its mass in grams?

$$M = 2.5 M_{\odot} \times \frac{2 \times 10^{33} \text{ gm}}{1 M_{\odot}} = 5.0 \times 10^{33} \text{ gm}$$

1 September 2009

omy 102 Fall 2009



How many Earths in the Sun?

That is, by what factor is the Sun more massive than the Farth?

Find the masses a couple of pages back, work it out and send the answer on your clicker.

1 September 2009

Astronomy 102, Fall 2009

17



Times and ages in astronomy

7 6 5	seconds	hours	days	years
Earth's rotation period	8.64×10 ⁴	24	1	
Moon's revolution period	2.3606×106	655.73	27.322	
Earth's revolution period	3.1558×10^{7}	8.7661×10^{3}	365.25	1
Century	3.16×10^9			100
Recorded human history	1.6×10^{11}			5000
Milky Way Galaxy's rotation period (at Sun's orbit)	7.5×10 ¹⁵			2.4×10 ⁸
Age of the Sun and Earth	1.44×10^{17}			4.56×10^{9}
Total lifetime of the Sun	4.7×10^{17}			1.5×10^{10}
Age of the Universe	4.4×10^{17}			1.4×10^{10}
1 September 2009	Astronomy 102, Fa	ill 2009		18

(c) University of Rochester

Typical timespans and important conversions

- lue Planetary revolution period: around 1 year
- $\hfill \Box$ Life expectancy, normal stars: around $10^{10}~{\rm years}$
- $\hfill \Box$ Life expectancy, giant stars: 10^6 10^8 years
- \square Rotation period of normal galaxies: 10^7 10^9 years
- \Box 1 year = 3.16x10⁷ seconds
- ☐ 1 hour = 3600 seconds

Example: How many seconds is a normal human lifespan (US)?

$$t = 75 \text{ years} \times \frac{3.16 \times 10^7 \text{ seconds}}{1 \text{ year}} = 2.37 \times 10^9 \text{ seconds}$$

1 September 2009

Astronomy 102, Fall 2009

How many lifespans in the Solar system's age?

That is, by what factor is the age of the Sun and Earth longer than an average human lifespan?

Find the numbers in the last couple of pages, work it out, and send the answer using your clicker.

1 September 2009

Astronomy 102, Fall 2009

09

The fundamental dimensions

Distance, time and mass are fundamental dimensions.

- ☐ Distances along each of the three different perpendicular directions of space determine the location of a given body with respect to others.
- ☐ Time determines the instant in the given body has that location.
- $\hfill \square$ A given body's mass determines how strongly the force of \hfill gravity influences it.
- ☐ Each given body has an additional fundamental dimension like mass, corresponding to each of the forces of nature. Electric charge, for example, dictates how strongly the electrostatic force influences a given body.

1 September 2009

Astronomy 102, Fall 2009

21

20

The fundamental dimensions (continued)

The dimensions of all other physical quantities are combinations of these fundamental dimensions.

- ☐ For instance: the dimension of **velocity**, and velocity's magnitude speed, is distance divided by time, as you
- \Box The dimension of **energy** is mass times distance squared, divided by time squared.
 - $\bullet\;$ i.e. mass times the square of the dimension of speed

☐ Units are the scales of the *quantities* that go with the qualities that are dimensions.

Thus: four fundamental dimensions for location (three space, one time), and in principle four for response to forces (gravity, electricity, and the strong and weak nuclear forces).

1 September 2009

Astronomy 102, Fall 2009

	Speeds in	astronom	y
	cm per second	km per second	miles per hour
NYS Thruway speed limit	3.0×10^{3}	3.0×10 ⁻²	65
Earth's rotational speed at the equator	4.7×10^{4}	0.47	1050
Speed of Earth in orbit	3×10^{6}	30	
Speed of Sun in orbit around center of Milky Way	2.5×10 ⁷	250	
Speed of Milky Way with respect to local Universe	5.5×10 ⁷	550	
Speed of light	2.9979×10^{10}	$2.9979{ imes}10^{5}$	
1 September 2009 Astronom	y 102, Fall 2009		23

Typical speeds and important conversions

- \square Planetary orbits in a solar system: tens of km/s
- $\hfill \square$ Stellar orbits in a normal galaxy: hundreds of km/s
- ☐ Speed between nearby galaxies: hundreds of km/s
- \square Speed of light: 2.99792458×10¹⁰ cm per second
- ☐ Conversion factors: use those given for distance and time.

Example: One mile is equal to 1.61 kilometers. What is the speed of light in miles per hour?

$$c = 2.9979 \times 10^{10} \frac{\text{cm}}{\text{sec}} \times \frac{\text{km}}{10^5 \text{ cm}} \times \frac{\text{mile}}{1.61 \text{ km}} \times \frac{3600 \text{ sec}}{\text{hour}}$$

 $=6.70\times10^8\,\frac{\text{mile}}{\text{c}}$ (670 million miles per hour) Astronomy 102, Fall 2009

1 September 2009

24

(c) University of Rochester

Work, heat and energy in astronomy



Hydrogen atom binding energy 1.6×10^{-12} ergDietary calorie 4.2×10^{10} ergBurn 1 kg anthracite coal 4.3×10^{14} ergDetonate H bomb (1 megaton) 4.2×10^{19} ergEarth-Sun binding energy 5.3×10^{40} ergSun's fuel supply at birth 2×10^{51} ergSupernova (exploding star) 10^{53} erg

1 September 2009

Astronomy 102, Fall 2009

Units of energy

In AST 102 our usual unit of energy will be the erg:

$$1 \text{ erg} = \frac{1 \text{ gram} \times (1 \text{ cm})^2}{(1 \text{ second})^2} = \text{gm cm}^2 \text{ sec}^{-2}$$

which is the unit of energy in the CGS (centimeter-gramsecond) system of units.

Possibly you are more familiar with the International System (SI, a.k.a. MKS for meter-kilogram-second) unit of energy, the **joule**:

1 joule =
$$\frac{1 \text{ kg} \times (1 \text{ m})^2}{(1 \text{ second})^2}$$
 = kg m² sec⁻² = 10⁷ erg

The others we have listed will find some uses too.

1 September 2009

Astronomy 102, Fall 2009

Luminosity (total power output) in astronomy

100 W light bulb

ergs per second (joules per second) (L_{\odot})

1.0×10 $^{\circ}$ 100

 $150\ horsepower\ car\ engine$ 1.2×10^{12} 1.2×10^{5} Large city 10^{15} 10^{8} H bomb (1 megaton, 0.01 4.2×10^{14} 1.1×10⁻¹² 4.2×10^{21} second) 3.8×10^{33} 3.8×10²⁶ Sun 105 4×10^{38} Largest stars 4×10^{31} 2×10^{10} Milky Way galaxy $8{\times}10^{43}$ 3C 273 (a typical quasar) $4{\times}10^{45}$ 10^{12}

For the astronomical objects, the power is emitted mostly in the form of light; hence the name.

1 September 2009

Astronomy 102, Fall 2009

27

25

26

Typical luminosities and important conversions

- \square Normal stars: around one solar luminosity (L_{\odot})
- \Box Giant stars: thousands to hundreds of thousands of L_{\odot}
- \Box Normal galaxies: 10^9 $10^{10}\,L_\odot$
- \blacksquare Quasars: 10^{12} 10^{13} L_{\odot}
- \Box 1 L_{\odot} = 3.8x10 33 erg/s = luminosity of the Sun
- \Box 1 watt = 10^7 erg/s

Example: Vega, the brightest star in the Northern summer sky, has a luminosity of about 1.9x10³⁵ erg/s. What's that in solar luminosities?

$$L = 1.9 \times 10^{35} \text{ erg/s} \times \frac{1 L_{\odot}}{3.8 \times 10^{33} \text{ erg/s}} = 50 L_{\odot}$$

1 September 2009

omy 102 Fall 2009

Rates

Speed and luminosity are examples of rates.

 \square Speed v is the rate of change of position x with time t:

$$v = \frac{\Delta x}{\Delta t}$$
 \Rightarrow $\Delta x = v\Delta t$ if v is constant.

position *interval* time (distance) *interval*

 \Box Luminosity *L* is the rate of change of energy *E* with time *t*:

$$L = \frac{\Delta E}{\Delta t}$$
 \Rightarrow $\Delta E = L\Delta t$ if L is constant.

1 September 2009

Astronomy 102, Fall 2009

Speed as a rate

Example: The radius of the Earth's orbit around the Sun is $1.5x10^{13}$ cm. What is its orbital speed (assumed constant)?

$$v = \frac{\Delta x}{\Delta t} = \frac{2\pi r}{\Delta t} = \frac{2 \times 3.14159 \times (1.5 \times 10^{13} \text{ cm})}{1 \text{ year}} \times \left(\frac{1 \text{ year}}{3.16 \times 10^{7} \text{ seconds}}\right)$$
$$= 3.0 \times 10^{6} \frac{\text{cm}}{\text{sec}} \times \left(\frac{\text{km}}{10^{5} \text{ cm}} \times \frac{1 \text{ mile}}{1.61 \text{ km}} \times \frac{3600 \text{ sec}}{\text{hour}}\right) = 66,800 \text{ mph.}$$

Example: How long should it take to get to Buffalo from here, at the Thruway speed limit?

$$\Delta t = \frac{\Delta x}{v} = \frac{60 \text{ miles}}{65 \frac{\text{miles}}{\text{hour}}} = 0.92 \text{ hour} \times \left(\frac{60 \text{ minutes}}{\text{hour}}\right) = 55 \text{ minutes}.$$

1 September 2009

Astronomy 102, Fall 2009

30



Now you try, with PRSs

There are eight furlongs in a mile, and two weeks in a fortnight. Suppose we take the furlong to be our unit of length, and a fortnight to be our unit of time.

Then, what are the units of speed?

A. Furlong fortnights B. Fortnights per furlong

C. Furlongs per fortnight D. Furlongs per second.

1 September 2009

Astronomy 102, Fall 2009



And again.

There are eight furlongs in a mile, and two weeks in a fortnight. Suppose we take the furlong to be our unit of length, and a fortnight to be our unit of time.

What is the NYS Thruway speed limit in this new system of units?

A. 1.5 furlongs per fortnight C. 8×10⁻⁴ furlong fortnights

B. 1.5×10⁵ furlongs per fortnight

D. 42 fortnights per furlong

1 September 2009

Astronomy 102, Fall 2009

32

Luminosity as a rate

Example: How long could the Sun live at its current luminosity, considering the fuel supply with which it was

$$\Delta t = \frac{\Delta E}{L} = \frac{2 \times 10^{51} \text{ erg}}{3.8 \times 10^{33} \frac{\text{erg}}{\text{sec}}} = 5.3 \times 10^{17} \text{ sec} \times \left(\frac{\text{year}}{3.16 \times 10^7 \text{ sec}}\right)$$

= 1.7×10^{10} years (17 billion years).

It has already lived 4.56 billion years.

Example: What is *your* "luminosity" in erg/sec, if you eat 3000 calories a day and don't gain or lose weight?

$$L = \frac{\Delta E}{\Delta t} = \frac{3000 \text{ Cal}}{1 \text{ day}} \times \left(\frac{1 \text{ day}}{86400 \text{ sec}} \times \frac{4.2 \times 10^{10} \text{ erg}}{\text{Cal}} \right) = 1.5 \times 10^9 \frac{\text{erg}}{\text{sec}}.$$

1 September 2009

Astronomy 102, Fall 2009

33

(c) University of Rochester

Remember the How Bi	ig is i nat sneet
---------------------	-------------------

Many important physical quantities that we will use frequently are collected on the The How Big Is That sheet, found under the "Constants and Equations" tab on the AST 102 Web site.

- ☐ You will always have access to this page while you're doing homework or exams. Thus you don't have to memorize all the numbers.
- ☐ However, to use the sheet effectively, and to understand our astronomical discussions, you must become familiar enough with them to know about how big most of them are
 - It would do you good to memorize at least the "typical" values of things, on the previous pages.

September 2009	Astronomy 102, Fall 2009