

Electrostatic

$$-\nabla^2 \phi = 4\pi\rho \quad \text{with } \vec{E} = -\vec{\nabla}\phi \quad (\text{statics only})$$

physical meaning of the potential ϕ

work done to move a test charge δq from $\vec{r}_1$ to $\vec{r}_2$
in presence of an electric field $\vec{E}$ is

$$W_{12} = \int_{\vec{r}_1}^{\vec{r}_2} d\vec{l} \cdot \vec{F}$$

where $\vec{F}$ is the force required to move the charge.

Since $\vec{E}$ exerts a force $\delta q \vec{E}$ on the charge,

$\vec{F}$ must counterbalance this electric force so

we can move the charge quasi statically $\Rightarrow \vec{F} = -\delta q \vec{E}$

$$W_{12} = -\delta q \int_{\vec{r}_1}^{\vec{r}_2} d\vec{l} \cdot \vec{E} = \delta q \int_{\vec{r}_1}^{\vec{r}_2} d\vec{l} \cdot \vec{\nabla}\phi = \delta q [\phi(\vec{r}_2) - \phi(\vec{r}_1)]$$

$$\phi(\vec{r}_2) - \phi(\vec{r}_1) = \frac{W_{12}}{\delta q}$$

difference in potential between
two points is the work per
unit charge to move a
test charge between the
two points

Green's Functions - part I

$$-\nabla^2 \phi = 4\pi \rho$$

We already know that for a point charge q at position $\vec{r}'$,
ie $\rho(\vec{r}) = q \delta(\vec{r} - \vec{r}')$, the solution to the above is

$$\phi(\vec{r}) = \frac{q}{|\vec{r} - \vec{r}'|} \quad \text{ie} \quad -\nabla^2 \left(\frac{1}{|\vec{r} - \vec{r}'|} \right) = 4\pi \delta(\vec{r} - \vec{r}')$$

We call the special solution for a point source
the Green function for the differential operator

$$-\nabla^2 G(\vec{r}, \vec{r}') = -4\pi \delta(\vec{r} - \vec{r}')$$

$G(\vec{r}, \vec{r}')$ gives the potential at position $\vec{r}$ due
to a unit source at position $\vec{r}'$

Generally, one also has to specify a desired
boundary condition for the Green function on
the boundary of the system.

For the Coulomb solution for a point charge
the implicit boundary condition is that the
potential vanish infinitely far from the charges

$$G(\vec{r}, \vec{r}') \rightarrow 0 \quad \text{as} \quad |\vec{r} - \vec{r}'| \rightarrow \infty$$

boundary of the system is taken to "infinity"

If one knows the Green's function, then one can find the solution for any distribution of sources $\rho(\vec{r})$

$$\phi(\vec{r}) = \int d^3r' G(\vec{r}, \vec{r}') \rho(\vec{r}')$$

proof: $-\nabla^2 \phi = \int d^3r' [\nabla^2 G(\vec{r}, \vec{r}')] \rho(\vec{r}')$

$$= \int d^3r' [4\pi \delta(\vec{r} - \vec{r}')] \rho(\vec{r}')$$

$$= 4\pi \rho(\vec{r})$$

We will return to concept of Green's function when we discuss solution of Poisson's eqn in

We will also see Green's functions again when we discuss solution of the inhomogeneous wave equation.

The Coulomb problem as a boundary value problem

Consider a conducting sphere of radius R with net charge q . (as $R \rightarrow 0$ we get a point charge).

What is $\phi(\vec{r})$? What is $E(\vec{r})$?

Review: Properties of conductors in electrostatics

- 1) $\vec{E} = 0$ inside conductor - if $\vec{E} \neq 0$ then a current $\vec{j} = \sigma \vec{E}$ flows and it is not statics (σ is conductivity)
- 2) $\rho = 0$ inside conductor - if $\vec{E} = 0$ inside, then $\vec{\nabla} \cdot \vec{E} = 4\pi\rho = 0$
- 3) Any net charge on the conductor must lie on the surface - follows from (2)
- 4) $\phi = \text{constant}$ throughout conductor - if $\vec{E} = 0$ then $\vec{E} = -\vec{\nabla}\phi \Rightarrow \phi$ is constant
- 5) Just outside the conductor, $\vec{E}$ is $\perp$ to surface.
- If $\vec{E}$ has a component $\parallel$ to surface then it exerts a force on electrons at the surface leading to a surface current - so would not be static

For conducting sphere, $\rho = 0$ for $r > R$ and $r < R$

all charge is on the surface $\Rightarrow \nabla^2 \phi = 0$ for $\begin{cases} r > R \\ r < R \end{cases}$

spherical symmetry $\Rightarrow$ expect spherically symmetric solution

$\Rightarrow \phi(\vec{r})$ depends only on $r = |\vec{r}|$

⇒ Solve Laplace's eqn by writing ∇^2 in spherical coords.
Only the radial terms do not vanish.

$$\nabla^2 \phi = \frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{d\phi}{dr} \right) = 0$$

$$r^2 \frac{d\phi}{dr} = -C_0 \quad \text{a constant}$$

$$\frac{d\phi}{dr} = -\frac{C_0}{r^2}$$

$$\phi(r) = \frac{C_0}{r} + C_1, \quad C_1 \text{ a constant}$$

"outside" $r > R$ $\phi_{(r)}^{\text{out}} = \frac{C_0^{\text{out}}}{r} + C_1^{\text{out}}$

"inside" $r < R$ $\phi(r) = \frac{C_0^{\text{in}}}{r} + C_1^{\text{in}}$

solution "outside" does not necessarily go smoothly into the solution "inside" because of the charge layer at $r=R$ that separates the two regions. We need to determine the constants $C_0^{\text{in}}, C_0^{\text{out}}, C_1^{\text{in}}, C_1^{\text{out}}$ by applying boundary conditions corresponding to the physical situation.

① For $r > R$, assume $\phi \rightarrow 0$ as $r \rightarrow \infty$ - boundary condition at infinity

$$\Rightarrow C_1^{\text{out}} = 0$$

$$\phi_{(r)}^{\text{out}} = \frac{C_0^{\text{out}}}{r} \quad \text{recover the expected Coulomb form.}$$


2) For $r < R$.

- i) we could use the fact that the region $r < R$ is a conductor with $\phi = \text{constant}$ to conclude $C_0^{\text{in}} = 0$
- ii) or, if we were dealing with a charged shell instead of a conductor, we could argue as follows:
no charge at origin $r=0 \Rightarrow$ expect ϕ should be finite at origin $\Rightarrow C_0^{\text{in}} = 0$

So $\phi^{\text{in}}(r) = C^{\text{in}}$ a constant


3) Now we need boundary condition at $r=R$ where "inside" and "outside" meet.

Review: Electric field and potential at a surface, charge layer



← a general surface S with surface charge density $\sigma(\vec{r})$ for $\vec{r}$ on S . $\sigma(\vec{r})da$ is total charge in area da on surface

i) Take "Gaussian pillbox" surface about point $\vec{r}$ on the surface S



Gauss' Law in integral form $\oint_S da \hat{n} \cdot \vec{E} = 4\pi Q_{\text{enclosed}}$

expect $\vec{E}$ is finite $\rightarrow$ contribution from sides of pillbox vanish as $dl \rightarrow 0$.

$$\oint_S d\vec{a} \cdot \vec{E} = \int_{\text{top}} d\vec{a} \cdot \vec{E} + \int_{\text{bottom}} d\vec{a} \cdot \vec{E}$$

$$= (\hat{n}^{\text{top}} \cdot \vec{E}^{\text{top}} + \hat{n}^{\text{bottom}} \cdot \vec{E}^{\text{bottom}}) da \quad \text{since } da \text{ is small}$$

$\vec{E}^{\text{top}}$ is electric field at $\vec{r}$ just above the surface S
 $\vec{E}^{\text{bottom}}$ is electric field at $\vec{r}$ just below the surface S


$\hat{n}^{\text{top}} \equiv \hat{n}$ is outward normal on top.

$\hat{n}^{\text{bottom}} = -\hat{n}$ is outward normal on bottom

$$\Rightarrow (\vec{E}^{\text{top}} - \vec{E}^{\text{bottom}}) \cdot \hat{n} da = 4\pi Q_{\text{enclosed}} = 4\pi \sigma(\vec{r}) da$$

$$\boxed{(\vec{E}^{\text{top}} - \vec{E}^{\text{bottom}}) \cdot \hat{n} = 4\pi \sigma(\vec{r})} \quad \text{discontinuity in normal component of } \vec{E}$$

ii) Take "Amperian loop" C at surface about point $\vec{r}$.



$$\vec{\nabla} \times \vec{E} = 0 \Rightarrow \oint_C d\vec{\ell} \cdot \vec{E}$$

since $\vec{E}$ is finite at surface,
 if take sides $dl' \rightarrow 0$ their
 contribution to integral vanishes

$$\Rightarrow \oint_C d\vec{\ell} \cdot \vec{E} = (\vec{E}^{\text{top}} - \vec{E}^{\text{bottom}}) \cdot d\vec{\ell} = 0$$

where $d\vec{\ell}$ is any infinitesimal tangent to the surface at $\vec{r}$.

⇒ tangential component of $\vec{E}$ is continuous

combine above to write $\boxed{\vec{E}^{\text{top}} - \vec{E}^{\text{bottom}} = 4\pi\sigma(F) \hat{n}}$

$$\text{iii) } \vec{E} = -\vec{\nabla}\phi \Rightarrow \phi(r_2) - \phi(r_1) = - \int_{r_1}^{r_2} d\vec{l} \cdot \vec{E}$$

Take $\vec{r}_2$ just above $\vec{r}$ on surface
 $\vec{r}_1$ just below $\vec{r}$ on surface $\left. \vphantom{\begin{matrix} \vec{r}_2 \\ \vec{r}_1 \end{matrix}} \right\} d\vec{l} \rightarrow 0$

Since $\vec{E}$ is finite $\Rightarrow \int d\vec{l} \cdot \vec{E} \rightarrow 0$

$$\Rightarrow \boxed{\phi^{\text{top}} = \phi^{\text{bottom}}}$$

potential ϕ is continuous at surface charge layer

can rewrite (i) as

$$(-\vec{\nabla}\phi^{\text{top}} + \vec{\nabla}\phi^{\text{bottom}}) \cdot \hat{n} = 4\pi\sigma$$

$$\boxed{-\frac{\partial\phi^{\text{top}}}{\partial n} + \frac{\partial\phi^{\text{bottom}}}{\partial n} = 4\pi\sigma}$$

↑ directional derivative of ϕ in direction $\hat{n}$

discontinuity in normal derivative of ϕ at surface

Apply to conducting sphere

$$\phi \text{ continuous} \Rightarrow \phi^{\text{in}}(R) = \phi^{\text{out}}(R)$$

$$C_1^{\text{in}} = \frac{C_0^{\text{out}}}{R}$$

only one unknown left

normal derivative of ϕ is discontinuous

$$-\frac{\partial \phi^{\text{top}}}{\partial n} + \frac{\partial \phi^{\text{bottom}}}{\partial n} = 4\pi\sigma$$

here $\hat{n} = \hat{r}$ the radial direction

$$\left[-\frac{d\phi^{\text{out}}}{dr} + \frac{d\phi^{\text{in}}}{dr} \right]_{r=R} = 4\pi\sigma$$

but $\frac{d\phi^{\text{in}}}{dr} = 0$ as $\phi^{\text{in}} = \text{constant}$

$$-\frac{d\phi^{\text{out}}}{dr} \Big|_{r=R} = 4\pi\sigma$$

charge q is uniformly distributed on surface at R

$$-\frac{d}{dr} \left(\frac{C_0^{\text{out}}}{r} \right) \Big|_{r=R} = \frac{C_0^{\text{out}}}{R^2} = 4\pi\sigma = 4\pi \left(\frac{q}{4\pi R^2} \right) = \frac{q}{R^2}$$

$$\Rightarrow C_0^{\text{out}} = q, \quad C^{\text{in}} = \frac{C_0^{\text{out}}}{R} = \frac{q}{R}$$

$$\phi(r) = \begin{cases} \frac{q}{R} & r < R \text{ inside} \\ \frac{q}{r} & r > R \text{ outside} \end{cases}$$

$$\Rightarrow \vec{E} = -\vec{\nabla}\phi = -\frac{d\phi}{dr} = \begin{cases} 0 & r < R \text{ inside} \\ \frac{q}{r^2} & r > R \text{ outside} \end{cases}$$

we get familiar Coulomb solution!

Summary We can view the preceding solution for ϕ_{out} as solving Laplace's eqn $\nabla^2 \phi = 0$ subject to a specified boundary condition on the normal derivative of ϕ at the boundary $r = R$ of the "outside" region of the system.

Alternate problem:

Another physical situation would be to connect conducting sphere to a battery that charges the sphere to a fixed voltage ϕ_0 (stat volts!) with respect to ground $\phi = 0$ at $r \rightarrow \infty$.

As before, outside the sphere $\phi = \frac{C_0}{r}$
Now the boundary condition is to specify the value of ϕ on the boundary of the outside region, i.e.

$$\phi(R) = \phi_0$$

$$\Rightarrow \frac{C_0}{R} = \phi_0, \quad C_0 = \phi_0 R$$

$$\phi(r) = \phi_0 \frac{R}{r}$$

(from preceding solution, we know that charging the sphere to voltage ϕ_0 (statvolts) induces a net charge $q = \phi_0 R$ on it)

These two versions of the conducting sphere problem are examples of a more general boundary value problem

Solve $\nabla^2 \phi = 0$ in a given region of space subject to one of the following two types of boundary conditions on the bounding surfaces of the region

i) Neumann boundary condition

$\frac{\partial \phi}{\partial n}$ - normal derivative of ϕ is specified on the bounding surfaces

ii) Dirichlet boundary condition

ϕ - value of ϕ is specified on the bounding surfaces

If the bounding surfaces consist of disjoint pieces, it is possible to specify either (i) or (ii) on each piece separately to get a mixed boundary value problem.