

$$kd \ll 1 \Rightarrow e^{-ik\hat{r} \cdot \vec{r}'} \approx 1 - ik\hat{r} \cdot \vec{r}' + \text{higher orders}$$

$$\vec{A}_\omega(\vec{r}) = \frac{e^{ikr}}{cr} \int d^3r' \vec{f}_\omega(\vec{r}') (1 - ik\hat{r} \cdot \vec{r}') (1 + \frac{\hat{r} \cdot \vec{r}'}{r})$$

$$= \frac{e^{ikr}}{cr} \int d^3r' \vec{f}_\omega(\vec{r}') \left[ 1 + \hat{r} \cdot \vec{r}' \left( \frac{1}{r} - ik \right) \right]$$

+ higher order in  $\frac{d}{r}$  or  $kd$

$$\vec{A}_\omega(\vec{r}) = \frac{e^{ikr}}{r} \left[ -\vec{I}_1 + \left( \frac{1}{r} - ik \right) \vec{I}_2 \right]$$

where  $\vec{I}_1 = \frac{1}{c} \int d^3r' \vec{f}_\omega(\vec{r}')$

$$\vec{I}_2 = \frac{1}{c} \int d^3r' \hat{r} \cdot \vec{r}' \vec{f}_\omega(\vec{r}')$$

Consider first  $\vec{I}_1$  i<sup>th</sup> component ( $\vec{I}_1$  vanishes in statics)

$$\int d^3r \vec{f}_i(\vec{r}) = - \int d^3r \vec{r}_i \vec{\nabla} \cdot \vec{f} \quad \text{integration by parts}$$

$$= \int d^3r \vec{r}_i \frac{\partial \vec{f}}{\partial t} \quad \text{as } \vec{\nabla} \cdot \vec{f} + \frac{\partial \vec{f}}{\partial t} = 0$$

$$\int d^3r \vec{f}_i(\vec{r}) = -i\omega \int d^3r \vec{r}_i f_\omega(\vec{r})$$

$$\Rightarrow \vec{I}_1 = -\frac{i\omega}{c} \int d^3r \vec{r} f_\omega(\vec{r}) = -\frac{i\omega}{c} \vec{P}_\omega$$

electric dipole moment

## Electric dipole approximation from $\vec{I}_1$

$$\vec{AEI}(\vec{r}) = \frac{e^{ikr}}{r} (-i\omega \vec{p}_\omega) = -i \vec{p}_\omega \frac{k e^{ikr}}{r} \quad \omega = ck$$

Consider  $\vec{I}_2$

$$\vec{I}_2 = \frac{1}{c} \int d^3 r' \hat{r} \cdot \vec{r}' \vec{f}_\omega(\vec{r}') = \frac{1}{c} \hat{r} \cdot \int d^3 r' \vec{r}' \vec{f}_\omega(\vec{r}')$$

we saw this tensor earlier when we did the <sup>tensor</sup>  
magnetic dipole approx, and when we derived the  
macroscopic Maxwell equations

$$\begin{aligned} \int d^3 \vec{r}' \vec{r}' \vec{f}_\omega(\vec{r}') &= - \int d^3 r' \vec{f}_\omega(\vec{r}') \vec{r}' - \int d^3 r' \vec{r}' \vec{r}' (\vec{\nabla}' \vec{f}_\omega(r')) \\ &= \frac{1}{2} \int d^3 r' [\vec{r}' \vec{f}_\omega - \vec{f}_\omega \vec{r}'] - \frac{1}{2} \int d^3 r' \epsilon_{\omega} \vec{r}' \vec{r}' f_\omega \end{aligned}$$

using  $\vec{\nabla}' \vec{f} = -\frac{\partial \vec{f}}{\partial t}$

$$\begin{aligned} \vec{I}_2 &= \frac{1}{2c} \int d^3 r' [(\hat{r} \cdot \vec{r}') \vec{f}_\omega - (\hat{r} \cdot \vec{f}_\omega) \vec{r}'] - \frac{1}{2} \epsilon_{\omega} \hat{r} \cdot \int d^3 r' (\vec{r}' \vec{r}') f_\omega(\vec{r}') \\ &= -\frac{1}{2c} \int d^3 r' [\hat{r} \times (\vec{r}' \times \vec{f}_\omega)] - \frac{1}{2} \frac{\epsilon_{\omega}}{c} \hat{r} \cdot \int d^3 r' (\vec{r}' \vec{r}') f_\omega(\vec{r}') \\ &= -\hat{r} \times \vec{m}_\omega - \frac{1}{2} \frac{i\omega}{3c} \hat{r} \cdot \vec{Q}'_\omega \end{aligned}$$

where  $\vec{m}_\omega = \frac{1}{2c} \int d^3 r' \vec{r}' \times \vec{f}_\omega(\vec{r}')$  is magnetic dipole moment

$$\vec{Q}'_\omega = \int d^3 r' 3\vec{r}' \vec{r}' f_\omega(\vec{r}') \quad \text{looks almost like electric quadrupole tensor}$$

to make it look like the proper quadrupole moment

$$\overleftrightarrow{Q}_w = \int d^3r' (3\vec{r}'\vec{r}' - r'^2 \vec{\mathbb{I}}) P_w(r')$$

we can write

$$\overleftrightarrow{Q}'_w = \overleftrightarrow{Q}_w + \vec{I} \left( \int d^3r' r'^2 f_w(r') \right)$$

$\vec{I}$  identity matrix  $I_{ij} = \delta_{ij}$

$$\vec{I}_2 = -\hat{r} \times \vec{m}_w - \frac{1}{2} \frac{i\omega}{3c} \hat{r} \cdot \overleftrightarrow{Q}_w - \frac{i\omega}{6c} \hat{r} C(w)$$

where  $C_w \equiv \int d^3r' r'^2 f_w(r')$   
is a scalar

Magnetic dipole approximation from  $\vec{I}_2$

$$\vec{A}_{M1}(\vec{r}) = \frac{e^{ikr}}{r} \left( \frac{1}{r} - ik \right) (-\hat{r} \times \vec{m}_w)$$

Electric quadrupole approximation from  $\vec{I}_2$

$$\vec{A}_{E2}(r) = \frac{e^{ikr}}{r} \left( \frac{1}{r} - ik \right) \left( -\frac{i\omega}{6c} \hat{r} \cdot \overleftrightarrow{Q}_w \right)$$

The last piece  $\frac{e^{ikr}}{r} \left( \frac{1}{r} - ik \right) \left( -\frac{i\omega}{6c} \hat{r} C_w \right)$

can always be ignored - it is a radial function  
and so its curl always vanishes  $\rightarrow$  gives

no contribution to  $\vec{B}$ . Similarly, since  $-\frac{i\omega}{c} \vec{E}_w = \vec{k} \times \vec{B}_w$

by Ampere's law, this term will give no  
contribution to  $\vec{E}$ .

So with these two approximations ① and ②

$$\vec{A}_w(\vec{r}) = \vec{A}_{E1}(\vec{r}) + \vec{A}_{M1}(\vec{r}) + \vec{A}_{E2}(\vec{r})$$

keeping higher order terms would give magnetic quadrupole, electric octopole etc.

Compare strengths of the terms above

Approx ③ Radiation zone: for from sources  $\frac{1}{r} \ll k$  so  $(\frac{1}{r} - ik) \approx -ik$  in  $\vec{A}_{M1}$  and  $\vec{A}_{E2}$   
only keep terms of  $O(\frac{1}{r})$

electric dipole  $\vec{P}_w \sim qd$   $\vec{A}_{E1} \sim qkd$

magnetic dipole  $\vec{m}_w \sim vqd$   $\vec{A}_{M1} \sim qhd\left(\frac{v}{c}\right)$

use  $v \sim \frac{d}{t} \sim dw \sim dkc \Rightarrow \vec{A}_{M1} \sim q(kd)^2$

electric quadrupole  $\vec{Q}_w \sim qd^2$   $\vec{A}_{E2} \sim qd^2k\frac{w}{c} \sim q(kd)^2$

Since Approx ② assumed  $kd \approx \frac{v}{c} \ll 1$   
above is expansion in powers of  $kd$

leading term is electric dipole

next order one [magnetic dipole  
electric quadrupole]

$$\frac{A_{M1}}{A_{E1}} \approx \frac{A_{E2}}{A_{E1}} \sim k$$

next order terms are smaller than  $A_{E1}$  by factor  $(kd)^2$

Electric Dipole Approximation - the leading term in non-relativistic expansion

$$\vec{A}_{EI}(\vec{r}) = -ik\vec{p}_w \frac{e^{ikr}}{r}$$

$$\vec{\nabla} \times (\phi \vec{F}) = (\vec{\nabla} \phi) \times \vec{F} + \phi \vec{\nabla} \times \vec{F}$$

$$\vec{B}_{EI} = \vec{\nabla} \times \vec{A}_{EI} = -ik \left( \vec{\nabla} \frac{e^{ikr}}{r} \right) \times \vec{p}_w$$

$$= -ik \left( ik - \frac{1}{r} \right) \frac{e^{ikr}}{r} \hat{r} \times \vec{p}_w$$

$$= k^2 \frac{e^{ikr}}{r} \left( 1 + \frac{i}{kr} \right) \hat{r} \times \vec{p}_w$$

In radiation zone approx.,  $kr, \gg 1$

$$\boxed{\vec{B}_{EI} \approx k^2 \frac{e^{ikr}}{r} \hat{r} \times \vec{p}_w}$$

To get electric field, use Ampere's Law

$$\vec{\nabla} \times \vec{B} = \frac{1}{c} \frac{\partial \vec{E}}{\partial t} \quad (\text{away from source where } \vec{F} = 0)$$

For oscillatory fields  $\vec{E} = E_w e^{-i\omega t}$

$$\vec{\nabla} \times \vec{B}_w = -\frac{c\omega}{c} \vec{E}_w \Rightarrow \vec{E}_{EI} = \frac{i}{k} \vec{\nabla} \times \vec{B}_{EI}$$

$$\vec{E}_{EI} = \frac{i}{k} \vec{\nabla} \times \left[ k^2 \frac{e^{ikr}}{r} \left( 1 + \frac{i}{kr} \right) \hat{r} \times \vec{p}_w \right]$$

$$\vec{E}_{EI} = \frac{i}{k} (\vec{\nabla} e^{ikr}) \times \left[ \frac{k^2}{r} \left(1 + \frac{i}{kr}\right) \hat{r} \times \vec{p}_\omega \right]$$

$$+ \frac{i}{k} e^{ikr} \vec{\nabla} \times \left[ \frac{k^2}{r} \left(1 + \frac{i}{kr}\right) \hat{r} \times \vec{p}_\omega \right]$$

this will always be of order  $1/r^2$   
so can ignore it in radiation zone approx

So in radiation zone approx

$$\vec{E}_{EI} = (\vec{\nabla} e^{ikr}) \times \left[ \frac{ik}{r} \hat{r} \times \vec{p}_\omega \right]$$

$$\boxed{\vec{E}_{EI} = - \frac{k^2}{r} e^{ikr} \hat{r} \times (\hat{r} \times \vec{p}_\omega)}$$

if do not make radiation zone approx, one gets


$$\vec{E}_{EI} = \frac{k^2 e^{ikr}}{r} \left[ \vec{p}_w - \hat{r}(\vec{p}_w \cdot \hat{r}) - \frac{i}{kr} \left( 1 + \frac{i}{kr} \right) (3\hat{r}(\vec{p}_w \cdot \hat{r}) - \vec{p}_w) \right]$$

Using radiation zone approx:

$$\vec{E}_{EI} = k^2 \frac{e^{-ikr}}{r} \hat{r} \times (\vec{p}_w \times \hat{r}) \quad |\vec{E}_{EI}| = |\vec{B}_{EI}|$$

$$\vec{B}_{EI} = -k^2 \frac{e^{-ikr}}{r} \vec{p}_w \times \hat{r} \quad \vec{E}_{EI} + \vec{B}_{EI}$$

If choose coordinates so that  $\vec{p}_w$  is along  $\hat{z}$  axis, then



$$\vec{E}_{EI} = -k^2 p_w \frac{e^{-ikr}}{r} \sin \theta \hat{\theta}$$

$$\vec{B}_{EI} = -k^2 p_w \frac{e^{-ikr}}{r} \sin \theta \hat{\phi}$$

Emitting power

radiating vector  $\vec{S}_{EI}(\vec{r}, t) = \frac{c}{4\pi} \operatorname{Re}\{\vec{E}_{EI}\} \times \operatorname{Re}\{\vec{B}_{EI}\}$

need to take real parts of complex expression  
before multiplying

$$\operatorname{Re}\{\vec{E}_{EI}(\vec{r}, t)\} = -k^2 p_w \frac{\cos(kr - wt)}{r} \sin \theta \hat{\theta}$$

$$\operatorname{Re}\{\vec{B}_{EI}(\vec{r}, t)\} = -k^2 p_w \frac{\cos(kr - wt)}{r} \sin \theta \hat{\phi}$$

$$\boxed{\vec{S}_{EI}(\vec{r}, t) = \frac{c}{4\pi} k^4 p_w^2 \frac{\cos^2(kr - wt)}{r^2} \sin^2 \theta \hat{r}}$$

$\vec{S}_{EI} \sim \hat{r} \Rightarrow$  energy flows radially outwards

$\vec{S}_{EI} \sim \frac{1}{r^2} \Rightarrow$  energy conserved

$$\oint da \hat{m} \cdot \vec{S}_{EI} = \text{constant for all } R$$

sphere  
radius  $R$

time averaged energy current

— Question - what about the  $\frac{1}{r^n}$ ,  $n > 2$ , terms if we do not make radiation zone approx?

$$\langle \vec{S}_{EI} \rangle = \frac{1}{T} \int_0^T dt \vec{S}_{EI}(\vec{r}_t) \quad T \text{ is period } T = \frac{2\pi}{\omega}$$

$$= \frac{C}{8\pi} k^4 p_w^2 \frac{\sin^2 \theta}{r^2} \hat{r}$$

$$\langle \cos^2(\theta) \rangle = \frac{1}{2}$$

average energy flowing through an element of area at spherical angles  $\theta, \phi$  is

$$dP_{EI} = \hat{r} \cdot \underbrace{\langle \vec{S}_{EI} \rangle r^2 \sin \theta d\theta d\phi}_{\text{area of surface element}}$$


$$= r^2 d\Omega \quad \Omega \text{ is solid angle}$$

$$= \hat{r} \cdot \langle \vec{S}_{EI} \rangle r^2 d\Omega$$

$$\frac{dP_{EI}}{d\Omega} = \hat{r} \cdot \langle \vec{S}_{EI} \rangle r^2 = \frac{C}{8\pi} k^4 p_w^2 \sin^2 \theta \sim \omega^4 \sin^2 \theta$$

$\theta = 0$

polar plot



rotationally symmetric  
about  $\hat{z}$  axis

$$\frac{dP_{EI}}{d\Omega} \sim \sin^2 \theta$$

most of power is directed outwards  
into plane  $\perp \vec{P}_{ws}$   
ie peaked about  $\theta = 90^\circ$


Total power radiated is

$$P_{EI} = \int \frac{dP_{EI}}{dS_2} dS_2 = \frac{ck^4 p_w^2}{8\pi} 2\pi \underbrace{\int_0^\pi \sin \theta \sin^2 \theta}_{\frac{4}{3}} \pi$$

$$P_{EI} = \frac{ck^4 p_w^2}{3} = \frac{p_w^2 \omega^4}{3c^3} \sim \omega^4$$

why the sky is blue - Lord Rayleigh

when look up at sky, you are seeing the indirect light of the sun, which is the light emitted by the atoms and molecules of the atmosphere as they oscillate and so radiate, due to the electric field of the direct light from the sun



power of the radiated indirect light is  $P \sim \omega^4 p_w^2$

$$\vec{F} = \alpha \vec{E} \quad \alpha \sim \frac{e^2}{m} \frac{1}{\omega_0^2 - \omega^2 - i\gamma\omega}$$

For molecules in atmosphere ( $N_2$  etc)  $\omega_0$  is typically at a freq higher than visible spectrum. Therefore, in visible spectrum  $\omega \sim \frac{e^2}{m\omega_0^2}$  indep of  $\omega$ .

$\Rightarrow$  power radiated is  $P \sim \omega^4$

$P \sim w^4$  largest at high freq

Since light from sun is "white light"  
it has components of all freqs. Of these  
freqs, the higher ones are scattered the  
most & make up the indirect light we see.

Since blue is the largest  $\omega$  in visible spectrum,  
the sky is blue!

When we look at sunrise or sunset, we  
are looking at the direct rays of the sun.  
Since these rays are least scattered at  
low  $\omega \Rightarrow$  sunset and sunrise are red!

## Magnetic Dipole approx - Radiation Zone $k r \gg 1$

$$\vec{A}_{M1} = \frac{e^{ikr}}{r} \left( \hat{r} - ik \right) \left( -\hat{r} \times \vec{m}_\omega \right)$$

$$\approx ik \hat{r} \times \vec{m}_\omega \frac{e^{ikr}}{r} \quad \text{in RZ}$$

$$\vec{B}_{M1} = \vec{\nabla} \times \vec{A}_{M1} = (\vec{\nabla} e^{ikr}) \times \left( ik \frac{\hat{r} \times \vec{m}_\omega}{r} \right)$$

$$+ e^{ikr} \vec{\nabla} \times \left( \frac{i k \hat{r} \times \vec{m}_\omega}{r} \right)$$

will give terms of  $\mathcal{O}(\frac{1}{r^2})$   
so ignore in RZ approx

$$\boxed{\vec{B}_{M1} = -k^2 \frac{e^{ikr}}{r} \hat{r} \times (\hat{r} \times \vec{m}_\omega)}$$

From Ampere's Law

$$\vec{E}_{M1} = \frac{i}{k} \vec{\nabla} \times \vec{B}_{M1} = -ik \left( \vec{\nabla} e^{ikr} \right) \times \left( \frac{\hat{r} \times [\hat{r} \times \vec{m}_\omega]}{r} \right)$$

$$-ik e^{ikr} \vec{\nabla} \times \left( \frac{\hat{r} \times [\hat{r} \times \vec{m}_\omega]}{r} \right)$$

will give terms of  $\mathcal{O}(\frac{1}{r^2})$   
so ignore in RZ approx

$$\vec{E}_{M1} = k^2 \frac{e^{ikr}}{r} \hat{r} \times (\hat{r} \times (\hat{r} \times \vec{m}_\omega))$$

triple product rule


$$= k^2 \frac{e^{ikr}}{r} \left\{ \hat{r} [\hat{r} \cdot (\hat{r} \times \vec{m}_\omega)] - (\hat{r} \times \vec{m}_\omega) [\hat{r} \cdot \hat{r}] \right\}$$

$$\boxed{\vec{E}_{M1} = -\frac{k^2}{r} e^{ikr} (\hat{r} \times \vec{m}_\omega)}$$

If align axes so that  $\vec{m}_\omega = m_\omega \hat{z}$  then

$$\vec{E}_{M1} = m_\omega \frac{k^2}{r} e^{ikr} \sin\theta \hat{\phi}$$

$$\vec{B}_{M1} = -m_\omega \frac{k^2}{r} e^{ikr} \sin\theta \hat{\theta}$$




Poynting vector

$$\vec{S}_{M1} = \frac{c}{4\pi} \operatorname{Re}\{\vec{E}_{M1}\} \times \operatorname{Re}\{\vec{B}_{M1}\}$$

$$= \frac{c}{4\pi} \frac{k^4 m_\omega^2}{r^2} \cos^2(kr - \omega t) \hat{r}$$

$$\langle \vec{S}_{M1} \rangle = \frac{c}{8\pi} \frac{k^4 m_\omega^2}{r^2} \sin^2\theta \hat{r}$$

$$\frac{dP_{M1}}{dS^2} = \hat{r} \cdot \langle \vec{S}_{M1} \rangle r^2 = \frac{c}{8\pi} k^4 m_\omega^2 \sin^2\theta \sim \omega^4 \sin^2\theta$$



$$P_{M1} = \int dS \frac{dP_{M1}}{dS^2} = \frac{c k^4 m_\omega^2}{3} = \frac{m_\omega^2 \omega^4}{3C^3}$$

$$\frac{P_{M1}}{P_{EI}} = \frac{m_\omega^2}{P_\omega^2} \sim \left(\frac{v}{c}\right)^2 \quad m_\omega \sim \frac{df}{c} \sim dg \frac{v}{c}$$

$$P_\omega \sim dg$$

## Electric Quadrupole radiation - Radiation zone approx

$$\vec{A}_{E2} = \frac{e^{ikr}}{r} (\hat{r} - ik) \left( -\frac{i\omega}{6c} \hat{r} \cdot \overleftrightarrow{Q}_\omega \right)$$

$$= -\frac{e^{ikr}}{r} \frac{k^3}{6} \hat{r} \cdot \overleftrightarrow{Q}_\omega \quad \text{in RZ approx}$$

$$\vec{B}_{E2} = \vec{\nabla} \times \vec{A}_{E2} = -(\vec{\nabla} e^{ikr}) \times \left[ \frac{k^2 \hat{r} \cdot \overleftrightarrow{Q}_\omega}{6r} \right]$$

$$- e^{ikr} \vec{\nabla} \times \left[ \frac{k^2 \hat{r} \cdot \overleftrightarrow{Q}_\omega}{6r} \right]$$

$\mathcal{O}(\frac{1}{r^2})$  so ignore in RZ approx

$$\boxed{\vec{B}_{E2} = -ik^3 \frac{e^{ikr}}{6r} \hat{r} \times (\hat{r} \cdot \overleftrightarrow{Q}_\omega)}$$

$$\vec{E}_{E2} = \frac{i}{k} \vec{\nabla} \times \vec{B}_{E2} = k^2 (\vec{\nabla} e^{ikr}) \times \left[ \frac{\hat{r} \times (\hat{r} \cdot \overleftrightarrow{Q}_\omega)}{6r} \right]$$

$$+ k^2 e^{ikr} \vec{\nabla} \times \left[ \frac{\hat{r} \times (\hat{r} \cdot \overleftrightarrow{Q}_\omega)}{6r} \right]$$

$\mathcal{O}(\frac{1}{r^2})$  so ignore in RZ approx

$$\vec{E}_{E2} = ik^3 \frac{e^{ikr}}{6r} \hat{r} \times \left[ \hat{r} \times (\hat{r} \cdot \overleftrightarrow{Q}_\omega) \right]$$

triple product rule

$$= \frac{ik^3 e^{ikr}}{6r} \left\{ \hat{r} [\hat{r} \cdot (\hat{r} \cdot \overleftrightarrow{Q}_\omega)] - (\hat{r} \cdot \overleftrightarrow{Q}_\omega) [\hat{r} \cdot \hat{r}] \right\}$$

$$\boxed{\vec{E}_{E2} = \frac{ik^3 e^{ikr}}{6r} \left\{ \hat{r} (\hat{r} \cdot \overleftrightarrow{Q}_\omega \cdot \hat{r}) - (\hat{r} \cdot \overleftrightarrow{Q}_\omega) \right\}}$$

## Poynting vector

$$\vec{S}_{E2} = +\frac{c}{4\pi} \operatorname{Re} \{ \vec{E}_{E2} \} \times \operatorname{Re} \{ \vec{B}_{E2} \}$$

$$= -\frac{(k\pi)^6}{36r^2} \sin^2(kr-wt) \left\{ \hat{r} (\hat{r} \cdot \vec{\mathbb{Q}}_w \cdot \hat{r}) - (\hat{r} \cdot \vec{\mathbb{Q}}_w) \right\} \times \left[ \hat{r} \times (\hat{r} \cdot \vec{\mathbb{Q}}_w) \right]$$

$$= -\frac{(k\pi)^6}{36r^2} \sin^2(kr-wt) \left\{ \hat{r} [\hat{r} (\hat{r} \cdot \vec{\mathbb{Q}}_w \cdot \hat{r}) \cdot (\hat{r} \cdot \vec{\mathbb{Q}}_w)] - (\hat{r} \cdot \vec{\mathbb{Q}}_w) [\hat{r} (\hat{r} \cdot \vec{\mathbb{Q}}_w \cdot \hat{r}) \cdot \hat{r}] \right.$$

$$\left. - \hat{r} [(\hat{r} \cdot \vec{\mathbb{Q}}_w) \cdot (\hat{r} \cdot \vec{\mathbb{Q}}_w)] + (\hat{r} \cdot \vec{\mathbb{Q}}_w) [(\hat{r} \cdot \vec{\mathbb{Q}}_w) \cdot \hat{r}] \right\}$$

$$= -\frac{(k\pi)^6}{36r^2} \sin^2(kr-wt) \left\{ (\hat{r} \cdot \vec{\mathbb{Q}}_w \cdot \hat{r})^2 \hat{r} - (\hat{r} \cdot \vec{\mathbb{Q}}_w) (\hat{r} \cdot \vec{\mathbb{Q}}_w \cdot \hat{r}) \right. \\ \left. - (\hat{r} \cdot \vec{\mathbb{Q}}_w \cdot \vec{\mathbb{Q}}_w \cdot \hat{r}) \hat{r} (\hat{r} \cdot \vec{\mathbb{Q}}_w) (\hat{r} \cdot \vec{\mathbb{Q}}_w \cdot \hat{r}) \right\}$$

$$\vec{S}_{E2} = \frac{-ck^6}{4\pi 36r^2} \sin^2(kr-wt) \left\{ (\hat{r} \cdot \vec{\mathbb{Q}}_w \cdot \hat{r})^2 - (\hat{r} \cdot \vec{\mathbb{Q}}_w)^2 \right\} \hat{r}$$

$$\langle \vec{S}_{E2} \rangle = -\frac{ck^6}{4\pi 72r^2} \left\{ (\hat{r} \cdot \vec{\mathbb{Q}}_w \cdot \hat{r})^2 - (\hat{r} \cdot \vec{\mathbb{Q}}_w)^2 \right\} \hat{r}$$

$$\frac{dP_{E2}}{ds} = \hat{r} \cdot \langle \vec{S}_{E2} \rangle r^2 = \frac{-ck^6}{4\pi 72} \left\{ (\hat{r} \cdot \vec{\mathbb{Q}}_w)^2 - (\hat{r} \cdot \vec{\mathbb{Q}}_w \cdot \hat{r})^2 \right\}$$


angular dependence of  $\frac{dP_{E2}}{ds}$  depends  
on specific form of the tensor  $\vec{\mathbb{Q}}_w$

For example: suppose  $\Omega_{ij} = 0$  except for  $\Omega_{zz}$

$$\Rightarrow \vec{\Omega}_\omega = \Omega_{zz} \hat{z} \hat{z}$$

$$(\hat{r} \cdot \vec{\Omega}_\omega \cdot \hat{r})^2 = (\Omega_{zz} \cos^2 \theta)^2$$

$$(\hat{r} \cdot \vec{\Omega}_\omega)^2 = \Omega_{zz}^2 \cos^2 \theta$$




$$\frac{dP_{E2}}{d\Omega} = \frac{ck^6}{4\pi f_2} \Omega_{zz}^2 [\cos^2 \theta - \cos^4 \theta]$$

$$= \frac{ck^6}{4\pi f_2} \Omega_{zz}^2 \cos^2 \theta \sin^2 \theta$$

$$\cos \theta \sin \theta = \frac{1}{2} \sin 2\theta$$

$$= \frac{ck^6}{4\pi f_2 288} \Omega_z^{22} \sin^2 2\theta$$



peak at  $45^\circ$

$\theta = 90^\circ$  rotationally invariant about  $\hat{z}$  axis

$$\frac{P_{E2}}{P_{E1}} \sim \frac{k^6 \Omega^2}{k^4 p^2} \sim \frac{k^2 (qd^2)^2}{(qd)^2} \sim k^2 d^2 \sim \left(\frac{v}{c}\right)^2$$

$$P_{E2} \sim P_{M1}$$

For more general case, choose axes so that  $\vec{Q}_w$  is diagonal - can always do this since  $\vec{Q}_w$  is symmetric

$$(\hat{r} \cdot \vec{Q}_w \cdot \hat{r}) = \hat{r} \cdot \begin{pmatrix} Q_{xx} & 0 & 0 \\ 0 & Q_{yy} & 0 \\ 0 & 0 & Q_{zz} \end{pmatrix} \cdot \hat{r}$$

$$= \hat{r} \cdot \begin{pmatrix} Q_{xx} \sin\theta \cos\varphi \\ Q_{yy} \sin\theta \sin\varphi \\ Q_{zz} \cos\theta \end{pmatrix} = Q_{xx} \sin^2\theta \cos^2\varphi + Q_{yy} \sin^2\theta \sin^2\varphi + Q_{zz} \cos^2\theta$$

$$(\hat{r} \cdot \vec{Q}_w)^2 = Q_{xx}^2 \sin^2\theta \cos^2\varphi + Q_{yy}^2 \sin^2\theta \sin^2\varphi + Q_{zz}^2 \cos^2\theta$$

$$\frac{dP_{E2}}{d\Omega} = \frac{k^6}{f^2} \left\{ Q_{zz}^2 (\cos^2\theta - \cos^4\theta) + Q_{xx}^2 (\sin^2\theta \cos^2\varphi - \sin^4\theta \cos^4\varphi) + Q_{yy}^2 (\sin^2\theta \sin^2\varphi - \sin^4\theta \sin^4\varphi) \right\}$$

$$\frac{dP_{E2}}{d\Omega} = \frac{k^6}{f^2} \left\{ Q_{zz}^2 \cos^2\theta \sin^2\theta + Q_{xx}^2 \sin^2\theta \cos^2\theta (1 - \sin^2\theta \cos^2\varphi) + Q_{yy}^2 \sin^2\theta \sin^2\theta (1 - \sin^2\theta \sin^2\varphi) \right\}$$

no special symmetries - varies with  $\theta$  and  $\varphi$