

Electromagnetism


Clearly \vec{E} + \vec{B} must transform into each other under Lorentz transf.

in inertial frame K
stationary line charg λ

$$\vec{E} \leftarrow \uparrow \nearrow \downarrow$$

$\swarrow \searrow$
cylindrical outward
electric field
no B -field

in frame K' moving with $\vec{v} \parallel$ to wire



moving line charge gives current
 $\Rightarrow B$ circulating around wire
as well as outward radial E

Lorentz force

$$\vec{F} = q\vec{E} + q\frac{\vec{v}}{c} \times \vec{B}$$

What is the velocity \vec{v} here? velocity with respect to what inertial frame? Clearly \vec{E} and \vec{B} must change from one inertial frame to another if this force law can make sense.

charge density

Consider charge ΔQ contained in a vol ΔV .

ΔQ is a Lorentz invariant scalar.

Consider the reference frame in which the charge is instantaneously at rest. In this frame

$$\Delta Q = \hat{\rho} \Delta V$$

$\hat{\rho}$ is charge density in the rest frame
 ΔV is volume in the rest frame

$\hat{\rho}$ is Lorentz invariant by definition

Now transform to another frame moving with \vec{v} with respect to rest frame

ΔQ remains the same

$$\Delta V = \frac{\hat{\Delta V}}{\gamma}$$

volume contracts in direction II to \vec{v}

$$\rho = \frac{\Delta Q}{\Delta V} = \frac{\Delta Q}{\hat{\Delta V}} \gamma = \hat{\rho} \gamma$$

$$\text{Current density is } \vec{j} = \rho \vec{v} = \gamma \hat{\vec{v}} \cdot \frac{\rho}{\gamma} = \hat{\rho} \hat{\vec{v}}$$

Define 4-current $j^\mu = (\vec{j}, i c \rho) = \hat{\rho} (\hat{\vec{v}}, i c \gamma)$

$$= \hat{\rho} u^\mu$$

j^μ is 4-vector since u^μ is 4-vector and $\hat{\rho}$ is Lorentz invariant scalar.

charge conservation

$$\vec{\nabla} \cdot \vec{j} + \frac{\partial \rho}{\partial t} = \boxed{\frac{\partial j^\mu}{\partial x_\mu} = 0}$$

Equation for potentials in Lorentz gauge

$$(\nabla^2 - \frac{1}{c^2} \frac{\partial^2}{\partial t^2}) \vec{A} = -\frac{4\pi}{c} \vec{f}$$

$$(\nabla^2 - \frac{1}{c^2} \frac{\partial^2}{\partial t^2}) \phi = -4\pi f$$

$$\frac{\partial^2}{\partial x_\mu^2} = (\nabla^2 - \frac{1}{c^2} \frac{\partial^2}{\partial t^2}) \text{ is Lorentz invariant operator}$$

4-potential

$$A_\mu = (\vec{A}, i\phi)$$

$$(\nabla^2 - \frac{1}{c^2} \frac{\partial^2}{\partial t^2}) A_\mu = -\frac{4\pi}{c} f_\mu = \frac{\partial^2 A_\mu}{\partial x_\mu^2}$$

Lorentz gauge condition is

$$\vec{\nabla} \cdot \vec{A} + \frac{\partial \phi}{c \partial t} = \frac{\partial A_\mu}{\partial x^\mu} = 0$$

Electric and magnetic fields

$$B_i = \frac{\partial A_k}{\partial x_j} - \frac{\partial A_j}{\partial x_k} \quad i, j, k \text{ cyclic permutation of } 1, 2, 3$$

$$E_i = -\frac{\partial \phi}{\partial x_i} - \frac{\partial A_i}{c \partial t} = i \left(\frac{\partial A_4}{\partial x_i} - \frac{\partial A_i}{\partial x_4} \right)$$

Define field stress tensor

$$F_{\mu\nu} = \frac{\partial A_\nu}{\partial x_\mu} - \frac{\partial A_\mu}{\partial x_\nu} = -F_{\nu\mu}$$

$$= \begin{pmatrix} 0 & B_3 & -B_2 & -iE_1 \\ -B_3 & 0 & B_1 & -iE_2 \\ B_2 & -B_1 & 0 & -iE_3 \end{pmatrix}$$

"curl" of a 4-ve
is a 4×4 anti
symmetric 2nd rank ten

In homogeneous Maxwell's equations can be written
in the form

$$\boxed{\frac{\partial F_{\mu\nu}}{\partial x_\nu} = \frac{4\pi}{c} j_\mu} \Rightarrow \boxed{\vec{\nabla} \cdot \vec{E} = 4\pi \rho \\ \vec{\nabla} \times \vec{B} - \frac{1}{c} \frac{\partial \vec{E}}{\partial t} = \frac{4\pi}{c} \vec{j}}$$

$$= \frac{\partial}{\partial x_\nu} \left(\frac{\partial A_\nu}{\partial x_\mu} - \frac{\partial A_\mu}{\partial x_\nu} \right) = \frac{\partial}{\partial x_\mu} \left(\frac{\partial A_\nu}{\partial x_\nu} \right) - \frac{\partial^2 A_\mu}{\partial x_\nu^2}$$

"0"

$$\Rightarrow - \frac{\partial^2 A_\mu}{\partial x_\nu^2} = \frac{4\pi}{c} j_\mu \text{ agrees with previous equation for } A_\mu$$

transformation law for 2nd rank tensor $F_{\mu\nu}$

$$\begin{aligned} F'_{\mu\nu} &= \frac{\partial A'_\nu}{\partial x^\mu} - \frac{\partial A'_\mu}{\partial x^\nu} \quad \text{use } A'_\mu = \alpha_{\mu 0} A_0 \\ &= \alpha_{\nu 2} \alpha_{\mu 0} \frac{\partial A_\lambda}{\partial x^\sigma} \quad \frac{\partial}{\partial x^\mu} = \alpha_{\mu 0} \lambda \frac{\partial}{\partial x_\lambda} \\ &\quad - \alpha_{\mu 0} \alpha_{\nu 2} \frac{\partial A_0}{\partial x_\lambda} \end{aligned}$$

$$F'_{\mu\nu} = \alpha_{\mu 0} \alpha_{\nu 2} F_{0\lambda} \quad \left. \begin{array}{l} \text{lets one find } \vec{E}' \text{ and } \vec{B}' \\ \text{if one knows } \vec{E} \text{ and } \vec{B} \end{array} \right\}$$

for n th rank tensor

$$T'_{\mu_1 \mu_2 \dots \mu_n} = \alpha_{\mu_1 0} \alpha_{\mu_2 2} \dots \alpha_{\mu_n n} T_{0, 2, \dots, n}$$

$\frac{\partial F_{\mu\nu}}{\partial x^\nu}$ is a 4-vector: proof:

$$\frac{\partial F_{\mu\nu}'}{\partial x'_\nu} = \alpha_{\mu\sigma} \alpha_{\nu\lambda} \alpha_{\lambda\tau} \frac{\partial F_{\sigma\lambda}}{\partial x_\tau}$$

but $\alpha_{\nu\lambda} = \bar{\alpha}_{\lambda\nu}$ since inverse = transpose

$$\alpha_{\nu\lambda} \alpha_{\lambda\tau} = \bar{\alpha}_{\lambda\nu} \alpha_{\lambda\tau} = \delta_{\nu\tau}$$

$$\frac{\partial F_{\mu\nu}'}{\partial x'_\nu} = \alpha_{\mu\sigma} \frac{\partial F_{\sigma\lambda}}{\partial x_\lambda} \delta_{\nu\lambda} = \alpha_{\mu\sigma} \frac{\partial F_{\sigma\lambda}}{\partial x_\lambda} \text{ transforms like 4-vector}$$

To write the homogeneous Maxwell Equations

Construct 3rd rank co-variant tensor

$$G_{\mu\nu\lambda} = \frac{\partial F_{\mu\nu}}{\partial x_\lambda} + \frac{\partial F_{\nu\mu}}{\partial x_\lambda} + \frac{\partial F_{\lambda\mu}}{\partial x_\nu}$$

transforms as $G'_{\mu\nu\lambda} = \alpha_{\mu\alpha} \alpha_{\nu\beta} \alpha_{\lambda\gamma} G_{\alpha\beta\gamma}$

In principle G has $4^3 = 64$ components

But can show that G is antisymmetric in exchange of any two indices

$$G_{\nu\mu\lambda} = \frac{\partial F_{\nu\mu}}{\partial x_\lambda} + \frac{\partial F_{\lambda\nu}}{\partial x_\mu} + \frac{\partial F_{\mu\lambda}}{\partial x_\nu}$$

$$= -\frac{\partial F_{\mu\nu}}{\partial x_\lambda} - \frac{\partial F_{\nu\lambda}}{\partial x_\mu} - \frac{\partial F_{\lambda\mu}}{\partial x_\nu} \quad \text{as } F \text{ anti-symmetric}$$

$$\therefore -G_{\mu\nu\lambda}$$

Also $G_{\mu\nu\lambda} = 0$ if any two indices are equal

\Rightarrow only 4 independent components

$$G_{012}, G_{013}, G_{023}, G_{123}$$

all other components either vanish or are \pm one of the above.

the 4 homogeneous Maxwell Equations:

$$\vec{\nabla} \cdot \vec{B} = 0, \quad \vec{\nabla} \times \vec{E} + \frac{1}{c} \frac{\partial \vec{B}}{\partial t} = 0$$

can be written as

$$\boxed{G_{\mu\nu\lambda} = 0}$$

to see, substitute in definition of G the definition of F

$$G_{\mu\nu\lambda} = \underbrace{\frac{\partial^2 A_\nu}{\partial x_\lambda \partial x_\mu} - \frac{\partial^2 A_\mu}{\partial x_\lambda \partial x_\nu}}_{+} + \underbrace{\frac{\partial^2 A_\mu}{\partial x_\nu \partial x_\lambda} - \frac{\partial^2 A_\lambda}{\partial x_\nu \partial x_\mu}}_{+} + \underbrace{\frac{\partial^2 A_\lambda}{\partial x_\mu \partial x_\nu} - \frac{\partial^2 A_\nu}{\partial x_\mu \partial x_\lambda}}_{+}$$

all terms cancel in pairs

$$= 0$$

$$G_{123} = 0 \Rightarrow \vec{\nabla} \cdot \vec{B} = 0$$

$$G_{012} = -i \left[\vec{\nabla} \times \vec{E} + \frac{1}{c} \frac{\partial \vec{B}}{\partial t} \right]_z = 0 \quad \text{3 component Faraday's law}$$

Another way to write homogeneous Maxwell Equations

Define $\epsilon_{\mu\nu\rho\sigma} = \begin{cases} +1 & \text{if } \mu\nu\rho\sigma \text{ is even permutation} \\ & \text{of } 1234 \\ -1 & \text{if } \mu\nu\rho\sigma \text{ is odd permutation} \\ & \text{of } 1234 \\ 0 & \text{otherwise} \end{cases}$

Define $\tilde{F}_{\mu\nu} = \frac{1}{2!} \epsilon_{\mu\nu\rho\sigma} F_{\rho\sigma}$ pseudo-tensor
 $= \begin{pmatrix} 0 & -E_3 & E_2 & -iB_1 \\ E_3 & 0 & -E_1 & -iB_2 \\ -E_2 & E_1 & 0 & -iB_3 \\ iB_1 & iB_2 & iB_3 & 0 \end{pmatrix}$ has wrong sign
under parity
transf

$\partial_\nu \tilde{F}_{\mu\nu} = 0$ gives homogeneous Maxwell equations

$$\frac{1}{2} F_{\mu\nu} F_{\mu\nu} = B^2 - E^2 \quad \left. \right\} \text{Lorentz invariant scalars}$$

$$-\frac{1}{4} F_{\mu\nu} \tilde{F}^{\mu\nu} = \vec{B} \cdot \vec{E} \quad \left. \right\}$$

From $F_{\mu\nu}^1 = \partial_\mu \partial_\nu F_0$, we can get
velocity transf for \vec{E} and \vec{B}

For a transformation from K to K' with K' moving
with v along x , with respect to K ,

$$E'_1 = E_1$$

$$B'_1 = B_1$$

$$E'_2 = \gamma(E_2 - \frac{v}{c} B_3)$$

$$B'_2 = \gamma(B_2 + \frac{v}{c} E_3)$$

$$E'_3 = \gamma(E_3 + \frac{v}{c} B_2)$$

$$B'_3 = \gamma(B_3 - \frac{v}{c} E_2)$$

Kinematics

"dot" is $\frac{d}{ds}$



4-momentum $p_\mu = m \dot{x}_\mu = m u_\mu = (m \gamma \vec{v}, \pm mc\gamma)$

$$p_\mu^2 = m^2 u_\mu^2 = -m^2 c^2$$

4-force $K_\mu = (\vec{K}, iK_0)$ "Minkowski force"

Newton's 2nd law

$$m \frac{d^2 x_\mu}{ds^2} = K_\mu$$

$$\Rightarrow m \frac{d u_\mu}{ds} = \frac{d p_\mu}{ds} = K_\mu$$

$$p_\mu^2 = -m^2 c^2 \Rightarrow \frac{d}{ds} (p_\mu^2) = p_\mu \frac{d p_\mu}{ds} = p_\mu K_\mu = 0$$

$$\Rightarrow m \gamma \vec{v} \cdot \vec{K} - mc\gamma K_0 = 0 \quad \text{or}$$

$$K_0 = \frac{\vec{v}}{c} \cdot \vec{K}$$

Define the usual 3-force by

$$\frac{d\vec{p}}{dt} = \vec{F}$$

$$\frac{d\vec{p}}{ds} = \vec{k} \text{ and } \frac{d\vec{p}}{ds} = \gamma \frac{d\vec{p}}{dt} = \gamma \vec{F} \Rightarrow \vec{k} = \gamma \vec{F}$$

$$K_0 = \gamma \frac{\vec{v}}{c} \cdot \vec{F}$$

Consider 4th component of Newton's eqn

$$\frac{m}{ds} u_4 = \frac{m}{ds} (ic\gamma) = iK_0 = i\gamma \frac{\vec{v}}{c} \cdot \vec{F}$$

$$d(m\gamma) = \gamma \frac{\vec{v}}{c^2} \cdot \vec{F} ds = dt \frac{\vec{v} \cdot \vec{F}}{c^2} = d\vec{r} \cdot \vec{F}$$

Work-energy theorem: $d(m\gamma c^2) = d\vec{r} \cdot \vec{F} = \text{work done}$

$\Rightarrow d(m\gamma c^2)$ is change in kinetic energy

$E = m\gamma c^2$ is relativistic kinetic energy

$\not{p}_\mu = (\vec{p}, \frac{iE}{c})$	$\vec{p} = m\gamma \vec{v}$
	$E = m\gamma c^2$

$$E = \frac{mc^2}{\sqrt{1-\frac{v^2}{c^2}}} \approx mc^2 \left(1 + \frac{1}{2} \frac{v^2}{c^2}\right) = mc^2 + \frac{1}{2} \frac{mv^2}{c^2}$$

↑ ↑

small v non-rel

$$\frac{dp_\mu}{ds} = k_\mu \rightarrow \text{therefore}$$

relativistic analog of Newton's 3rd law
as well as law of conservation of energy

rest mass kinetic energy

Lorentz force

$$\frac{dp_\mu}{ds} = K_\mu$$

what is the K_μ that represents the Lorentz force
and how can we write it in ~~relative~~ Lorentz
covariant way?

K_μ should depend on the fields $F_{\mu\nu}$
and the particles trajectory x_μ

$$\text{as } \vec{r} \rightarrow 0 \quad \vec{K} = q \vec{E}$$

K_μ can't depend directly on x_μ as should be
indep of origin of coords. So can
depend only on \dot{x}_μ , \ddot{x}_μ , etc.

as $v \rightarrow 0$, K does not depend on the
acceleration, so K does not depend on \dot{x}_μ

K_μ only depends on $F_{\mu\nu}$ and \dot{x}_μ

we need to form a 4-vector out of
 $F_{\mu\nu}$ and \dot{x}_μ that is linear in the fields $F_{\mu\nu}$
and proportional to the charge q .

The only possibility is

$$q f(x_\mu^2) F_{\mu\nu} \dot{x}_\nu$$

But $\dot{x}_\mu^2 = c^2$ is a constant. Choose $f(x_\mu^2) = \frac{1}{c}$

$K_\mu = \frac{g}{c} F_{\mu\nu} \dot{x}_\nu$ is only possibility

This gives force

$$\vec{F} = \frac{1}{c} \vec{K}$$

$$F_i = \frac{1}{c} K_i = \frac{g}{c} (F_{ij} \dot{x}_j + F_{i4} \dot{x}_4)$$

$$= \frac{g}{c} \left(\frac{\partial A_j - \partial A_i}{\partial x_j} \right) \dot{x}_j + \frac{g}{c} (-iE_i)(ic)$$

$$= \frac{g}{c} [\epsilon_{ijk} B_k \gamma_0 j] + \frac{g}{c} E_i c \gamma$$

$$= g E_i + g \epsilon_{ijk} \frac{v_j}{c} B_k$$

$$\vec{F} = g \vec{E} + g \frac{\vec{v}}{c} \times \vec{B}$$

Lorentz force is the same form in all inertial frames.
No relativistic modification is needed.

Relativistic Larmor's formula

$$\text{non-relativistic } P = \frac{2}{3} \frac{e^2 [\text{alt o}]}{\epsilon^3} r^2$$

Consider inertial frame in which charge is instantaneously at rest. Call this rest frame K.

$$\text{power radiated in } K \text{ is } P = \frac{d\overset{\circ}{E}(t)}{dt}$$

where $\overset{\circ}{E}$ is energy radiated. In K , the momentum density $\overset{\circ}{T} = \frac{1}{4\pi c} \overset{\circ}{E} \times \overset{\circ}{B} \sim \overset{\circ}{F}$ is in outward radial direction. Integrating over all directions, the radiated momentum vanishes

$$\overset{\circ}{P} = 0$$

energy-momentum is a 4-vector $(\overset{\circ}{P}, \frac{1}{c}\overset{\circ}{E})$

To get radiated energy in original frame K we can use Lorentz transf

$$\frac{\overset{\circ}{E}}{c} = \gamma \left(\frac{\overset{\circ}{E}}{c} - \frac{\overset{\circ}{v}}{c} \cdot \overset{\circ}{P} \right) \Rightarrow \overset{\circ}{E} = \gamma \overset{\circ}{E} \text{ as } \overset{\circ}{P} = 0$$

and $dt = \gamma dt'$ is time interval in K

$(dt' = 0 \text{ as charge stays at origin in K})$

$$\text{So } \frac{d\overset{\circ}{E}}{dt} = \frac{\gamma d\overset{\circ}{E}}{\gamma dt'} = \frac{d\overset{\circ}{E}}{dt} \Rightarrow P = \overset{\circ}{P}$$

radiated power \propto Lorentz moment!

in \hat{K} we can use non-relativistic Larmor's formula since $v=0$. So

$$P = \frac{2}{3} \frac{8 \hat{a}^2}{c^3}$$

\hat{a} = acceleration in \hat{K}

To write an expression without explicitly making mention of frame \hat{K} , we need to find a Lorentz invariant scalar that reduces to a^2 as $v \rightarrow 0$.

Only choice is α_μ^2 the 4-acceleration $\alpha_\mu = \frac{du_\mu}{ds}$

$$\alpha_\mu = \frac{du_\mu}{ds} = \gamma \frac{du_\mu}{dt} = \gamma \frac{d}{dt} (\gamma \vec{v}, cc\gamma)$$

$$\vec{\alpha} = \gamma^2 \frac{d\vec{v}}{dt} + \gamma \vec{v} \frac{d\gamma}{dt}$$

$$\alpha_4 = cc\gamma \frac{d\gamma}{dt}$$

$$\frac{d\gamma}{dt} = \frac{d}{dt} \left(\frac{1}{\sqrt{1-v^2/c^2}} \right) = \frac{\vec{v} \cdot \frac{d\vec{v}}{dt}}{(1-v^2/c^2)^{3/2}} = \frac{1}{c^2} \gamma^3 \vec{v} \cdot \vec{a}$$

as $\vec{v} \rightarrow 0$, $\gamma \rightarrow 1$, $\frac{d\gamma}{dt} \rightarrow 0$, so $\left\{ \vec{\alpha} \rightarrow \frac{d\vec{v}}{dt} = \vec{a} \right.$

$$\alpha_\mu^2 \rightarrow |\vec{a}|^2 \text{ as desired}$$

$$\left. \alpha_4 \rightarrow 0 \right\}$$

Relativistic Larmor's formula

$$P = \frac{2}{3} \frac{8}{c^3} \alpha_\mu^2 = \frac{2}{3} \frac{8}{c^3} (\alpha_\mu)^2$$

$$\alpha_\mu = \left(\gamma^2 \frac{d\vec{v}}{dt} + \gamma \vec{v} \frac{d\gamma}{dt}, -c\gamma \frac{d\gamma}{dt} \right)$$

$$\frac{d\gamma}{dt} = \frac{1}{c^2} \gamma^3 \vec{v} \cdot \vec{a}$$

$$\alpha_\mu = \left(\gamma^2 \vec{a} + \gamma^4 \frac{1}{c^2} (\vec{v} \cdot \vec{a}) \vec{v}, -c \frac{\gamma^4}{c^2} \vec{v} \cdot \vec{a} \right)$$

$$\begin{aligned}\alpha_\mu^2 &= \gamma^4 a^2 + \gamma^8 \frac{(\vec{v} \cdot \vec{a})^2 v^2}{c^4} + \frac{2\gamma^6}{c^2} (\vec{v} \cdot \vec{a})^2 - \frac{\gamma^8 (\vec{v} \cdot \vec{a})^2}{c^2} \\ &= \gamma^4 \left[a^2 + \gamma^4 \frac{(\vec{v} \cdot \vec{a})^2}{c^2} \left(\frac{v^2}{c^2} - 1 \right) + \frac{2\gamma^2}{c^2} (\vec{v} \cdot \vec{a})^2 \right] \\ &= \gamma^4 \left[a^2 - \gamma^2 \frac{(\vec{v} \cdot \vec{a})^2}{c^2} + \frac{2\gamma^2}{c^2} (\vec{v} \cdot \vec{a})^2 \right]\end{aligned}$$

$$\alpha_\mu^2 = \gamma^4 \left[a^2 + \gamma^2 \frac{(\vec{v} \cdot \vec{a})^2}{c^2} \right]$$

$$\text{as } \vec{v} \rightarrow 0 \rightarrow \alpha_\mu^2 \rightarrow a^2$$

$\alpha_\mu^2 = a^2$ Lorentz invariant

a = acceleration in instantaneous rest

For a charge accelerating in linear motion, $\alpha_\mu^2 = v^2 a^2$

$$\alpha_\mu^2 = \gamma^4 a^2 \left(1 + \gamma^2 \frac{v^2}{c^2} \right) = \gamma^6 a^2$$

$$P = \frac{2}{3} \frac{a^2}{c^3} \gamma^6$$

For a charge in circular motion $(\vec{v} \cdot \vec{a}) = 0$

$$\alpha_\mu^2 = \gamma^4 a^2$$

$$P = \frac{2}{3} \frac{a^2}{c^3} \gamma^4$$