

Green's Functions - part I

$$-\nabla^2 \phi = 4\pi \rho$$

We already know that for a point charge q at position $\vec{r}'$,
ie $\rho(\vec{r}) = q \delta(\vec{r} - \vec{r}')$, the solution to the above is

$$\phi(\vec{r}) = \frac{q}{|\vec{r} - \vec{r}'|} \quad \text{ie} \quad -\nabla^2 \left(\frac{1}{|\vec{r} - \vec{r}'|} \right) = 4\pi \delta(\vec{r} - \vec{r}')$$

We call the special solution for a point source
the Green function for the differential operator

$$-\nabla^2 G(\vec{r}, \vec{r}') = -4\pi \delta(\vec{r} - \vec{r}')$$

$G(\vec{r}, \vec{r}')$ gives the potential at position $\vec{r}$ due
to a unit source at position $\vec{r}'$

Generally, one also has to specify a desired
boundary condition for the Green function on
the boundary of the system.

For the Coulomb solution for a point charge
the implicit boundary condition is that the
potential vanish infinitely far from the charges

$$G(\vec{r}, \vec{r}') \rightarrow 0 \quad \text{as} \quad |\vec{r} - \vec{r}'| \rightarrow \infty$$

boundary of the system is taken to infinity

If one knows the Green's function, then one can find the solution for any distribution of sources $\rho(\vec{r})$

$$\phi(\vec{r}) = \int d^3r' G(\vec{r}, \vec{r}') \rho(\vec{r}')$$

proof:
$$-\nabla^2 \phi = \int d^3r' \left[-\nabla^2 G(\vec{r}, \vec{r}') \right] \rho(\vec{r}')$$
$$= \int d^3r' \left[4\pi \delta(\vec{r} - \vec{r}') \right] \rho(\vec{r}')$$
$$= 4\pi \rho(\vec{r})$$

We will return to concept of Green's function when we discuss solution of Poisson's eqn in a finite volume

We will also see Green's functions again when we discuss solution of the inhomogeneous wave equation.

The Coulomb problem as a boundary value problem

Consider a conducting sphere of radius R with net charge q . (as $R \rightarrow 0$ we get a point charge).

What is $\phi(\vec{r})$? What is $E(\vec{r})$?

Review: Properties of conductors in electrostatics

- 1) $\vec{E} = 0$ inside conductor - if $\vec{E} \neq 0$ then a current $\vec{j} = \sigma \vec{E}$ flows and it is not statics (σ is conductivity)
- 2) $\rho = 0$ inside conductor - if $\vec{E} = 0$ inside, then $\vec{\nabla} \cdot \vec{E} = 4\pi\rho = 0$
- 3) Any net charge on the conductor must lie on the surface - follows from (2)
- 4) $\phi = \text{constant}$ throughout conductor - if $\vec{E} = 0$ then $\vec{E} = -\vec{\nabla}\phi \Rightarrow \phi$ is constant
- 5) Just outside the conductor, $\vec{E}$ is $\perp$ to surface.
- If $\vec{E}$ has a component $\parallel$ to surface then it exerts a force on electrons at the surface, leading to a surface current - so would not be static

For conducting sphere, $\rho = 0$ for $r > R$ and $r < R$
all charge is on the surface $\Rightarrow \nabla^2\phi = 0$ for $\begin{cases} r > R \\ r < R \end{cases}$

spherical symmetry $\Rightarrow$ expect spherically symmetric solution

$\Rightarrow \phi(\vec{r})$ depends only on $r = |\vec{r}|$

⇒ Solve Laplace's eqn by writing ∇^2 in spherical coords.
Only the radial terms do not vanish.

$$\nabla^2 \phi = \frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{d\phi}{dr} \right) = 0$$

$$r^2 \frac{d\phi}{dr} = -C_0 \quad \text{a constant}$$

$$\frac{d\phi}{dr} = -\frac{C_0}{r^2}$$

$$\phi(r) = \frac{C_0}{r} + C_1, \quad C_1 \text{ a constant}$$

"outside" $r > R$ $\phi^{\text{out}}(r) = \frac{C_0^{\text{out}}}{r} + C_1^{\text{out}}$

"inside" $r < R$ $\phi^{\text{in}}(r) = \frac{C_0^{\text{in}}}{r} + C_1^{\text{in}}$

solution "outside" does not necessarily go smoothly into the solution "inside" because of the charge layer at $r=R$ that separates the two regions. We need to determine the constants $C_0^{\text{in}}, C_0^{\text{out}}, C_1^{\text{in}}, C_1^{\text{out}}$ by applying boundary conditions corresponding to the physical situation.

① For $r > R$, assume $\phi \rightarrow 0$ as $r \rightarrow \infty$ - boundary condition at infinity

$$\Rightarrow C_1^{\text{out}} = 0$$

$$\phi^{\text{out}}(r) = \frac{C_0^{\text{out}}}{r} \quad \text{recover the expected Coulomb form.}$$

2) For $r < R$.

i) we could use the fact that the region $r < R$ is a conductor with $\phi = \text{constant}$ to conclude $C_0^{\text{in}} = 0$


ii) or, if we were dealing with a charged shell instead of a conductor, we could argue as follows:

no charge at origin $r=0 \Rightarrow$ expect ϕ should be finite at origin $\Rightarrow C_0^{\text{in}} = 0$

So $\phi^{\text{in}}(r) = C^{\text{in}}$ a constant


3) Now we need boundary condition at $r=R$ where "inside" and "outside" meet.

Review: Electric field and potential at a surface charge layer



← a general surface S with surface charge density $\sigma(\vec{r})$ for $\vec{r}$ on S . $\sigma(\vec{r})da$ is total charge in area da on surface

i) Take "Gaussian pillbox" surface about point $\vec{r}$ on the surface S



Gauss' Law in integral form $\oint_S d\vec{a} \cdot \hat{n} \cdot \vec{E} = 4\pi Q_{\text{enclosed}}$

expect $\vec{E}$ is finite $\rightarrow$ contribution from sides of pillbox vanish as $dl \rightarrow 0$.

$$\oint_S d\vec{a} \cdot \vec{E} = \int_{\text{top}} d\vec{a} \cdot \vec{E} + \int_{\text{bottom}} d\vec{a} \cdot \vec{E}$$

$$= (\hat{n}^{\text{top}} \cdot \vec{E}^{\text{top}} + \hat{n}^{\text{bottom}} \cdot \vec{E}^{\text{bottom}}) da \quad \text{since } da \text{ is small}$$


$\vec{E}^{\text{top}}$ is electric field at $\vec{r}$ just above the surface S
 $\vec{E}^{\text{bottom}}$ is electric field at $\vec{r}$ just below the surface S

$\hat{n}^{\text{top}} \equiv \hat{n}$ is outward normal on top
 $\hat{n}^{\text{bottom}} = -\hat{n}$ is outward normal on bottom

$$\Rightarrow (\vec{E}^{\text{top}} - \vec{E}^{\text{bottom}}) \cdot \hat{n} da = 4\pi Q_{\text{enclosed}} = 4\pi \sigma(\vec{r}) da$$

$$\boxed{(\vec{E}^{\text{top}} - \vec{E}^{\text{bottom}}) \cdot \hat{n} = 4\pi \sigma(\vec{r})} \quad \text{discontinuity in normal component of } \vec{E}$$

ii) Take "Amperian loop" C at surface about point $\vec{r}$.



$$\vec{\nabla} \times \vec{E} = 0 \Rightarrow \oint_C d\vec{l} \cdot \vec{E}$$

since $\vec{E}$ is finite at surface,
 if take sides $dl' \rightarrow 0$ their contribution to integral vanishes,

$$\Rightarrow \oint_C d\vec{l} \cdot \vec{E} = \boxed{(\vec{E}^{\text{top}} - \vec{E}^{\text{bottom}}) \cdot d\vec{l} = 0}$$

where $d\vec{l}$ is any infinitesimal tangent to the surface at $\vec{r}$.

⇒ tangential component of $\vec{E}$ is continuous

combine above to write $\boxed{\vec{E}^{\text{top}} - \vec{E}^{\text{bottom}} = 4\pi\sigma(F) \hat{n}}$

iii) $\vec{E} = -\vec{\nabla}\phi \Rightarrow \phi(r_2) - \phi(r_1) = - \int_{r_1}^{r_2} d\vec{l} \cdot \vec{E}$

Take $\vec{r}_2$ just above $\vec{r}$ on surface
 $\vec{r}_1$ just below $\vec{r}$ on surface $\left. \vphantom{\begin{matrix} \vec{r}_2 \\ \vec{r}_1 \end{matrix}} \right\} d\vec{l} \rightarrow 0$

Since $\vec{E}$ is finite $\Rightarrow \int d\vec{l} \cdot \vec{E} \rightarrow 0$

$$\Rightarrow \boxed{\phi^{\text{top}} = \phi^{\text{bottom}}}$$

potential ϕ is continuous at surface charge layer

can rewrite (i) as

$$(-\vec{\nabla}\phi^{\text{top}} + \vec{\nabla}\phi^{\text{bottom}}) \cdot \hat{n} = 4\pi\sigma$$

$$\boxed{-\frac{\partial\phi^{\text{top}}}{\partial n} + \frac{\partial\phi^{\text{bottom}}}{\partial n} = 4\pi\sigma}$$

↑ directional derivative of ϕ in direction $\hat{n}$

discontinuity in normal derivative of ϕ at surface

Apply to conducting sphere

$$\phi \text{ continuous} \Rightarrow \phi^{\text{in}}(R) = \phi^{\text{out}}(R)$$

$$C_1^{\text{in}} = \frac{C_0^{\text{out}}}{R}$$

only one unknown left

normal derivative of ϕ is discontinuous

$$-\frac{\partial \phi^{\text{top}}}{\partial n} + \frac{\partial \phi^{\text{bottom}}}{\partial n} = 4\pi\sigma$$

here $\hat{n} = \hat{r}$ the radial direction

$$\left[-\frac{d\phi^{\text{out}}}{dr} + \frac{d\phi^{\text{in}}}{dr} \right]_{r=R} = 4\pi\sigma$$

but $\frac{d\phi^{\text{in}}}{dr} = 0$ as $\phi^{\text{in}} = \text{constant}$

$$-\frac{d\phi^{\text{out}}}{dr} \Big|_{r=R} = 4\pi\sigma$$

charge q is uniformly distributed on surface at R

$$-\frac{d}{dr} \left(\frac{C_0^{\text{out}}}{r} \right)_{r=R} = \frac{C_0^{\text{out}}}{R^2} = 4\pi\sigma = 4\pi \left(\frac{q}{4\pi R^2} \right) = \frac{q}{R^2}$$

$$\Rightarrow C_0^{\text{out}} = q, \quad C_0^{\text{in}} = \frac{q^{\text{out}}}{R} = \frac{q}{R}$$

$$\phi(r) = \begin{cases} \frac{q}{R} & r < R \text{ inside} \\ \frac{q}{r} & r > R \text{ outside} \end{cases}$$

$$\Rightarrow \vec{E} = -\vec{\nabla}\phi = -\frac{d\phi}{dr} = \begin{cases} 0 & r < R \text{ inside} \\ \frac{q}{r^2} & r > R \text{ outside} \end{cases}$$

we get familiar Coulomb solution!

Summary We can view the preceding solution for ϕ_{out} as solving Laplace's eqn $\nabla^2 \phi = 0$ subject to a specified boundary condition on the normal derivative of ϕ at the boundary $r=R$ of the "outside" region of the system.

Alternate problem:

Another physical situation would be to connect conducting sphere to a battery that charges the sphere to a fixed voltage ϕ_0 (stat volts!) with respect to ground $\phi=0$ at $r \rightarrow \infty$.

As before, outside the sphere $\phi = \frac{C_0}{r}$
Now the boundary condition is to specify the value of ϕ on the boundary of the outside region, i.e.

$$\phi(R) = \phi_0$$

$$\Rightarrow \frac{C_0}{R} = \phi_0, \quad C_0 = \phi_0 R$$

$$\phi(r) = \phi_0 \frac{R}{r}$$

(from preceding solution, we know that charging the sphere to voltage ϕ_0 (statvolts) induces a net charge $q = \phi_0 R$ on it)

These two versions of the conducting sphere problem are examples of a more general boundary value problem

Solve $\nabla^2 \phi = 0$ in a given region of space subject to one of the following two types of boundary conditions on the bounding surfaces of the region

i) Neumann boundary condition

$\frac{\partial \phi}{\partial n}$ - normal derivative of ϕ is specified on the bounding surface

ii) Dirichlet boundary condition

ϕ - value of ϕ is specified on the bounding surfaces

If the bounding surfaces consist of disjoint pieces, it is possible to specify either (i) or (ii) on each piece separately to get a mixed boundary value problem.

Some more problems

infinite conducting wire of radius R with line charge density λ = charge per unit length



$$\text{surface charge } \sigma = \frac{\lambda}{2\pi R}$$

Expect cylindrical symmetry $\Rightarrow \phi$ depends only on cylindrical coord r .

$$\nabla^2 \phi = 0 \quad \text{for } r > R, \quad r < R$$

use ∇^2 in cylindrical coords - only radial term non vanishing

$$\nabla^2 \phi = \frac{1}{r} \frac{d}{dr} \left(r \frac{d\phi}{dr} \right) = 0$$

$$r \frac{d\phi}{dr} = C_0 \quad \text{constant}$$

$$\frac{d\phi}{dr} = \frac{C_0}{r}$$

$$\phi(r) = C_0 \ln r + C_1 \quad \text{const}$$

note: one cannot now choose $\phi \rightarrow 0$ as $r \rightarrow \infty$!

one needs to fix zero of ϕ at some other radius. a convenient choice is $r = R$, but any other choice could also be made.

$$\phi^{\text{out}} = C_0^{\text{out}} \ln r + C_1^{\text{out}}$$

$$\phi^{\text{in}} = C_0^{\text{in}} \ln r + C_1^{\text{in}}$$

$$\phi^{\text{in}} = \text{const in conductor} \rightarrow C_0^{\text{in}} = 0$$

$$\text{or } \phi^{\text{in}} \text{ should not diverge as } r \rightarrow 0 \Rightarrow C_0^{\text{in}} = 0$$

$$\text{so } \phi^{\text{in}} = C_1^{\text{in}} \text{ constant}$$

boundary condition at $r=R$

$$\left[-\frac{d\phi^{\text{out}}}{dr} + \frac{d\phi^{\text{in}}}{dr} \right]_{r=R} = 4\pi\sigma$$

$$\Rightarrow -\frac{C_0^{\text{out}}}{R} = 4\pi\sigma = 4\pi \left(\frac{\lambda}{2\pi R} \right) = \frac{2\lambda}{R}$$

$$C_0^{\text{out}} = -2\lambda$$

$$\phi^{\text{out}}(r) = -2\lambda \ln r + C_1^{\text{out}}$$

continuity of ϕ

$$\phi^{\text{in}}(R) = \phi^{\text{out}}(R) \Rightarrow C_1^{\text{in}} = -2\lambda \ln R + C_1^{\text{out}}$$

Remaining const C_1^{out} is not too important as it is just a common additive constant to both ϕ^{in} and $\phi^{\text{out}} \Rightarrow$ does not change $\vec{E} = -\vec{\nabla}\phi$.


If we use the condition $\phi(R)=0$ then we can solve for C_1^{out} .

$$0 = -2\lambda \ln R + C_1^{\text{out}}$$

$$\Rightarrow C_1^{\text{out}} = 2\lambda \ln R$$

$$\Rightarrow \phi(r) = \begin{cases} -2\lambda \ln(r/R) & r \geq R \\ 0 & r < R \end{cases}$$

infinite conducting half space $\Rightarrow \vec{E}(\vec{r}) = \begin{cases} \frac{2\lambda}{r} \hat{r} & r \geq R \\ 0 & r < R \end{cases}$



expect ϕ depends only on x

$$\nabla^2 \phi = \frac{d^2 \phi}{dx^2} = 0$$

$$\rightarrow \begin{cases} \phi^>(x) = C_0^> x + C_1^> & x > 0 \\ \phi^<(x) = C_0^< x + C_1^< & x < 0 \end{cases}$$

for $x < 0$, $\phi = \text{const}$ in conductor $\Rightarrow C_0^< = 0$

at $x=0$, ϕ continuous $\Rightarrow \phi^<(0) = \phi^>(0)$

$$C_1^< = C_1^>$$

$\frac{d\phi}{dx}$ discontinuous $\Rightarrow$

$$-\left. \frac{d\phi}{dx} \right|_{x=0} = 4\pi\sigma$$

$$C_0^> = -4\pi\sigma$$

$$\Rightarrow \phi(x) = \begin{cases} -4\pi\sigma x + C_1^> & x > 0 \\ C_1^> & x < 0 \end{cases}$$

const $C_1^>$ does not change value of $\vec{E}$

as for the wire, we cannot choose $\phi \rightarrow 0$ as $x \rightarrow \infty$.

~~we can set $\phi = 0$ at~~

$$-\vec{\nabla}\phi = \vec{E} = \begin{cases} 4\pi\sigma \hat{x} & x > 0 \\ 0 & x < 0 \end{cases}$$

infinite charged plane

similar to previous problem, but now no conductor at $x < 0$, just free space on both sides of the charged plane at $x = 0$.

~~expect ϕ depends on $|x|$ by symmetry~~

$$\nabla^2\phi = \frac{d^2\phi}{dx^2} = 0 \Rightarrow \begin{aligned} \phi^> &= C_0^> x + C_1^> & x > 0 \\ \phi^< &= C_0^< x + C_1^< & x < 0 \end{aligned}$$

continuity of ϕ at $x = 0$

$$\Rightarrow \phi^>(0) = \phi^<(0) \Rightarrow C_1^> = C_1^<$$

discontinuity of $d\phi/dx$ at $x = 0$

$$-\frac{d\phi^>}{dx} + \frac{d\phi^<}{dx} = 4\pi\sigma$$

$$-C_0^> + C_0^< = 4\pi\sigma$$

$$\text{Define } \bar{C}_0 = \frac{C_0^> + C_0^<}{2}$$

Then we can write

$$\begin{aligned} C_0^< &= \bar{C}_0 + 2\pi\sigma \\ C_0^> &= \bar{C}_0 - 2\pi\sigma \end{aligned}$$

$$\phi = \begin{cases} -2\pi\sigma x + \bar{C}_0 x + C_i^> & x > 0 \\ 2\pi\sigma x + \bar{C}_0 x + C_i^< & x < 0 \end{cases}$$

$$\Rightarrow -\frac{d\phi}{dx} = \vec{E} = \begin{cases} (2\pi\sigma - \bar{C}_0) \hat{x} & x > 0 \\ (-2\pi\sigma - \bar{C}_0) \hat{x} & x < 0 \end{cases}$$

Const $C_i^>$ does not effect $\vec{E}$ - additive const to ϕ

$\bar{C}_0$ represents const uniform electric field $-\bar{C}_0 \hat{x}$,
that exists independently of the charged surface

If we assumed that all $\vec{E}$ fields are just those arising from the plane, then we can set $\bar{C}_0 = 0$.
Equivalently, if the plane is the only source of $\vec{E}$, then we expect ϕ depends only on $|x|$ by symmetry.
 $\Rightarrow C_0^< = -C_0^>$ and again $\bar{C}_0 = 0$. In this case

$$\phi(x) = \begin{cases} -2\pi\sigma x & x > 0 \\ 2\pi\sigma x & x < 0 \end{cases} \quad \left(\begin{array}{l} \text{we also set} \\ C_i^> = 0 \text{ here} \\ \text{corresponding} \\ \text{to } \phi(0) = 0 \end{array} \right)$$

$$\vec{E}(x) = \begin{cases} 2\pi\sigma \hat{x} & x > 0 \\ -2\pi\sigma \hat{x} & x < 0 \end{cases}$$

$\vec{E}$ is constant ~~by~~ but oppositely directed on either side of the charged plane