

Electric Dipole Approximation - the leading term in non-relativistic expansion

$$\vec{A}_{E1}(\vec{r}) = -ik \vec{p}_\omega \frac{e^{ikr}}{r}$$

$$\vec{\nabla} \times (\phi \vec{F}) = (\vec{\nabla} \phi) \times \vec{F} + \phi \vec{\nabla} \times \vec{F}$$

$$\vec{B}_{E1} = \vec{\nabla} \times \vec{A}_{E1} = -ik \left(\vec{\nabla} \frac{e^{ikr}}{r} \right) \times \vec{p}_\omega$$

$$= -ik \left(ik - \frac{1}{r} \right) \frac{e^{ikr}}{r} \hat{r} \times \vec{p}_\omega$$

$$= k^2 \frac{e^{ikr}}{r} \left(1 + \frac{i}{kr} \right) \hat{r} \times \vec{p}_\omega$$

In radiation zone approx, $kr, \gg 1$

$$\boxed{\vec{B}_{E1} \approx k^2 \frac{e^{ikr}}{r} \hat{r} \times \vec{p}_\omega}$$

To get electric field, use Ampere's Law

$$\vec{\nabla} \times \vec{B} = \frac{1}{c} \frac{\partial \vec{E}}{\partial t} \quad (\text{away from source where } \vec{J} = 0)$$

For oscillating fields $\vec{E} = E_\omega e^{-i\omega t}$

$$\vec{\nabla} \times \vec{B}_\omega = -\frac{i\omega}{c} \vec{E}_\omega \Rightarrow E_{E1} = \frac{i}{k} \vec{\nabla} \times \vec{B}_{E1}$$

$$\vec{E}_{E1} = \frac{i}{k} \vec{\nabla} \times \left[k^2 \frac{e^{ikr}}{r} \left(1 + \frac{i}{kr} \right) \hat{r} \times \vec{p}_\omega \right]$$

$$\vec{E}_{E1} = \frac{i}{k} (\vec{\nabla} e^{ikr}) \times \left[\frac{k^2}{r} \left(1 + \frac{i}{kr}\right) \hat{r} \times \vec{p}_\omega \right]$$

← ignore in RZ approx

$$+ \frac{i}{k} e^{ikr} \vec{\nabla} \times \left[\frac{k^2}{r} \left(1 + \frac{i}{kr}\right) \hat{r} \times \vec{p}_\omega \right]$$

this will always be of order $1/r^2$

so can ignore it in radiation zone approx

So in radiation zone approx

$$\vec{E}_{E1} = (\vec{\nabla} e^{ikr}) \times \left[\frac{ik}{r} \hat{r} \times \vec{p}_\omega \right]$$

$$\boxed{\vec{E}_{E1} = -k^2 \frac{e^{ikr}}{r} \hat{r} \times (\hat{r} \times \vec{p}_\omega)}$$

if do not make radiation zone approx, one gets

$$\vec{E}_{EI} = \frac{k^2 e^{ikr}}{r} \left[\vec{p}_w - \hat{r}(\vec{p}_w \cdot \hat{r}) - \frac{i}{kr} \left(1 + \frac{i}{kr}\right) (3\hat{r}(\vec{p}_w \cdot \hat{r}) - \vec{p}_w) \right]$$

Using radiation zone approx:


$$\vec{E}_{EI} = k^2 \frac{e^{ikr}}{r} \hat{r} \times (\vec{p}_w \times \hat{r})$$

$$|\vec{E}_{EI}| = |\vec{B}_{EI}|$$

$$\vec{B}_{EI} = -k^2 \frac{e^{ikr}}{r} \vec{p}_w \times \hat{r}$$

$$\vec{E}_{EI} \perp \vec{B}_{EI}$$

If choose coordinates so that $\vec{p}_w$ is along $\hat{z}$ axis, then



$$\vec{E}_{EI} = -k^2 p_w \frac{e^{ikr}}{r} \sin \theta \hat{\theta}$$

$$\vec{B}_{EI} = -k^2 p_w \frac{e^{ikr}}{r} \sin \theta \hat{\phi}$$

Emitted power

pointing vector $\vec{S}_{EI}(\vec{r}, t) = \frac{c}{4\pi} \text{Re}\{\vec{E}_{EI}\} \times \text{Re}\{\vec{B}_{EI}\}$

need to take real parts of complex expression before multiplying

$$\text{Re}\{\vec{E}_{EI}(\vec{r}, t)\} = -k^2 p_w \frac{\cos(kr - \omega t)}{r} \sin \theta \hat{\theta}$$

$$\text{Re}\{\vec{B}_{EI}(\vec{r}, t)\} = -k^2 p_w \frac{\cos(kr - \omega t)}{r} \sin \theta \hat{\phi}$$

$$\boxed{\vec{S}_{EI}(\vec{r}, t) = \frac{c}{4\pi} k^4 p_w^2 \frac{\cos^2(kr - \omega t)}{r^2} \sin^2 \theta \hat{r}}$$

$\vec{S}_{EI} \sim \hat{r} \Rightarrow$ energy flows radially outwards

$\vec{S}_{EI} \sim \frac{1}{r^2} \Rightarrow$ energy conserved

$$\oint_{\text{sphere radius } R} da \hat{m} \cdot \langle \vec{S}_{EI} \rangle = \text{constant for all } R$$

Question - what about the

$\frac{1}{r^n}, n \neq 2$, terms if we do not make radiation zone approx?

time averaged energy current

$$\langle \vec{S}_{EI} \rangle = \frac{1}{T} \int_0^T dt \vec{S}_{EI}(\vec{r}, t)$$

T is period $T = \frac{2\pi}{\omega}$

$$\langle \cos^2(\cdot) \rangle = \frac{1}{2}$$

$$= \frac{c}{8\pi} k^4 p_\omega^2 \frac{\sin^2 \theta}{r^2} \hat{r}$$

average energy flowing through an element of area at spherical angles θ, ϕ is

$$dP_{EI} = \hat{r} \cdot \langle \vec{S}_{EI} \rangle \underbrace{r^2 \sin \theta d\theta d\phi}_{\text{area of surface element}}$$


area of surface element

$= r^2 d\Omega$ Ω is solid angle

$$= \hat{r} \cdot \langle \vec{S}_{EI} \rangle r^2 d\Omega$$

$$\frac{dP_{EI}}{d\Omega} = \hat{r} \cdot \langle \vec{S}_{EI} \rangle r^2 = \frac{c}{8\pi} k^4 p_\omega^2 \sin^2 \theta \sim \omega^4 \sin^2 \theta$$

polar plot



rotationally symmetric about $\hat{z}$ or $\hat{p}_\omega$

$$\frac{dP_{EI}}{d\Omega} \sim \sin^2 \theta$$

most of power is directed outwards into plane $\perp \vec{p}_\omega$, i.e. peaked about $\theta = 90^\circ$


Total power radiated is

$$P_{EI} = \int \frac{dP_{EI}}{d\Omega} d\Omega = \frac{ck^4 p_w^2}{8\pi} 2\pi \int_0^\pi \sin\theta \sin^2\theta d\theta$$

$$P_{EI} = \frac{ck^4 p_w^2}{3} = \frac{p_w^2 \omega^4}{3c^3} \sim \omega^4$$

why the sky is blue - Lord Rayleigh

When look up at sky, you are seeing the indirect light of the sun, which is the light emitted by the atoms and molecules of the atmosphere as they oscillate, and so radiate, due to the electric field of the direct light from the sun



power of the radiated indirect light is $P \sim \omega^4 p_0^2$

$$\vec{p} = \alpha \vec{E} \quad \alpha \sim \frac{e^2}{m} \frac{1}{\omega_0^2 - \omega^2 - i\omega\gamma}$$

For molecules in atmosphere (N_2 etc) ω_0 is typically at a freq higher than visible spectrum. Therefore, in visible spectrum $\alpha \sim \frac{e^2}{m\omega_0^2}$ indep of ω .

$\Rightarrow$ power radiated is $P \sim \omega^4$

$P \sim \omega^4$ largest at high freq

Since light from sun is "white light"

it has components of all freqs. Of these freqs, the higher ones are scattered the most & make up the indirect light we see.

Since blue is the largest ω in visible spectrum, the sky is blue!

When we look at sunrise or sunset, we are looking at the direct rays of the sun. Since these rays are least scattered at low $\omega \Rightarrow$ sunset and sunrise are red!

Magnetic Dipole approx - Radiation Zone $kr \gg 1$

$$\vec{A}_M = \frac{e^{ikr}}{r} \left(\frac{1}{r} - ik \right) (-\hat{r} \times \vec{m}_\omega)$$

$$\approx ik \hat{r} \times \vec{m}_\omega \frac{e^{ikr}}{r} \quad \text{in RZ}$$

$$\vec{B}_M = \vec{\nabla} \times \vec{A}_M = (\vec{\nabla} e^{ikr}) \times \left(\frac{ik \hat{r} \times \vec{m}_\omega}{r} \right)$$

$$+ e^{ikr} \vec{\nabla} \times \left(\frac{ik \hat{r} \times \vec{m}_\omega}{r} \right)$$

will give terms of $o(\frac{1}{r^2})$
so ignore in RZ approx

$$\vec{B}_M = -k^2 \frac{e^{ikr}}{r} \hat{r} \times (\hat{r} \times \vec{m}_\omega)$$

From Ampere's Law

$$\vec{E}_M = \frac{1}{k} \vec{\nabla} \times \vec{B}_M = -ik (\vec{\nabla} e^{ikr}) \times \left(\frac{\hat{r} \times [\hat{r} \times \vec{m}_\omega]}{r} \right)$$

$$- ik e^{ikr} \vec{\nabla} \times \left(\frac{\hat{r} \times [\hat{r} \times \vec{m}_\omega]}{r} \right)$$

will give terms of $o(\frac{1}{r^2})$
so ignore in RZ approx

$$\vec{E}_M = k^2 \frac{e^{ikr}}{r} \hat{r} \times (\hat{r} \times (\hat{r} \times \vec{m}_\omega))$$

triple product rule


$$= k^2 \frac{e^{ikr}}{r} \left\{ \hat{r} [\hat{r} \cdot (\hat{r} \times \vec{m}_\omega)] - (\hat{r} \times \vec{m}_\omega) [\hat{r} \cdot \hat{r}] \right\}$$

$$\vec{E}_M = -\frac{k^2}{r} e^{ikr} (\hat{r} \times \vec{m}_\omega)$$

If align axes so that $\vec{m}_\omega = m_\omega \hat{z}$ then

$$\vec{E}_{M1} = m_\omega \frac{k^2}{r} e^{i(kr - \omega t)} \sin \theta \hat{\phi}$$

$$\vec{B}_{M1} = -m_\omega \frac{k^2}{r} e^{i(kr - \omega t)} \sin \theta \hat{\theta}$$




Poynting vector

$$\vec{S}_{M1} = \frac{c}{4\pi} \text{Re}\{\vec{E}_{M1}\} \times \text{Re}\{\vec{B}_{M1}\}$$

$$= \frac{c}{4\pi} \frac{k^4 m_\omega^2}{r^2} \cos^2(kr - \omega t) \sin^2 \theta \hat{r}$$

$$\langle \vec{S}_{M1} \rangle = \frac{c}{8\pi} \frac{k^4 m_\omega^2}{r^2} \sin^2 \theta \hat{r}$$

$$\frac{dP_{M1}}{d\Omega} = \hat{r} \cdot \langle \vec{S}_{M1} \rangle r^2 = \frac{c}{8\pi} k^4 m_\omega^2 \sin^2 \theta \sim \omega^4 \sin^2 \theta$$



rotationally symmetric
about $\hat{z}$ axis

$$P_{M1} = \int d\Omega \frac{dP_{M1}}{d\Omega} = \frac{c k^4}{3} m_\omega^2 = \frac{m_\omega^2 \omega^4}{3c^3}$$

$$\frac{P_{M1}}{P_{E1}} = \frac{m_\omega^2}{p_\omega^2} \sim \left(\frac{v}{c}\right)^2 \quad m_\omega \sim \frac{df}{dg} \sim dg \frac{v}{c}$$

$$p_\omega \sim dg$$

Electric Quadrupole radiation - radiation zone approx

$$\vec{A}_{E2} = \frac{e^{ikr}}{r} \left(\frac{1}{r} - ik \right) \left(\frac{-i\omega}{6c} \hat{r} \cdot \vec{Q}_\omega \right)$$

$$= -\frac{e^{ikr}}{r} \frac{k^2}{6} \hat{r} \cdot \vec{Q}_\omega \quad \text{in RZ approx}$$

$$\begin{aligned} \vec{B}_{E2} &= \vec{\nabla} \times \vec{A}_{E2} = -(\vec{\nabla} e^{ikr}) \times \left[\frac{k^2 \hat{r} \cdot \vec{Q}_\omega}{6r} \right] \\ &\quad - e^{ikr} \vec{\nabla} \times \left[\frac{k^2 \hat{r} \cdot \vec{Q}_\omega}{6r} \right] \end{aligned}$$

$$\boxed{\vec{B}_{E2} = -ik^3 \frac{e^{ikr}}{6r} \hat{r} \times (\hat{r} \cdot \vec{Q}_\omega)} \quad \text{O}\left(\frac{1}{r^2}\right) \text{ so ignore in RZ approx}$$

$$\begin{aligned} \vec{E}_{E2} &= \frac{i}{k} \vec{\nabla} \times \vec{B}_{E2} = k^2 (\vec{\nabla} e^{ikr}) \times \left[\frac{\hat{r} \times (\hat{r} \cdot \vec{Q}_\omega)}{6r} \right] \\ &\quad + k^2 e^{ikr} \vec{\nabla} \times \left[\frac{\hat{r} \times (\hat{r} \cdot \vec{Q}_\omega)}{6r} \right] \end{aligned}$$

O(1/r) so ignore in RZ approx

$$\vec{E}_{E2} = ik^3 \frac{e^{ikr}}{6r} \hat{r} \times [\hat{r} \times (\hat{r} \cdot \vec{Q}_\omega)]$$

triple product rule

$$= \frac{ik^3 e^{ikr}}{6r} \left\{ \hat{r} [\hat{r} \cdot (\hat{r} \cdot \vec{Q}_\omega)] - (\hat{r} \cdot \vec{Q}_\omega) [\hat{r} \cdot \hat{r}] \right\}$$

$$\boxed{\vec{E}_{E2} = \frac{ik^3 e^{ikr}}{6r} \left\{ \hat{r} (\hat{r} \cdot \vec{Q}_\omega \cdot \hat{r}) - (\hat{r} \cdot \vec{Q}_\omega) \right\}}$$

Poynting vector

$$\vec{S}_{E2} = +\frac{c}{4\pi} \operatorname{Re} \{ \vec{E}_{E2} \} \times \operatorname{Re} \{ \vec{B}_{E2} \}$$

$$\stackrel{(94\pi)}{=} \frac{-k^6}{36r^2} \sin^2(kr - \omega t) \left\{ \hat{r} (\hat{r} \cdot \vec{Q}_\omega \cdot \hat{r}) - (\hat{r} \cdot \vec{Q}_\omega) \right\} \times \left[\hat{r} \times (\hat{r} \cdot \vec{Q}_\omega) \right]$$

$$\stackrel{(94\pi)}{=} \frac{-k^6}{36r^2} \sin^2(kr - \omega t) \left\{ \hat{r} \left[\hat{r} (\hat{r} \cdot \vec{Q}_\omega \cdot \hat{r}) \cdot (\hat{r} \cdot \vec{Q}_\omega) \right] \right. \\ \left. - (\hat{r} \cdot \vec{Q}_\omega) \left[\hat{r} (\hat{r} \cdot \vec{Q}_\omega \cdot \hat{r}) \cdot \hat{r} \right] \right.$$

$$\left. - \hat{r} \left[(\hat{r} \cdot \vec{Q}_\omega) \cdot (\hat{r} \cdot \vec{Q}_\omega) \right] \right. \\ \left. + (\hat{r} \cdot \vec{Q}_\omega) \left[(\hat{r} \cdot \vec{Q}_\omega) \cdot \hat{r} \right] \right\}$$

$$\stackrel{(94\pi)}{=} \frac{-k^6}{36r^2} \sin^2(kr - \omega t) \left\{ (\hat{r} \cdot \vec{Q}_\omega \cdot \hat{r})^2 \hat{r} - (\hat{r} \cdot \vec{Q}_\omega) (\hat{r} \cdot \vec{Q}_\omega \cdot \hat{r}) \right. \\ \left. - (\hat{r} \cdot \vec{Q}_\omega \cdot \vec{Q}_\omega \cdot \hat{r}) \hat{r} + (\hat{r} \cdot \vec{Q}_\omega) (\hat{r} \cdot \vec{Q}_\omega \cdot \hat{r}) \right\}$$

$$\vec{S}_{E2} = \frac{-ck^6}{4\pi 36r^2} \sin^2(kr - \omega t) \left\{ (\hat{r} \cdot \vec{Q}_\omega \cdot \hat{r})^2 - (\hat{r} \cdot \vec{Q}_\omega)^2 \right\} \hat{r}$$

$$\langle \vec{S}_{E2} \rangle = \frac{-ck^6}{4\pi 72r^2} \left\{ (\hat{r} \cdot \vec{Q}_\omega \cdot \hat{r})^2 - (\hat{r} \cdot \vec{Q}_\omega)^2 \right\} \hat{r}$$

$$\frac{dP_{E2}}{d\Omega} = \hat{r} \cdot \langle \vec{S}_{E2} \rangle r^2 = \frac{-ck^6}{4\pi 72} \left\{ (\hat{r} \cdot \vec{Q}_\omega)^2 - (\hat{r} \cdot \vec{Q}_\omega \cdot \hat{r})^2 \right\}$$


angular dependence of $\frac{dP_{E2}}{d\Omega}$ depends
on specific form of the tensor $\vec{Q}_\omega$

For example: suppose $Q_{ij} = 0$ except for Q_{zz}

$$\Rightarrow \vec{Q}_w = Q_{zz} \hat{z} \hat{z}$$

$$(\hat{r} \cdot \vec{Q}_w \cdot \hat{r})^2 = (Q_{zz} \cos^2 \theta)^2$$

$$(\hat{r} \cdot \vec{Q}_w)^2 = Q_{zz}^2 \cos^2 \theta$$




$$\frac{dP_{E2}}{d\Omega} = \frac{c k^6}{4\pi^2} Q_{zz}^2 [\cos^2 \theta - \cos^4 \theta]$$

$$= \frac{c k^6}{4\pi^2} Q_{zz}^2 \cos^2 \theta \sin^2 \theta$$

$$\cos \theta \sin \theta = \frac{1}{2} \sin 2\theta$$

$$= \frac{c k^6}{4\pi^2} Q_{zz}^2 \sin^2 2\theta$$



peak at 45°

rotationally invariant
about $\hat{z}$ axis

$$\frac{P_{E2}}{P_{E1}} \sim \frac{k^6 Q^2}{k^4 p^2} \sim \frac{k^2 (q d^2)^2}{(q d)^2} \sim k^2 d^2 \sim \left(\frac{v}{c}\right)^2$$

$$P_{E2} \sim P_{M1}$$

For more general case, choose axes so that $\vec{Q}$ is diagonal - can always do this since $\vec{Q}$ is symmetric

$$(\hat{r} \cdot \vec{Q} \cdot \hat{r}) = \hat{r} \cdot \begin{pmatrix} Q_{xx} & 0 & 0 \\ 0 & Q_{yy} & 0 \\ 0 & 0 & Q_{zz} \end{pmatrix} \cdot \hat{r}$$

$$= \hat{r} \cdot \begin{pmatrix} Q_{xx} \sin\theta \cos\varphi \\ Q_{yy} \sin\theta \sin\varphi \\ Q_{zz} \cos\theta \end{pmatrix} = Q_{xx} \sin^2\theta \cos^2\varphi + Q_{yy} \sin^2\theta \sin^2\varphi + Q_{zz} \cos^2\theta$$

$$(\hat{r} \cdot \vec{Q})^2 = Q_{xx}^2 \sin^2\theta \cos^2\varphi + Q_{yy}^2 \sin^2\theta \sin^2\varphi + Q_{zz}^2 \cos^2\theta$$

$$\frac{dP_{E2}}{d\Omega} = \frac{k^6}{72} \left\{ Q_{zz}^2 (\cos^2\theta - \cos^4\theta) + Q_{xx}^2 (\sin^2\theta \cos^2\varphi - \sin^4\theta \cos^4\varphi) + Q_{yy}^2 (\sin^2\theta \sin^2\varphi - \sin^4\theta \sin^4\varphi) \right\}$$

$$\frac{dP_{E2}}{d\Omega} = \frac{k^6}{72} \left\{ Q_{zz}^2 \cos^2\theta \sin^2\theta + Q_{xx}^2 \sin^2\theta \cos^2\varphi (1 - \sin^2\theta \cos^2\varphi) + Q_{yy}^2 \sin^2\theta \sin^2\varphi (1 - \sin^2\theta \sin^2\varphi) \right\}$$

no special symmetries - varies with θ and φ