

Note: Lorentz gauge condition does not uniquely determine \vec{A} and ϕ . If one constructs \vec{A} and ϕ obeying Lorentz gauge condition, and then constructs

$$\begin{aligned}\vec{A}' &= \vec{A} + \vec{\nabla}\chi \\ \phi' &= \phi - \frac{1}{c} \frac{\partial \chi}{\partial t}\end{aligned}$$

then \vec{A}' and ϕ' will also be in Lorentz gauge provided $\square^2 \chi = 0$ (proof left to reader)

2) Coulomb Gauge

gauge constraint: require $\vec{\nabla} \cdot \vec{A} = 0$
if \vec{A} is in the Coulomb Gauge, then $\vec{A}' = \vec{A} + \vec{\nabla}\chi$ will also be in Coulomb gauge provided $\nabla^2 \chi = 0$.

Then Gauss' law becomes

$$\nabla^2 \phi + \frac{1}{c} \frac{\partial}{\partial t} (\vec{\nabla} \cdot \vec{A}) = -4\pi\rho$$

$$\Rightarrow \boxed{\nabla^2 \phi = -4\pi\rho} \quad \text{same as electrostatics!}$$

$$\Rightarrow \phi(\vec{r}, t) = \int d^3r' \frac{\rho(\vec{r}', t)}{|\vec{r} - \vec{r}'|}$$

no matter what motion the source $\rho(\vec{r}, t)$ has! ϕ is given by the instantaneous Coulomb potential even though electromagnetic fields have a finite velocity of propagation c !

Ampere's law becomes:

$$-\nabla^2 A + \frac{1}{c^2} \frac{\partial^2 \vec{A}}{\partial t^2} = \frac{4\pi}{c} \vec{j} - \vec{\nabla} (\vec{\nabla} \cdot \vec{A} + \frac{1}{c} \frac{\partial \phi}{\partial t})$$

$$\Rightarrow \square^2 A = \frac{4\pi}{c} \vec{j} - \frac{1}{c} \vec{\nabla} \left(\frac{\partial \phi}{\partial t} \right)$$

$$\text{where } \vec{\nabla} \left(\frac{\partial \phi}{\partial t} \right) = \vec{\nabla} \left[\int d^3r' \frac{\partial \rho}{\partial t} \frac{1}{|\vec{r} - \vec{r}'|} \right]$$

$$= - \vec{\nabla} \left[\int d^3r' \frac{\vec{\nabla}' \cdot \vec{j}(\vec{r}', t)}{|\vec{r} - \vec{r}'|} \right] \quad \text{by continuity eqn.}$$

To see the meaning of this term, recall - any vector function \vec{j} can be written as the sum of a curlfree and a divergenceless part

$$\vec{j} = \vec{j}_{||} + \vec{j}_{\perp} \quad \text{where } \vec{\nabla} \times \vec{j}_{||} = 0 \quad \text{curlfree}$$

$$\vec{\nabla} \cdot \vec{j}_{\perp} = 0 \quad \text{divergenceless}$$

where

$$\vec{j}_{||}(\vec{r}) = -\frac{1}{4\pi} \vec{\nabla} \int d^3r' \frac{\vec{\nabla}' \cdot \vec{j}(\vec{r}')}{|\vec{r} - \vec{r}'|} \quad \text{longitudinal part}$$

$$\vec{j}_{\perp}(\vec{r}) = \frac{1}{4\pi} \vec{\nabla} \times \int d^3r' \frac{\vec{\nabla}' \times \vec{j}(\vec{r}')}{|\vec{r} - \vec{r}'|} \quad \text{transverse part}$$

$$\text{So } \vec{\nabla} \left(\frac{\partial \phi}{\partial t} \right) = 4\pi \vec{j}_{||}, \quad \text{and}$$

$$\square^2 A = \frac{4\pi}{c} \vec{j} - \frac{4\pi}{c} \vec{j}_{||} = \frac{4\pi}{c} \vec{j}_{\perp}$$

Transverse + Longitudinal Parts of vector functions

To prove the preceding claim, $\vec{f} = \vec{f}_{\parallel} + \vec{f}_{\perp}$, where $\vec{\nabla} \times \vec{f}_{\parallel} = 0$ and $\vec{\nabla} \cdot \vec{f}_{\perp} = 0$, we first desire to prove Helmholtz Theorem.

Helmholtz Theorem: For a vector function $\vec{f}(\vec{r})$ if one knows the divergence and curl of \vec{f} then one can ~~uniquely~~ uniquely determine \vec{f} itself.

That is, if

$$\vec{\nabla} \cdot \vec{f} = 4\pi D(\vec{r}) \quad \text{where } D(\vec{r}) \text{ is a known scalar function}$$

$$\vec{\nabla} \times \vec{f} = 4\pi \vec{C}(\vec{r}) \quad \text{where } \vec{C}(\vec{r}) \text{ is a known vector function}$$

~~Then one can solve for~~

And if well defined boundary conditions on \vec{f} are known (here we will assume $\vec{f}(\vec{r}) \rightarrow 0$ as $r \rightarrow \infty$), then there is a unique solution for $\vec{f}(\vec{r})$.

We prove this by construction!

Assume a solution of the form

$$\vec{f} = -\vec{\nabla}\phi + \vec{\nabla} \times \vec{W} \quad \text{where } \phi \text{ is a scalar and } \vec{W} \text{ a vector}$$

Now we show that we can find such a solution

First consider

$$\vec{\nabla} \cdot \vec{f} = -\nabla^2 \varphi + \vec{\nabla} \cdot (\vec{\nabla} \times \vec{W}) = -\nabla^2 \varphi + 0 = 4\pi D(\vec{r})$$

So $-\nabla^2 \varphi = 4\pi D(\vec{r})$ This is just Poisson's eqn we saw in electrostatics

Solution when $\varphi(\vec{r}) \rightarrow 0$ as $r \rightarrow \infty$ is given by

$$\varphi(\vec{r}) = \int d^3r' \frac{D(\vec{r}')}{|\vec{r} - \vec{r}'|}$$

Coulomb-like
integral solution

Now consider

$$\begin{aligned} \vec{\nabla} \times \vec{f} &= -\vec{\nabla} \times \vec{\nabla} \varphi + \vec{\nabla} \times (\vec{\nabla} \times \vec{W}) = 0 - \nabla^2 \vec{W} + \vec{\nabla} (\vec{\nabla} \cdot \vec{W}) \\ &= 4\pi \vec{C}(\vec{r}) \end{aligned}$$

Choose a gauge in which $\vec{\nabla} \cdot \vec{W} = 0$ (just like Coulomb gauge in magnetostatics)

$$\text{Then } -\nabla^2 \vec{W} = 4\pi \vec{C}(\vec{r})$$

$$\vec{W}(\vec{r}) = \int d^3r' \frac{\vec{C}(\vec{r}')}{|\vec{r} - \vec{r}'|}$$

just like solution for vector pot \vec{A} in magnetostatics

So we have constructed a solution

$$\vec{f}(\vec{r}) = -\vec{\nabla} \varphi + \vec{\nabla} \times \vec{W}$$

$$= -\vec{\nabla} \int d^3r' \frac{D(\vec{r}')}{|\vec{r} - \vec{r}'|} + \vec{\nabla} \times \int d^3r' \frac{\vec{C}(\vec{r}')}{|\vec{r} - \vec{r}'|}$$

$$\text{where } \vec{\nabla} \cdot \vec{f} = 4\pi D \text{ and } \vec{\nabla} \times \vec{f} = 4\pi \vec{C}$$

Note: For above solution to be well defined, the integrals must converge. They will converge if the "sources" $D(\vec{r})$ and $\vec{C}(\vec{r})$ are sufficiently "localized" in space, i.e. $D(\vec{r}) \rightarrow 0$, $\vec{C}(\vec{r}) \rightarrow 0$ sufficiently fast as $\vec{r} \rightarrow \infty$.

Now we show that the above solution is unique.

Suppose there was another solution \vec{g} such that

$$\vec{\nabla} \cdot \vec{g} = 4\pi D \quad \text{and} \quad \vec{\nabla} \times \vec{g} = 4\pi \vec{C}$$

Consider $\vec{h} \equiv \vec{f} - \vec{g}$ then

$$\vec{\nabla} \cdot \vec{h} = 0 \quad \text{and} \quad \vec{\nabla} \times \vec{h} = 0$$

Can show that only such \vec{h} that also has $\vec{h}(\vec{r}) \rightarrow 0$ as $\vec{r} \rightarrow \infty$ is $\vec{h} \equiv 0$, so $\vec{g} = \vec{f}$ and solution is unique.

As a consequence of Helmholtz Theorem we have also shown the following

① Any vector function \vec{f} can be written in terms of a scalar and vector potential

$$\vec{f} = -\vec{\nabla} \phi + \vec{\nabla} \times \vec{W}$$

or equivalently

(2) Any vector function \vec{F} can be written in terms of a curl free and a divergenceless part

$$\vec{F} = \vec{F}_{||} + \vec{F}_{\perp} \quad \text{where} \quad \vec{\nabla} \times \vec{F}_{||} = 0 \quad \text{curl free}$$

$$\vec{\nabla} \cdot \vec{F}_{\perp} = 0 \quad \text{divergenceless}$$

$$\text{where} \quad \left\{ \begin{array}{l} \vec{F}_{||}(\vec{r}) = -\vec{\nabla} \Phi(\vec{r}) = -\vec{\nabla} \int \frac{d^3 r'}{4\pi} \frac{[\vec{\nabla}' \cdot \vec{f}(\vec{r}')] }{|\vec{r} - \vec{r}'|} \\ \vec{F}_{\perp}(\vec{r}) = \vec{\nabla} \times \vec{W}(\vec{r}) = \vec{\nabla} \times \int \frac{d^3 r'}{4\pi} \frac{[\vec{\nabla}' \times \vec{f}(\vec{r}')] }{|\vec{r} - \vec{r}'|} \end{array} \right.$$

where in above we used $\bar{D}(\vec{r}') = \frac{1}{4\pi} \vec{\nabla}' \cdot \vec{f}(\vec{r}')$

$$\bar{C}(\vec{r}') = \frac{1}{4\pi} \vec{\nabla}' \times \vec{f}(\vec{r}')$$

~~where~~ $\vec{F}_{||}$ is called the longitudinal part of \vec{F}

\vec{F}_{\perp} is called the transverse part of \vec{F}

to understand the reason for these names, we need to consider the Fourier transform

Above can be generalized to situations where \vec{F} satisfies other boundary conditions, say has a specified value on a given boundary surface. One just replaces $\frac{1}{|\vec{r} - \vec{r}'|}$ by the appropriate Green's function — see more to come!

Returning to Ampere's law we see that the term

$$\vec{\nabla} \left(\frac{\partial \phi}{\partial t} \right) = -\vec{\nabla} \int d^3r' \left[\frac{\vec{\nabla}' \cdot \vec{j}(r', t)}{|\vec{r} - \vec{r}'|} \right]$$

$$= 4\pi \vec{j}_{||}(\vec{r}, t)$$

So Ampere's law becomes

$$\square^2 \vec{A} = \frac{4\pi}{c} \vec{j} - \frac{4\pi}{c} \vec{j}_{||}$$

$$\square^2 \vec{A} = \frac{4\pi}{c} \vec{j}_{\perp}$$

In Coulomb gauge, only the transverse part of \vec{j} serves as a source for \vec{A} .

\vec{A} describes the transverse modes, i.e. the EM radiation (recall in EM waves, the fields are always \perp direction of propagation)

ϕ describes the longitudinal modes

Coulomb gauge is not Lorentz invariant - if $\vec{\nabla} \cdot \vec{A} = 0$ in one inertial reference frame, in general $\vec{\nabla} \cdot \vec{A} \neq 0$ in another.

In Coulomb gauge, if $\rho = 0$, then $\phi = 0$ and

$$\vec{E} = -\frac{1}{c} \frac{\partial \vec{A}}{\partial t}$$

Electrostatic

$$-\nabla^2 \phi = 4\pi\rho \quad \text{with} \quad \vec{E} = -\vec{\nabla}\phi \quad (\text{statics only})$$

physical meaning of the potential ϕ

work done to move a test charge δq from \vec{r}_1 to \vec{r}_2 in presence of an electric field \vec{E} is

$$W_{12} = \int_{\vec{r}_1}^{\vec{r}_2} d\vec{l} \cdot \vec{F}$$

where \vec{F} is the force required to move the charge.

Since \vec{E} exerts a force $\delta q \vec{E}$ on the charge,

\vec{F} must counterbalance this electric force so

we can move the charge quasi-statically $\Rightarrow \vec{F} = -\delta q \vec{E}$

$$W_{12} = -\delta q \int_{\vec{r}_1}^{\vec{r}_2} d\vec{l} \cdot \vec{E} = \delta q \int_{\vec{r}_1}^{\vec{r}_2} d\vec{l} \cdot \vec{\nabla}\phi = \delta q [\phi(\vec{r}_2) - \phi(\vec{r}_1)]$$

$$\phi(\vec{r}_2) - \phi(\vec{r}_1) = \frac{W_{12}}{\delta q}$$

difference in potential between two points is the work per unit charge to move a test charge between the two points

Green's Functions - part I

$$-\nabla^2 \phi = 4\pi \rho$$

We already know that for a point charge q at position \vec{r}' ,
ie $\rho(\vec{r}) = q \delta(\vec{r} - \vec{r}')$, the solution to the above is

$$\phi(\vec{r}) = \frac{q}{|\vec{r} - \vec{r}'|} \quad \text{ie} \quad -\nabla^2 \left(\frac{1}{|\vec{r} - \vec{r}'|} \right) = 4\pi \delta(\vec{r} - \vec{r}')$$

We call the special solution for a point source
the Green function for the differential operator

$$-\nabla^2 G(\vec{r}, \vec{r}') = -4\pi \delta(\vec{r} - \vec{r}')$$

$G(\vec{r}, \vec{r}')$ gives the potential at position \vec{r} due
to a unit source at position \vec{r}'

Generally, one also has to specify a desired
boundary condition for the Green function on
the boundary of the system.

For the Coulomb solution for a point charge
the implicit boundary condition is that the
potential vanish infinitely far from the charges

$$G(\vec{r}, \vec{r}') \rightarrow 0 \quad \text{as} \quad |\vec{r} - \vec{r}'| \rightarrow \infty$$

boundary of the system is taken to infinity

If one knows the Green's function, then one can find the solution for any distribution of sources $\rho(\vec{r})$

$$\phi(\vec{r}) = \int d^3r' G(\vec{r}, \vec{r}') \rho(\vec{r}')$$

proof: $-\nabla^2 \phi = \int d^3r' [-\nabla^2 G(\vec{r}, \vec{r}')] \rho(\vec{r}')$

$$= \int d^3r' [4\pi \delta(\vec{r} - \vec{r}')] \rho(\vec{r}')$$
$$= 4\pi \rho(\vec{r})$$

We will return to concept of Green's function when we discuss solution of Poisson's eqn in a finite volume

We will also see Green's functions again when we discuss solution of the inhomogeneous wave equation.