

Helmholtz Theorem

Suppose $\vec{\nabla} \cdot \vec{E} = f(r)$ { for $\vec{r} \in V$
 $\vec{\nabla} \times \vec{E} = g(r)$ } f and g known functions
 $\vec{E}(\vec{r}) = \vec{h}(\vec{r})$ for $\vec{r} \in S$ the boundary
of V

Suppose we had two different solutions

\vec{E} and \vec{E}' to the above equations.

Then consider $\vec{G} = \vec{E} - \vec{E}'$.

\vec{G} must satisfy

$$\begin{aligned}\vec{\nabla} \cdot \vec{G} &= 0 & \text{for } \vec{r} \in V \\ \vec{\nabla} \times \vec{G} &= 0 \\ \vec{G} &= 0 & \text{for } \vec{r} \in S\end{aligned}$$

Now $\vec{\nabla} \times \vec{G} = 0 \Rightarrow \vec{G} = \vec{\nabla} \phi$ for some scalar potential ϕ .

$$\vec{\nabla} \cdot \vec{G} = 0 \Rightarrow \vec{\nabla}^2 \phi = 0 \quad \text{in } V \quad \left. \begin{array}{l} \Rightarrow \phi = 0 \\ \phi = 0 \quad \text{on } S \end{array} \right\}$$

by properties of harmonic function

for harmonic function $\phi(\vec{r}) = \frac{1}{4\pi R^2} \oint_S d\vec{a}' \phi(\vec{r}')$ where S is sphere of radius R centered at \vec{r}

$\Rightarrow \phi(\vec{r})$ can have no extremum inside a volume $V \Rightarrow$ all extrema must be on boundary of V . If $\phi = 0$ on boundary $\Rightarrow \phi_{\max} = \phi_{\min} = \phi = 0$

Magneto statics

Lorentz Force

~~Electricity~~
~~Magnetism~~

a charge q , in motion with velocity \vec{v} , feels the force

$$\vec{F} = q(\vec{E} + k_4 \vec{v} \times \vec{B}) \quad \leftarrow \text{Lorentz force}$$

\vec{B} is the magnetic field at the position of the charge.
 k_4 is a universal constant.

Just as the constant k_1 fixed the units of charge q ,
the constant k_4 can be viewed as fixing the units of B
magnetic field. By choosing the units of q and B
appropriately, we are free to choose any values for k_1 and k_4 .

Magnetic field \vec{B} is generated by moving charge.
A charge q' with velocity \vec{v}' ($v' \ll c$) located at
the origin $\vec{r}'=0$ produces a magnetic field at
position \vec{r} ,

holds only $\rightarrow \vec{B}(\vec{r}) = k_5 q' \frac{\vec{v}' \times \vec{r}}{r^3} = \frac{k_5}{k_1} \vec{v}' \times \vec{E}(\vec{r})$
nonrelativistically

k_5 is a universal constant. we will see that
it cannot be chosen independently of k_1 and k_4 .
(since k_1 fixed units of q , and k_4 fixed units of \vec{B} ,
there are no further new quantities whose units
could be adjusted to allow us to fix k_5 arbitrarily)

(11)

The force on a charge q at position \vec{r} , moving with velocity \vec{v} , due to a charge q' at the origin moving with velocity \vec{v}' is, in non-relativistic limit ($v, v' \ll c$),

$$\vec{F} = k_1 q q' \frac{\vec{r}}{r^3} + k_4 k_5 q q' \frac{\vec{v} \times (\vec{v}' \times \vec{r})}{r^3}$$

\vec{r}
Coulomb force

\vec{v}
magnetic analog of Coulomb force

The magnetic part is just the point charge equivalent of the Biot-Savart law for the force between current carrying wires. If we regard $q\vec{v} = \vec{I}$ as the current of charge q , and $q'\vec{v}' = \vec{I}'$ as the current of charge q' , then the magnetic force is $k_4 k_5 \frac{\vec{I} \times (\vec{I}' \times \vec{r})}{r^3}$, which is the Biot-Savart Law.

Re-write above force as

$$\vec{F} = k_1 \left(1 + \frac{k_4 k_5}{k_1} \vec{v} \times \vec{v}' \times \right) \frac{\vec{r}}{r^3} q q'$$

we see that $\left(\frac{k_4 k_5}{k_1} \right)$ has units of $(\text{velocity})^{-2}$

it must be independent of whatever convention one used to choose the units of q or B (ie independent of choices for k_1 ad k_4). Experimentally it is found that

$$\left(\frac{k_4 k_5}{k_1} \right) = \frac{1}{c^2}$$

c = speed of light
in vacuum

Continuum current density

For charges q_i at positions $\vec{r}_i(t)$ with $\vec{v}_i = \frac{d\vec{r}_i}{dt}$
we define the current density

$$\vec{j}(\vec{r}, t) = \sum_i q_i \vec{v}_i(t) \delta(\vec{r} - \vec{r}_i(t))$$

units of \vec{j} are (charge) ($\frac{\text{length}}{\text{time}}$) ($\frac{1}{\text{length}^3}$) = $\frac{(\text{charge})}{(\text{area} \cdot \text{time})}$

charge per unit area per unit time

For a surface S'

$$\int_S d\vec{a} \hat{n} \cdot \vec{j} = I \quad \begin{matrix} \text{current (charge per unit time)} \\ \text{passing through surface } S' \end{matrix}$$

Charge Conservation vol V bounded by surface S'

$$\frac{d}{dt} \int_V d^3r j(\vec{r}, t) = - \oint_S d\vec{a} \hat{n} \cdot \vec{j}$$

rate of change of total charge in V = $(-)$ charge flowing out of V
through S' per unit time

$$\text{use } \oint_S d\vec{a} \hat{n} \cdot \vec{j} = \int_V d^3r \nabla \cdot \vec{j} = - \int_V \frac{\partial \rho}{\partial t}$$

\Rightarrow local charge conservation

$$\boxed{\frac{\partial \rho}{\partial t} + \nabla \cdot \vec{j} = 0}$$

A static situation has $\frac{\partial \vec{B}}{\partial t} = 0$

\Rightarrow magnetostatics is defined by the condition $\vec{\nabla} \cdot \vec{J} = 0$

Differential formulation of Biot-Savart

For a set of charges q_i at \vec{r}_i we have

$$\vec{B}(\vec{r}) = \sum_i k_s q_i \vec{v}_i \times \frac{(\vec{r} - \vec{r}_i)}{|\vec{r} - \vec{r}_i|^3}$$

$$= k_s \int d^3 r' \vec{J}(\vec{r}') \times \frac{(\vec{r} - \vec{r}')}{|\vec{r} - \vec{r}'|^3}$$

$$= k_s \int d^3 r' \vec{J}(\vec{r}') \times \vec{\nabla} \left(\frac{1}{|\vec{r} - \vec{r}'|} \right)$$

$$\vec{B}(\vec{r}) = k_s \vec{\nabla} \times \left[\int d^3 r' \frac{\vec{J}(r')}{|\vec{r} - \vec{r}'|} \right]$$

where we used $\vec{\nabla} \times (\vec{A} \phi) = -\vec{A} \times \vec{\nabla} \phi$ when \vec{A} is indep of \vec{r}

$\Rightarrow \boxed{\vec{\nabla} \cdot \vec{B} = 0}$ since $\vec{\nabla} \cdot (\vec{\nabla} \times \vec{A}) = 0$ for any vector function \vec{A}

integral form $\oint da \vec{n} \cdot \vec{B} = 0$

$$\vec{\nabla} \times \vec{B} = k_s \vec{\nabla} \times \left[\vec{\nabla} \times \left(\int d^3 r' \frac{\vec{J}(r')}{|\vec{r} - \vec{r}'|} \right) \right]$$

$$\text{use } \vec{\nabla} \times (\vec{\nabla} \times \vec{A}) = \vec{\nabla} (\vec{\nabla} \cdot \vec{A}) - \vec{\nabla}^2 \vec{A}$$

(14)

$$\vec{\nabla} \times \vec{B} = k_5 \vec{\nabla} \left[\int d^3r' \vec{\nabla} \cdot \left(\frac{\vec{f}(\vec{r}')}{|\vec{r}-\vec{r}'|} \right) \right]$$

$$= -k_5 \int d^3r' \vec{f}(\vec{r}') \nabla^2 \left(\frac{1}{|\vec{r}-\vec{r}'|} \right)$$

$$\text{in the 2nd term, } \nabla^2 \left(\frac{1}{|\vec{r}-\vec{r}'|} \right) = -4\pi \delta(\vec{r}-\vec{r}')$$

$$\text{in the 1st term, } \vec{\nabla} \cdot \frac{\vec{f}(\vec{r}')}{|\vec{r}-\vec{r}'|} = \vec{f}(\vec{r}') \cdot \vec{\nabla} \left(\frac{1}{|\vec{r}-\vec{r}'|} \right)$$

$$= -\vec{f}(\vec{r}') \cdot \vec{\nabla}' \left(\frac{1}{|\vec{r}-\vec{r}'|} \right)$$

$$\text{since } \vec{\nabla} = -\vec{\nabla}'$$

$$\text{So, } \int d^3r' \vec{\nabla} \cdot \left(\frac{\vec{f}(\vec{r}')}{|\vec{r}-\vec{r}'|} \right) = - \int d^3r' \vec{f}(\vec{r}') \cdot \vec{\nabla}' \left(\frac{1}{|\vec{r}-\vec{r}'|} \right)$$

$$\text{integrate by parts} \quad = \int d^3r' \left(\vec{\nabla}' \cdot \vec{f}(\vec{r}') \right) \left(\frac{1}{|\vec{r}-\vec{r}'|} \right)$$

surface term $\rightarrow 0$ as

we take surface $\rightarrow \infty$

since $\vec{f} \rightarrow 0$ as $r \rightarrow \infty$

But for magnetostatics $\vec{\nabla} \cdot \vec{f} = 0 \Rightarrow$ only 2nd term remains

Thus, for magnetostatics

$$\vec{\nabla} \times \vec{B} = 4\pi k_5 \vec{f} \quad \text{Amperes' law}$$

$$\text{integral form} \quad \oint_C d\vec{l} \cdot \vec{B} = 4\pi k_5 \int_S da \hat{n} \cdot \vec{f}$$

\curvearrowleft curve bounding surface

Although above diff eqns were derived starting from a "non-relativistic"

point-charge Biot-Savart law, the accuracy remain true for all magneto static situations

(15)

So far electrostatics $\left\{ \begin{array}{l} \vec{\nabla} \cdot \vec{E} = 4\pi k_1 \rho \text{ Gauss} \\ \vec{\nabla} \times \vec{E} = 0 \end{array} \right.$

magnetostatics $\left\{ \begin{array}{l} \vec{\nabla} \cdot \vec{B} = 0 \\ \vec{\nabla} \times \vec{B} = 4\pi k_5 \vec{J} \text{ Amperes} \end{array} \right.$

charge current conservation $\frac{\partial \vec{B}}{\partial t} + \vec{\nabla} \cdot \vec{J} = 0$

Time Dependent situations

Faraday's law of induction $\vec{\nabla} \times \vec{E} \neq 0$! mag flux

$$\oint_C d\vec{l} \cdot \vec{E} = -k_3 \frac{\partial}{\partial t} \iint_S da \hat{n} \cdot \vec{B}$$

$$\hookrightarrow \text{emf around loop} \quad \text{emf} = -\frac{d\Phi}{dt}$$

voltage around closed loop \sim time rate of change of magnetic flux through loop

$$\Rightarrow \boxed{\vec{\nabla} \times \vec{E} = -k_3 \frac{\partial \vec{B}}{\partial t}}$$

k_3 is universal constant

Maxwell correction to Ampere's law

In our derivation of $\vec{\nabla} \times \vec{B} = 4\pi k_5 \vec{J}$
we used $\vec{\nabla} \cdot \vec{J} = 0$. This is only true for magnetostatics - it is not true in general

Alternatively, since $\vec{\nabla} \cdot (\vec{\nabla} \times \vec{B}) = 0$ always,
if Ampere's law was true, we would necessarily conclude that $\vec{\nabla} \cdot \vec{J} = 0$. But $\vec{\nabla} \cdot \vec{J} = -\frac{\partial \rho}{\partial t} \neq 0$ in general.