

## The Coulomb problem as a boundary value problem

Consider a conducting sphere of radius  $R$  with net charge  $q$ . (as  $R \rightarrow 0$  we get a point charge). What is  $\phi(\vec{r})$ ? What is  $\vec{E}(\vec{r})$ ?

### Review: Properties of conductors in electrostatics

- 1)  $\vec{E} = 0$  inside conductor - if  $\vec{E} \neq 0$  then a current  $\vec{j} = \sigma \vec{E}$  flows and it is not statics ( $\sigma$  is conductivity)
- 2)  $\rho = 0$  inside conductor - if  $\vec{E} = 0$  inside, then  $\vec{\nabla} \cdot \vec{E} = 4\pi\rho = 0$
- 3) Any net charge on the conductor must lie on the surface - follows from (2)
- 4)  $\phi = \text{constant}$  throughout conductor - if  $\vec{E} = 0$  then  $\vec{E} = -\vec{\nabla}\phi \Rightarrow \phi$  is constant
- 5) Just outside the conductor,  $\vec{E}$  is  $\perp$  to surface.
  - If  $\vec{E}$  has a component  $\parallel$  to surface then it exerts a force on electrons at the surface leading to a surface current - so would not be static

For conducting sphere,  $\rho = 0$  for  $r > R$  and  $r < R$   
all charge is on the surface  $\Rightarrow \nabla^2\phi = 0$  for  $\begin{cases} r > R \\ r < R \end{cases}$

spherical symmetry  $\Rightarrow$  expect spherically symmetric solution

$\Rightarrow \phi(\vec{r})$  depends only on  $r = |\vec{r}|$

⇒ Solve Laplace's eqn by writing  $\nabla^2$  in spherical coords.  
Only the radial terms do not vanish.

$$\nabla^2 \phi = \frac{1}{r^2} \frac{d}{dr} \left( r^2 \frac{d\phi}{dr} \right) = 0$$

$$r^2 \frac{d\phi}{dr} = -C_0 \quad \text{a constant}$$

$$\frac{d\phi}{dr} = -\frac{C_0}{r^2}$$

$$\phi(r) = \frac{C_0}{r} + C_1, \quad C_1 \text{ a constant}$$

"outside"  $r > R$   $\phi_{(r)}^{\text{out}} = \frac{C_0^{\text{out}}}{r} + C_1^{\text{out}}$

"inside"  $r < R$   $\phi_{(r)}^{\text{in}} = \frac{C_0^{\text{in}}}{r} + C_1^{\text{in}}$

solution "outside" does not necessarily go smoothly into the solution "inside" because of the charge layer at  $r=R$  that separates the two regions. We need to determine the constants  $C_0^{\text{in}}, C_0^{\text{out}}, C_1^{\text{in}}, C_1^{\text{out}}$  by applying boundary conditions corresponding to the physical situation.

- ① For  $r > R$ , assume  $\phi \rightarrow 0$  as  $r \rightarrow \infty$  - boundary condition at infinity

$$\Rightarrow C_1^{\text{out}} = 0$$

$$\phi_{(r)}^{\text{out}} = \frac{C_0^{\text{out}}}{r} \quad \text{recover the expected Coulomb form.}$$

2) For  $r < R$ .

i) we could use the fact that the region  $r < R$  is a conductor with  $\phi = \text{constant}$  to conclude  $C_0^{\text{in}} = 0$


ii) or, if we were dealing with a charged shell instead of a conductor, we could argue as follows:

no charge at origin  $r=0 \Rightarrow$  expect  $\phi$  should be finite at origin  $\Rightarrow C_0^{\text{in}} = 0$

So  $\phi^{\text{in}}(r) = C^{\text{in}}$  a constant

3) Now we need boundary condition at  $r=R$  where "inside" and "outside" meet.


Review: Electric field and potential at a surface charge layer



← a general surface  $S$  with surface charge density  $\sigma(\vec{r})$  for  $\vec{r}$  on  $S$ ,  $\sigma(\vec{r})da$  is total charge in area  $da$  on surface

i) Take "Gaussian pillbox" surface about point  $\vec{r}$  on the surface  $S'$

top and bottom areas of pill box  $da$



Gauss' Law in integral form  $\oint da \hat{n} \cdot \vec{E} = 4\pi Q_{\text{enclosed}}$

expect  $\vec{E}$  is finite  $\Rightarrow$  contribution from sides of pillbox vanish as  $dl \rightarrow 0$ .

$$\oint da \hat{n} \cdot \vec{E} = \int_{\text{top}} da \hat{n} \cdot \vec{E} + \int_{\text{bottom}} da \hat{n} \cdot \vec{E}$$

$$= (\hat{n}_{\text{top}} \cdot \vec{E}_{\text{top}} + \hat{n}_{\text{bottom}} \cdot \vec{E}_{\text{bottom}}) da \quad \text{since } da \text{ is small}$$

$\vec{E}_{\text{top}}$  is electric field at  $\vec{r}$  just above the surface  $S$

$\vec{E}_{\text{bottom}}$  is electric field at  $\vec{r}$  just below the surface  $S$


$\hat{n}_{\text{top}} = \hat{n}$  is outward normal on top

$\hat{n}_{\text{bottom}} = -\hat{n}$  is outward normal on bottom

$$\Rightarrow (\vec{E}_{\text{top}} - \vec{E}_{\text{bottom}}) \cdot \hat{n} da = 4\pi Q_{\text{enclosed}} = 4\pi \sigma(\vec{r}) da$$

$$(\vec{E}_{\text{top}} - \vec{E}_{\text{bottom}}) \cdot \hat{n} = 4\pi \sigma(\vec{r}) \quad \boxed{\text{discontinuity in normal component of } E}$$

ii) Take "Amperian loop"  $C$  at surface about point  $\vec{r}$ .



$\nabla \times \vec{E} = 0 \Rightarrow \oint_C d\vec{l} \cdot \vec{E} = 0$  since  $\vec{E}$  is finite at surface,  
if take sides  $dl' \rightarrow 0$  their contribution to integral vanishes

$$\oint_C d\vec{l} \cdot \vec{E} = \boxed{(\vec{E}_{\text{top}} - \vec{E}_{\text{bottom}}) \cdot d\vec{l} = 0}$$

where  $d\vec{l}$  is any infinitesimal tangent to the surface at  $\vec{r}$ .

$\Rightarrow$  tangential component of  $\vec{E}$  is continuous

combine above to write

$$\vec{E}^{\text{top}} - \vec{E}^{\text{bottom}} = 4\pi\sigma(F) \hat{m}$$

iii)  $\vec{E} = -\vec{\nabla}\phi \Rightarrow \phi(r_2) - \phi(r_1) = - \int_{r_1}^{r_2} d\vec{l} \cdot \vec{E}$

Take  $\vec{r}_2$  just above  $\vec{r}$  on surface  
 $\vec{r}_1$  just below  $\vec{r}$  on surface  $\} d\vec{l} \rightarrow 0$

Since  $\vec{E}$  is finite  $\Rightarrow \int d\vec{l} \cdot \vec{E} \rightarrow 0$

$$\Rightarrow \phi^{\text{top}} = \phi^{\text{bottom}}$$

potential  $\phi$  is continuous at surface charge layer

can rewrite (i) as

$$(-\vec{\nabla}\phi^{\text{top}} + \vec{\nabla}\phi^{\text{bottom}}) \cdot \hat{m} = 4\pi\sigma$$

$$-\frac{\partial\phi^{\text{top}}}{\partial m} + \frac{\partial\phi^{\text{bottom}}}{\partial m} = 4\pi\sigma$$

1 directional derivative of  $\phi$  in direction  $\hat{m}$

discontinuity in normal derivative of  $\phi$  at surface

Apply to conducting sphere

$$\phi \text{ continuous} \Rightarrow \phi^{\text{in}}(R) = \phi^{\text{out}}(R)$$

$$C_1^{\text{in}} = \frac{C_0^{\text{out}}}{R}$$

only one unknown left

normal derivative of  $\phi$  is discontinuous

$$-\frac{\partial \phi^{\text{top}}}{\partial n} + \frac{\partial \phi^{\text{bottom}}}{\partial n} = 4\pi\sigma$$

here  $\hat{n} = \hat{r}$  the radial direction

$$\left[ -\frac{d\phi^{\text{out}}}{dr} + \frac{d\phi^{\text{in}}}{dr} \right]_{r=R} = 4\pi\sigma$$

but  $\frac{d\phi^{\text{in}}}{dr} = 0$  as  $\phi^{\text{in}} = \text{constant}$

$$-\frac{d\phi^{\text{out}}}{dr} \Big|_{r=R} = 4\pi\sigma$$

charge  $q$  is uniformly distributed on surface at  $R$

$$-\frac{d}{dr} \left( \frac{C_0^{\text{out}}}{r} \right)_{r=R} = \frac{C_0^{\text{out}}}{R^2} = 4\pi\sigma = 4\pi \left( \frac{q}{4\pi R^2} \right) = \frac{q}{R^2}$$

$$\Rightarrow C_0^{\text{out}} = q, \quad C^{\text{in}} = \frac{C_0^{\text{out}}}{R} = \frac{q}{R}$$

$$\phi(r) = \begin{cases} \frac{q}{R} & r < R \text{ inside} \\ \frac{q}{r} & r > R \text{ outside} \end{cases}$$

$$\Rightarrow \vec{E} = -\vec{\nabla}\phi = -\frac{d\phi}{dr} = \begin{cases} 0 & r < R \text{ inside} \\ \frac{q}{r^2} & r > R \text{ outside} \end{cases}$$

we get familiar Coulomb solution!

Summary We can view the preceding solution for  $\phi_{\text{out}}$  as solving Laplace's eqn  $\nabla^2 \phi = 0$  subject to a specified boundary condition on the normal derivative of  $\phi$  at the boundary  $r=R$  of the "outside" region of the system.

Alternate problem:

Another physical situation would be to connect a conducting sphere to a battery that charges the sphere to a fixed voltage  $\phi_0$  (statvolts!) with respect to ground  $\phi=0$  at  $r \rightarrow \infty$ .

As before, outside the sphere  $\phi = \frac{C_0}{r}$

Now the boundary condition is to specify the value of  $\phi$  on the boundary of the outside region, i.e.

$$\phi(R) = \phi_0$$

$$\Rightarrow \frac{C_0}{R} = \phi_0, \quad C_0 = \phi_0 R$$

$$\phi(r) = \phi_0 \frac{R}{r}$$

(from preceding solution, we know that charging the sphere to voltage  $\phi_0$  (statvolts) induces a net charge  $q = \phi_0 R$  on it)

These two versions of the conducting sphere problem are examples of a more general boundary value problem

Solve  $\nabla^2\phi = 0$  in a given region of space subject to one of the following two types of boundary conditions on the boundary surfaces of the region

i) Neumann boundary condition

$\frac{\partial \phi}{\partial n}$  - normal derivative of  $\phi$  is specified on the boundary surface

ii) Dirichlet boundary condition

$\phi$  - value of  $\phi$  is specified on the boundary surfaces

If the boundary surfaces consist of disjoint pieces, it is possible to specify either (i) or (ii) on each piece separately to get a mixed boundary value problem.

## Some more problems

infinite conducting wire of radius  $R$  with line charge density  $\lambda$  = charge per unit length



$$\text{surface charge } \sigma = \frac{\lambda}{2\pi R}$$

x

Expect cylindrical symmetry  $\Rightarrow \phi$  depends only on cylindrical coord  $r$ .

$$\nabla^2 \phi = 0 \text{ for } r > R, r < R$$

use  $\nabla^2$  in cylindrical coords - only radial term non vanishing

$$\nabla^2 \phi = \frac{1}{r} \frac{d}{dr} \left( r \frac{d\phi}{dr} \right) = 0$$

$$r \frac{d\phi}{dr} = C_0 \text{ constant}$$

$$\frac{d\phi}{dr} = \frac{C_0}{r}$$

$$\phi(r) = C_0 \ln r + C_1 \text{ const}$$

note: one cannot now choose  $\phi \rightarrow 0$  as  $r \rightarrow \infty$ !

one needs to fix zero of  $\phi$  at some other radius. a convenient choice is  $r=R$ , but any other choice could also be made.

$$\begin{aligned}\phi^{\text{out}} &= C_0^{\text{out}} \ln r + C_1^{\text{out}} \\ \phi^{\text{in}} &= C_0^{\text{in}} \ln r + C_1^{\text{in}}\end{aligned}$$

$$\phi^{\text{in}} = \text{const in conductor} \Rightarrow C_0^{\text{in}} = 0$$

or  $\phi^{\text{in}}$  should not diverge as  $r \rightarrow 0 \Rightarrow C_0^{\text{in}} = 0$

$$\text{so } \phi^{\text{in}} = C_1^{\text{in}} \text{ constant}$$

boundary condition at  $r=R$

$$\left[ -\frac{d\phi^{\text{out}}}{dr} + \frac{d\phi^{\text{in}}}{dr} \right]_{r=R} = 4\pi\sigma$$

$$\Rightarrow -\frac{C_0^{\text{out}}}{R} = 4\pi\sigma = 4\pi \left( \frac{2}{2\pi R} \right) = \frac{2\lambda}{R}$$

$$C_0^{\text{out}} = -2\lambda$$

$$\phi^{\text{out}}(r) = -2\lambda \ln r + C_1^{\text{out}}$$

continuity of  $\phi$

$$\phi^{\text{in}}(R) = \phi^{\text{out}}(R) \Rightarrow C_1^{\text{in}} = -2\lambda \ln R + C_1^{\text{out}}$$

Remaining const  $C_1^{\text{out}}$  is not too important as it is just a common additive constant to both  $\phi^{\text{in}}$  and  $\phi^{\text{out}}$   $\rightarrow$  does not change  $\vec{E} = -\vec{\nabla}\phi$


If use the condition  $\phi(R)=0$  then we can solve for  $C_1^{\text{out}}$ .

$$0 = -2\lambda \ln R + C_1^{\text{out}} \Rightarrow C_1^{\text{out}} = 2\lambda \ln R$$

$$\Rightarrow \phi(r) = \begin{cases} -2\lambda \ln(r/R) & r \geq R \\ 0 & r < R \end{cases}$$

$\Rightarrow \vec{E}(\vec{r}) = \begin{cases} \frac{2\lambda}{r} \hat{r} & r \geq R \\ 0 & r < R \end{cases}$

infinite conducting half space



$$\nabla^2 \phi = \frac{d^2 \phi}{dx^2} = 0$$

$$\Rightarrow \begin{cases} \phi^>(x) = C_0^>x + C_1^> & x > 0 \\ \phi^<(x) = C_0^<x + C_1^< & x < 0 \end{cases}$$

for  $x < 0$ ,  $\phi = \text{const}$  in conductor  $\Rightarrow C_0^< = 0$

at  $x=0$ ,  $\phi$  continuous  $\Rightarrow \phi^<(0) = \phi^>(0)$   
 $C_1^< = C_1^>$

$\frac{d\phi}{dx}$  discontinuous  $\Rightarrow$

$$-\left. \frac{d\phi}{dx} \right|_{x=0}^> = 4\pi\sigma$$

$$C_0^> = -4\pi\sigma$$

$$\Rightarrow \phi(x) = \begin{cases} -4\pi\sigma x + C_1^> & x > 0 \\ C_1^> & x < 0 \end{cases}$$

const  $C_1^>$  does not change value of  $\vec{E}$

as for the wire, we cannot choose  $\phi \rightarrow 0$  as  $x \rightarrow \infty$ .  
 we can set  $\phi = 0$  at

$$-\vec{\nabla}\phi = \vec{E} = \begin{cases} 4\pi\sigma \hat{x} & x > 0 \\ 0 & x < 0 \end{cases}$$

### infinite charged plane

similar to previous problem, but now no conductor  
 at  $x < 0$ , just free space on both sides of the  
 charged plane at  $x = 0$ .

~~expect underdamped response by symmetry~~

$$\nabla^2\phi = \frac{d^2\phi}{dx^2} = 0 \Rightarrow \phi^> = c_0^>x + c_1^> \quad x > 0$$

$$\phi^< = c_0^<x + c_1^< \quad x < 0$$

continuity of  $\phi$  at  $x = 0$

$$\Rightarrow \phi^>(0) = \phi^<(0) \Rightarrow c_1^> = c_1^<$$

discontinuity of  $d\phi/dx$  at  $x = 0$

$$-\frac{d\phi^>}{dx} + \frac{d\phi^<}{dx} = 4\pi\sigma$$

$$-c_0^> + c_0^< = 4\pi\sigma$$

$$\text{Define } \bar{c}_0 = \frac{c_0^> + c_0^<}{2}$$

Then we can write

$$c_0^< = \bar{c}_0 + 2\pi\sigma$$

$$c_0^> = \bar{c}_0 - 2\pi\sigma$$

$$\phi = \begin{cases} -2\pi\sigma x + \bar{c}_0 x + c_i^> & x > 0 \\ 2\pi\sigma x + \bar{c}_0 x + c_i^> & x < 0 \end{cases}$$

$$\Rightarrow -\frac{d\phi}{dx} = \vec{E} = \begin{cases} (2\pi\sigma - \bar{c}_0) \hat{x} & x > 0 \\ (-2\pi\sigma - \bar{c}_0) \hat{x} & x < 0 \end{cases}$$

Const  $c_i^>$  does not effect  $\vec{E}$  - additive const to  $\phi$

$\bar{c}_0$  represents const uniform electric field  $-\bar{c}_0 \hat{x}$ ,  
that exists independently of the charged surface

If we assumed that all  $\vec{E}$  fields are just those  
arising from the plane, then we can set  $\bar{c}_0 = 0$ .  
Equivalently, if the plane is the only source of  $\vec{E}$ ,  
then we expect  $\phi$  depends only on  $|x|$  by symmetry.

$$\Rightarrow c_0^< = -c_0^> \text{ and again } \bar{c}_0 = 0. \text{ In this case}$$

$$\phi(x) = \begin{cases} -2\pi\sigma x & x > 0 \\ 2\pi\sigma x & x < 0 \end{cases}$$

(we also set  $c_i^> = 0$  here corresponding to  $\phi(0) = 0$ )

$$\vec{E}(x) = \begin{cases} 2\pi\sigma \hat{x} & x > 0 \\ -2\pi\sigma \hat{x} & x < 0 \end{cases}$$

$\vec{E}$  is constant <sup>but</sup> oppositely directed on  
either side of the charged plane

## Green's theorem, Uniqueness, Green function - part II

We want to show that the boundary value problem we described is well posed - i.e. there is a unique solution. We start by deriving Greens Theorem

$$\text{Consider } \int_V d^3r \vec{\nabla} \cdot \vec{A} = \oint_S da \hat{n} \cdot \vec{A} \quad \text{Gauss theorem}$$

$$\text{let } \vec{A} = \phi \vec{\nabla} \psi \quad \phi, \psi \text{ any two scalar functions}$$

$$\Rightarrow \vec{\nabla} \cdot \vec{A} = \phi \nabla^2 \psi + \vec{\nabla} \phi \cdot \vec{\nabla} \psi$$

$$\phi \vec{\nabla} \psi \cdot \hat{n} = \phi \frac{\partial \psi}{\partial n}$$

$$\Rightarrow \int_V d^3r (\phi \nabla^2 \psi + \vec{\nabla} \phi \cdot \vec{\nabla} \psi) = \oint_S da \phi \frac{\partial \psi}{\partial n} \quad \left. \right\} \text{Green's 1st identity}$$

$$\text{let } \phi \leftrightarrow \psi$$

$$\int_V d^3r (\psi \nabla^2 \phi + \vec{\nabla} \phi \cdot \vec{\nabla} \psi) = \oint_S da \psi \frac{\partial \phi}{\partial n}$$

subtract

$$\int_V d^3r (\phi \nabla^2 \psi - \psi \nabla^2 \phi) = \oint_S da \left( \phi \frac{\partial \psi}{\partial n} - \psi \frac{\partial \phi}{\partial n} \right) \quad \left. \right\} \text{Green's 2nd identity}$$

Apply Green's 2nd identity with  $\psi = \frac{1}{|\vec{r} - \vec{r}'|}$ ,

$\vec{r}'$  is integration variable,  $\phi$  is the scalar potential with  $\nabla^2 \phi = -4\pi\rho$ . Use  $\nabla^2 \psi = \nabla'^2 \psi = -4\pi \delta(\vec{r} - \vec{r}')$

$$\int_V d^3r' \left[ \phi(r') [-4\pi \delta(r - r')] - \left( \frac{1}{|\vec{r} - \vec{r}'|} \right) (-4\pi \rho(r')) \right]$$

$$= \oint_S da' \left[ \phi \frac{\partial}{\partial n'} \left( \frac{1}{|\vec{r} - \vec{r}'|} \right) - \frac{1}{|\vec{r} - \vec{r}'|} \frac{\partial \phi}{\partial n'} \right]$$

If  $\vec{r}$  lies within the volume  $V$ , then

$$(*) \quad \phi(\vec{r}) = \int_V d^3r' \frac{\rho(\vec{r}')}{|\vec{r}-\vec{r}'|} + \oint_S da' \left[ \frac{1}{4\pi} \frac{\partial \phi}{\partial n'} - \phi \frac{\partial}{\partial n'} \left( \frac{1}{|\vec{r}-\vec{r}'|} \right) \right]$$

Note: if  $\vec{r}$  lies outside the volume  $V$ , then

$$0 = \int_V d^3r' \frac{\rho(\vec{r}')}{|\vec{r}-\vec{r}'|} + \oint_S da' \left[ \frac{1}{4\pi} \frac{\partial \phi}{\partial n'} - \phi \frac{\partial}{\partial n'} \left( \frac{1}{|\vec{r}-\vec{r}'|} \right) \right]$$

potential from a  
surface charge density

$$\sigma = \frac{1}{4\pi} \frac{\partial \phi}{\partial n'}$$

potential from a  
surface dipole layer of  
dipole strength density

$$\frac{\phi}{4\pi}$$

From (\*), if  $S \rightarrow \infty$  and  $E \sim \frac{\partial \phi}{\partial n'} \rightarrow 0$  faster than  $\frac{1}{r}$ ,

then the surface integral vanishes and we recover

Coulombs law  $\phi(\vec{r}) = \int_V d^3r' \frac{\rho(\vec{r}')}{|\vec{r}-\vec{r}'|}$

(\*) gives the generalization of Coulombs law to a system  
with a finite boundary

For a charge free volume  $V$ , i.e.  $\rho(r) = 0$  in  $V$ ,  
the potential everywhere is determined by the  
potential and its normal derivative on the surface.

But one cannot in general freely specify both  
 $\phi$  and  $\frac{\partial \phi}{\partial n'}$  on the boundary surface since the  
resulting  $\phi$  from (\*) would not in general obey  
 $\nabla^2 \phi = 0$ .

Specifying both  $\phi$  ad  $\frac{\partial \phi}{\partial n}$  on surface is known as "Cauchy" boundary conditions — for Laplace's eqn, Cauchy b.c. overspecify the problem + a solution cannot in general be found.

### Uniqueness

If we have a system of charges in vol  $V$ , and either the potential  $\phi$ , or its normal derivative  $\frac{\partial \phi}{\partial n}$ , is specified on the surfaces of  $V$ , then there is a unique solution to Poisson's equation inside  $V$ . Specifying  $\phi$  is known as Dirichlet boundary conditions. Specifying  $\frac{\partial \phi}{\partial n}$  is known as Neumann boundary conditions.

proof: Suppose we had two solutions  $\phi_1$  ad  $\phi_2$ , both with  $-\nabla^2 \phi = 4\pi\rho$  inside  $V$ , ad obeying specified b.c. on surface of  $V$ .

Define  $U = \phi_2 - \phi_1 \rightarrow \nabla^2 U = 0$  inside  $V$

and  $U=0$  on surface  $S$  - for Dirichlet b.c.  
or  $\frac{\partial U}{\partial n} = 0$  on surface  $S$  - for Neumann b.c.

Use Green's 1st identity with  $\phi = \psi = U$

$$\int_V d^3r (U \nabla^2 U + \vec{\nabla} U \cdot \vec{\nabla} U) = \oint_S U \frac{\partial U}{\partial n}$$

$$\text{as } \nabla^2 U = 0$$

$$\text{as } \oint_S U \frac{\partial U}{\partial n} = 0$$

$$\Rightarrow \int_V d^3r |\vec{\nabla}u|^2 = 0 \Rightarrow \vec{\nabla}u = 0 \Rightarrow u = \text{const}$$

For Dirichlet b.c.,  $u=0$  on surface  $S$ , so const = 0  
and  $\phi_1 = \phi_2$ . Solution is unique

For Neumann b.c.,  $\phi_1$  and  $\phi_2$  differ only by an arbitrary constant. Since  $\vec{E} = -\vec{\nabla}\phi$ , the electric fields  $\vec{E}_1 = -\vec{\nabla}\phi_1$  and  $\vec{E}_2 = -\vec{\nabla}\phi_2$  are the same.

~~bullet~~ If boundary ~~state~~ surface  $S$  consists of several disjoint pieces, then solution is unique if specify  $\phi$  on some pieces and  $\frac{\partial\phi}{\partial n}$  on other pieces.

Solution of Poisson's equation with both  $\phi$  and  $\frac{\partial\phi}{\partial n}$  specified on the same surface  $S$  (Cauchy b.c.) does not in general exist, since specifying either  $\phi$  or  $\frac{\partial\phi}{\partial n}$  alone is enough to give a unique solution.