

When atoms have intrinsic magnetic moments due to electron spin, we can add these to  $\vec{M}$  in obvious way

When molecules are neutral,  $q_n = 0$ , the "bound current" is given by

$$\vec{j}_{\text{bound}} = \sum_n \langle \vec{j}_n \rangle = c \vec{\nabla} \times \vec{M} + \frac{\partial \vec{P}}{\partial t}$$

Note that the  $\frac{\partial \vec{P}}{\partial t}$  term is crucial to give conservation of bound charge

$$\begin{aligned} \vec{\nabla} \cdot \vec{j}_{\text{bound}} &= c \vec{\nabla} \cdot (\vec{\nabla} \times \vec{M}) + \vec{\nabla} \cdot \frac{\partial \vec{P}}{\partial t} \\ &= 0 + \frac{\partial}{\partial t} (\vec{\nabla} \cdot \vec{P}) \end{aligned}$$

$$= -\frac{\partial \rho_{\text{bound}}}{\partial t} \quad \text{where } \rho_{\text{bound}} = -\vec{\nabla} \cdot \vec{P} \text{ is bound charge density}$$

$$\text{So } \boxed{\vec{\nabla} \cdot \vec{j}_{\text{bound}} + \frac{\partial \rho_{\text{bound}}}{\partial t} = 0}$$

and bound charge is conserved.

Since total average charge must be conserved, i.e.

$$\vec{\nabla} \cdot \langle \vec{j}_0 \rangle - \frac{\partial \langle \rho_0 \rangle}{\partial t} = 0, \quad \text{and } \langle \vec{j}_0 \rangle = \vec{j} + \vec{j}_{\text{bound}}$$

↑  
free current


$$\langle \rho_0 \rangle = \rho + \rho_{\text{bound}}$$

↑  
free charge

$$\Rightarrow \boxed{\vec{\nabla} \cdot \vec{j} + \frac{\partial \rho}{\partial t} = 0}$$

Free charge is also conserved

At a surface of a magnetic material



$\hat{m}$  outward normal to surface

take  $\hat{z} \equiv \hat{dl} \times \hat{m}$  out of page

Amperian loop  $C$  bounding surface of area  $da$

$$\begin{aligned} c \int_S da \hat{z} \cdot (\nabla \times \vec{M}) &= \int_S da \hat{z} \cdot \vec{J}_{\text{bound}} = da \hat{z} \cdot \vec{J}_{\text{bound}} \\ &= (\hat{dl} \times \hat{m}) \cdot \vec{K}_{\text{bound}} \quad \text{as width of loop} \\ &\quad \rightarrow 0 \\ &= (\hat{m} \times \vec{K}_{\text{bound}}) \cdot d\vec{l} \end{aligned}$$

But by Stokes theorem

$$c \int_S da \hat{z} \cdot (\nabla \times \vec{M}) = c \int_C d\vec{l} \cdot \vec{M} = c d\vec{l} \cdot \vec{M} \quad \text{since width} \rightarrow 0$$

and  $\vec{M} = 0$  outside

$$\Rightarrow c d\vec{l} \cdot \vec{M} = (\hat{m} \times \vec{K}_{\text{bound}}) \cdot d\vec{l} \quad \text{for any } d\vec{l} \text{ in plane of surface}$$

$$\Rightarrow c \vec{M}_\pm = \hat{m} \times \vec{K}_{\text{bound}}$$

where  $\vec{M}_\pm$  is component of  $\vec{M}$  tangential to the surface (since  $\vec{K}_b$  is in plane of surface,  $\hat{m} \times \vec{K}$  is also entirely in the plane of the surface)

$$\begin{aligned} \Rightarrow c \hat{m} \times \vec{M}_\pm &= c \hat{m} \times \vec{M} = \hat{m} \times (\hat{m} \times \vec{K}_{\text{bound}}) \\ &= -\vec{K}_{\text{bound}} \end{aligned}$$

$$\Rightarrow \begin{cases} \vec{K}_{\text{bound}} = c \vec{M} \times \hat{m} \\ \vec{H}_{\text{bound}} = c \nabla \times \vec{M} \end{cases}$$

Total bound charge vanishes

$$Q_{\text{bound}} = \int_V d^3r \rho_{\text{bound}} + \int_S da \sigma_{\text{bound}}$$


$\uparrow$  vol of dielectric       $\leftarrow$  surface of dielectric

$$= \int_V d^3r -\vec{\nabla} \cdot \vec{P} + \int da \hat{n} \cdot \vec{P}$$

but by Gauss theorem  $\int_V d^3r \vec{\nabla} \cdot \vec{P} = \int da \hat{n} \cdot \vec{P}$

so  $Q_{\text{bound}} = -\int da \hat{n} \cdot \vec{P} + \int da \hat{n} \cdot \vec{P} = 0$

Total bound current vanishes



$\hat{n}$  is normal to crosssection  
 $\hat{n}'$  is normal to surface

total current flowing through  $S$  is

$$\int_S da \hat{n} \cdot \vec{j}_{\text{bound}} + \int_C dl \vec{K}_{\text{bound}} \cdot \hat{n}$$

$$= c \int_S da \hat{n} \cdot (\vec{\nabla} \times \vec{M}) + c \int_C dl \hat{n} \cdot (\vec{M} \times \hat{n}')$$

$$= c \int_C d\vec{l} \cdot \vec{M} + c \int_C dl (\hat{n}' \times \hat{n}) \cdot \vec{M}$$

$= -\hat{x}$  mit tangent,  $d\vec{l} = dl \hat{x}$

$$= c \int_C d\vec{l} \cdot \vec{M} - c \int_C dl \cdot \vec{M} = 0$$

## Macroscopic Maxwell Equations


$$\begin{aligned}\vec{\nabla} \cdot \vec{B} &= 0 & \vec{\nabla} \times \vec{E} + \frac{1}{c} \frac{\partial \vec{B}}{\partial t} &= 0 \\ \vec{\nabla} \times \vec{H} &= \frac{4\pi}{c} \vec{j} + \frac{1}{c} \frac{\partial \vec{D}}{\partial t} & \vec{\nabla} \cdot \vec{D} &= 4\pi \rho\end{aligned}$$

where  $\rho$  and  $\vec{j}$  are macroscopic charge + current densities  
do not include bound charges or currents

$$\begin{aligned}\vec{D} &= \vec{E} + 4\pi \vec{P} & \vec{P} & \text{is polarization density} \\ \vec{H} &= \vec{B} - 4\pi \vec{M} & \vec{M} & \text{is magnetization density}\end{aligned}$$

## Boundary conditions for statics

electrostatics: at surface of a dielectric, or at interface between two different dielectrics



$$\vec{\nabla} \times \vec{E} = 0 \Rightarrow \hat{t} \cdot \vec{E}_{\text{above}} = \hat{t} \cdot \vec{E}_{\text{below}}$$

tangential component  $\vec{E}$  is continuous

$$\vec{\nabla} \cdot \vec{D} = 4\pi \rho \Rightarrow \hat{n} \cdot (\vec{D}_{\text{above}} - \vec{D}_{\text{below}}) = 4\pi \sigma$$

normal component of  $\vec{D}$  jumps by  $4\pi \sigma$

magneto statics: at surface or interface of magnetic materials

$$\vec{\nabla} \cdot \vec{B} = 0 \Rightarrow \hat{n} \cdot \vec{B}_{\text{above}} - \hat{n} \cdot \vec{B}_{\text{below}}$$

normal component of  $\vec{B}$  is continuous

$$\vec{\nabla} \times \vec{H} = \frac{4\pi}{c} \vec{j} \Rightarrow \hat{t} \cdot (\vec{H}_{\text{above}} - \vec{H}_{\text{below}}) = \frac{4\pi}{c} (\vec{k} \times \hat{n}) \cdot \hat{t}$$


tangential component of  $\vec{H}$  jumps by  $\frac{4\pi}{c} \vec{k} \times \hat{n}$

if  $\sigma = 0$ , i.e. no free surface charge, then  $\hat{n} \cdot \vec{D}$  continuous

if  $\vec{k} = 0$ , i.e. no free surface current, then  $\hat{t} \cdot \vec{H}$  continuous

## Examples

① Uniformly polarized sphere of radius  $R$   $\vec{P} = P\hat{z}$



bound charge  $\rho_b = -\vec{\nabla} \cdot \vec{P} = 0$  as  $\vec{P}$  constant

$$\sigma_b = \hat{n} \cdot \vec{P} = \hat{r} \cdot \vec{P} = P \cos \theta$$

we saw earlier that a sphere with surface charge  $\sigma(\theta) = \sigma_0 \cos \theta$  gives an electric field like a pure dipole for  $r > R$ , and is constant for  $r < R$ .

$$\vec{E}(\vec{r}) = \begin{cases} \left( \frac{4}{3} \pi R^3 P \right) \left[ \frac{2 \cos \theta \hat{r} + \sin \theta \hat{\theta}}{r^3} \right] & r > R \\ -\frac{4\pi P}{3} \hat{z} & r < R \end{cases}$$

total dipole moment is  $\vec{p} = \frac{4}{3} \pi R^3 P \hat{z}$

check behavior at boundary

Tangential component  $\vec{E}$

$$\vec{E}_{\text{above}}^{\perp} = \left( \frac{4}{3} \pi R^3 P \right) \frac{\sin \theta \hat{\theta}}{R^3} = \frac{4\pi P}{3} \sin \theta \hat{\theta}$$

$$\vec{E}_{\text{below}}^{\perp} = -\frac{4\pi P}{3} (\hat{z} \cdot \hat{\theta}) \hat{\theta} = \frac{4\pi P}{3} \sin \theta \hat{\theta}$$

$\Rightarrow$  Tangential component  $\vec{E}$  is continuous

normal component of  $\vec{D}$

outside:  $\vec{P} = 0 \Rightarrow \vec{D} = \vec{E}$

$$\Rightarrow \hat{n} \cdot \vec{D} = \hat{r} \cdot \vec{E} = \left( \frac{4}{3} \pi R^3 P \right) \frac{2 \cos \theta \hat{r}}{R^3} = \frac{8}{3} \pi P \cos \theta$$

$$\text{inside: } \vec{E} = -\frac{4\pi\vec{P}}{3} \Rightarrow \vec{P} = -\frac{3}{4\pi}\vec{E}$$

$$\vec{D} = \vec{E} + 4\pi\vec{P} = \vec{E} - 3\vec{E} = -2\vec{E} = \frac{8\pi P}{3}\hat{z}$$

$$\hat{m} \cdot \vec{D} = \hat{r} \cdot \left( \frac{8\pi P}{3}\hat{z} \right) = \frac{8\pi P}{3} \cos\theta$$

$\Rightarrow$  normal component  $\vec{D}$  is continuous

Note: normal component of  $\vec{E}$  should jump by  $4\pi\sigma_b = 4\pi P \cos\theta$   
inside

to check this:  $\hat{m} \cdot \vec{E} = \hat{r} \cdot \left( -\frac{4}{3}\pi P \hat{z} \right) = -\frac{4}{3}\pi P \cos\theta$

$$\hat{m} \cdot (\vec{E}^{\text{above}} - \vec{E}^{\text{below}}) = \frac{8}{3}\pi P \cos\theta + \frac{4}{3}\pi P \cos\theta$$

$$= \frac{12}{3}\pi P \cos\theta = 4\pi P \cos\theta = \sigma_b(\theta)$$

② Uniformly magnetized sphere of radius  $R$   $\vec{M} = M \hat{z}$



bound current  $\vec{j}_b = c \vec{\nabla} \times \vec{M} = 0$  as  $\vec{M}$  constant  
 $\vec{K}_b = c \vec{M} \times \hat{n} = cM (\hat{z} \times \hat{r})$   
 $= cM \sin \theta \hat{\phi}$

We saw earlier that a sphere with surface current  $\vec{K}_b = K_0 \sin \theta \hat{\phi}$  gives a magnetic field that is pure dipole for  $r > R$ , and is constant for  $r < R$ .

$$\vec{B}(\vec{r}) = \begin{cases} \left( \frac{4}{3} \pi R^3 M \right) \left[ \frac{2 \cos \theta \hat{r} + \sin \theta \hat{\theta}}{r^3} \right] & r > R \\ \frac{8}{3} \pi M \hat{z} & r < R \end{cases}$$

total dipole moment is  $\vec{m} = \frac{4}{3} \pi R^3 \vec{M}$

check behavior at boundary

normal component of  $\vec{B}$

$$\hat{m} \cdot \vec{B}^{\text{above}} = \hat{r} \cdot \vec{B}^{\text{above}} = \frac{8}{3} \pi M \cos \theta$$

$$\hat{m} \cdot \vec{B}^{\text{below}} = \hat{r} \cdot \vec{B}^{\text{below}} = \frac{8}{3} \pi M (\hat{r} \cdot \hat{z}) = \frac{8}{3} \pi M \cos \theta$$


$\Rightarrow$  normal component of  $\vec{B}$  is continuous

tangential component of  $\vec{H}$

outside:  $\vec{M} = 0 \Rightarrow \vec{H} = \vec{B}$

$$\vec{H}_{\text{above}}^t = \left(\frac{4}{3} \pi M\right) \sin \theta \hat{\theta}$$

inside:  $\vec{H} = \vec{B} - 4\pi \vec{M} = \vec{B} - 4\pi \left(\frac{3}{8\pi} \vec{B}\right) = \vec{B} - \frac{3}{2} \vec{B} = -\frac{1}{2} \vec{B}$   
 $= -\frac{4\pi M}{3} \hat{z}$



so  $\vec{H}_{\text{below}}^t = -\frac{4\pi}{3} M (\hat{z} \cdot \hat{\theta}) = \frac{4\pi}{3} M \sin \theta \hat{\theta}$

$\Rightarrow$  tangential component  $\vec{H}$  is continuous

Note: tangential component  $\vec{B}$  should jump by  $\frac{4\pi}{c} \vec{K}_b \times \hat{m} = 4\pi M \sin \theta \hat{\theta}$

inside:

to check:  $\vec{B}_{\text{below}}^t = \frac{8}{3} \pi M (\hat{z} \cdot \hat{\theta}) \hat{\theta} = -\frac{8}{3} \pi M \sin \theta \hat{\theta}$

$$\vec{H}_{\text{above}}^t = \vec{B}_{\text{above}}^t \Rightarrow \vec{B}_{\text{above}}^t - \vec{B}_{\text{below}}^t = \frac{4}{3} \pi M \sin \theta \hat{\theta} + \frac{8}{3} \pi M \sin \theta \hat{\theta}$$

$$= 4\pi M \sin \theta \hat{\theta} = \frac{4\pi}{c} \vec{K}_b$$