

For  $T \leq T_c$ ,  $z = 1$  so  $\frac{dz}{dT} = 0$  and only 1st term remains

$$\frac{T}{\lambda^3} \propto T^{5/2} \text{ so } \frac{d}{dT} \left( \frac{T}{\lambda^3} \right) = \frac{5}{2} \left( \frac{T}{\lambda^3} \right) \frac{1}{T} = \frac{5}{2} \frac{1}{\lambda^3}$$

↙  $z=1$  here for all  $T \leq T_c$

$$\Rightarrow \frac{C_V}{Nk_B} = \frac{3}{2} \nu \left( \frac{5}{2} \frac{1}{\lambda^3} \right) g_{5/2}(1) = \frac{15}{4} g_{5/2}(1) \frac{\nu}{\lambda^3}$$

$$= \frac{15}{4} g_{5/2}(1) \nu \left( \frac{2\pi m k_B T}{h^2} \right)^{3/2}$$

Note, at  $T_c$ ,  $n = \frac{g_{3/2}(1)}{\lambda_c^3}$ , and  $\nu = \frac{1}{n}$

$$\frac{C_V(T_c)}{Nk_B} = \frac{15}{4} \frac{g_{5/2}(1)}{g_{3/2}(1)} = \frac{15}{4} \frac{1.341}{2.612} = 1.925 \leftarrow \text{this is larger than the classical ideal gas value of } \frac{3}{2}$$

So  $\boxed{\frac{C_V}{Nk_B} = 1.925 \left( \frac{T}{T_c} \right)^{3/2} \quad T \leq T_c}$

For  $T \geq T_c$ ,  $z$  varies with  $T$  and we need to evaluate the 2nd term as well

1st term gives  $\frac{15}{4} g_{5/2}(z(T)) \frac{\nu}{\lambda^3}$  ↙ here  $z$  depends on  $T$  for  $T > T_c$

2nd term: from Pathria Appendix D Eq(10),

$$z \frac{d}{dz} [g_\nu(z)] = g_{\nu-1}(z)$$

$$\Rightarrow \frac{d g_{5/2}}{dz} \frac{dz}{dT} = g_{3/2} \frac{1}{z} \frac{dz}{dT}$$

To find  $\frac{1}{z} \frac{dz}{dT}$  consider our earlier result for the density when  $T > T_c$ :

$$n = \frac{g_{3/2}(z)}{\lambda^3} \quad \leftarrow \text{determines } z(T) \text{ for fixed } n$$

$$\text{for } n \text{ fixed} \Rightarrow 0 = \frac{dn}{dT} = \frac{d}{dT} \left( \frac{1}{\lambda^3} \right) g_{3/2} + \frac{1}{\lambda^3} \frac{dg_{3/2}}{dz} \frac{dz}{dT}$$

$$0 = \frac{3}{2} \frac{1}{\lambda^3 T} g_{3/2} + \frac{1}{\lambda^3} g_{1/2} \frac{1}{z} \frac{dz}{dT}$$

$$\Rightarrow \frac{1}{z} \frac{dz}{dT} = -\frac{3}{2} \frac{g_{3/2}}{g_{1/2}} \frac{1}{T}$$

$$\frac{C_V}{Nk_B} = \frac{15}{4} g_{5/2}(z) \frac{v}{\lambda^3} + \frac{3}{2} v \frac{T}{\lambda^3} g_{3/2}(z) \left( -\frac{3}{2} \right) \frac{g_{3/2}(z)}{g_{1/2}(z)} \frac{1}{T}$$

$$\text{use } n = \frac{1}{v} = \frac{g_{3/2}(z)}{\lambda^3} \Rightarrow \frac{v}{\lambda^3} = \frac{1}{g_{3/2}(z)}$$

$$\boxed{\frac{C_V}{Nk_B} = \frac{15}{4} \frac{g_{5/2}(z)}{g_{3/2}(z)} - \frac{9}{4} \frac{g_{3/2}(z)}{g_{1/2}(z)} \quad T > T_c}$$

$$\text{Note } g_{1/2}(1) = \sum_{\ell=1}^{\infty} \frac{1}{\ell^{1/2}} \rightarrow \infty$$

So as  $T \rightarrow T_c^+$  from above, and  $z \rightarrow 1$

$$\frac{C_V}{Nk_B}(T_c^+) = \frac{15}{4} \frac{g_{5/2}(1)}{g_{3/2}(1)} - \frac{9}{4} \frac{g_{3/2}(1)}{\infty} = \frac{15}{4} \frac{1.341}{2.612} = 1.925$$

$$\Rightarrow \boxed{C_V \text{ is continuous at } T_c}$$

Finally we want to show that although  $C_V$  is continuous at  $T_c$ ,  $\frac{dC_V}{dT}$  is discontinuous

For  $T \leq T_c$   $\frac{C_V}{Nk_B} = 1.925 \left( \frac{T}{T_c} \right)^{3/2}$

$$\frac{d}{dT} \left( \frac{C_V}{Nk_B} \right) = \frac{3}{2} (1.925) \left( \frac{T}{T_c} \right)^{1/2} \frac{1}{T_c} = 2.89 \left( \frac{T}{T_c} \right)^{1/2} \frac{1}{T_c}$$

so slope at  $T_c^-$  (just below  $T_c$ )

w:  $\boxed{\frac{d}{dT} \left( \frac{C_V}{Nk_B} \right) = \frac{2.89}{T_c}, \quad T = T_c^-}$

For  $T > T_c$

$$\frac{C_V}{Nk_B} = \frac{15}{4} \frac{g_{5/2}(z)}{g_{3/2}(z)} - \frac{9}{4} \frac{g_{3/2}(z)}{g_{1/2}(z)}$$

$$\frac{d}{dT} \left( \frac{C_V}{Nk_B} \right) = \frac{15}{4} \frac{g_{3/2} \frac{dg_{5/2}}{dz} \frac{dz}{dT} - g_{5/2} \frac{dg_{3/2}}{dz} \frac{dz}{dT}}{(g_{3/2}(z))^2}$$

$$- \frac{9}{4} \frac{g_{1/2} \frac{dg_{3/2}}{dz} \frac{dz}{dT} - g_{3/2} \frac{dg_{1/2}}{dz} \frac{dz}{dT}}{(g_{1/2}(z))^2}$$

$$= \frac{1}{z} \frac{dz}{dT} \left\{ \frac{15}{4} \left( \frac{g_{3/2}^2 - g_{5/2} g_{1/2}}{g_{3/2}^2} \right) - \frac{9}{4} \left( \frac{g_{1/2}^2 - g_{3/2} g_{-1/2}}{g_{1/2}^2} \right) \right\}$$

use  $\frac{1}{z} \frac{dz}{dT} = -\frac{3}{2} \frac{g_{3/2}}{g_{1/2}} \frac{1}{T}$  as found earlier

$$\frac{d}{dT} \left( \frac{C_V}{Nk_B} \right) = -\frac{3}{8T} \frac{g_{3/2}}{g_{1/2}} \left\{ 15 \left( 1 - \frac{g_{5/2} g_{1/2}}{g_{3/2}^2} \right) - 9 \left( 1 - \frac{g_{3/2} g_{-1/2}}{g_{1/2}^2} \right) \right\}$$

Now as  $T \rightarrow T_c^+$  from above,  $z \rightarrow 1$ , we have

$g_{5/2}(1)$  and  $g_{3/2}(1)$  are finite, but  $g_{1/2}(1)$  and

$g_{-1/2}(1) \rightarrow \infty$

$\Rightarrow$  at  $T_c^+$

$$\frac{d}{dT} \left( \frac{C_V}{Nk_B} \right) = \frac{45}{8T_c} \frac{g_{5/2}(1)}{g_{3/2}(1)} - \frac{27}{8T_c} \frac{g_{3/2}(1)^2}{g_{1/2}(1)} \frac{g_{-1/2}(1)}{g_{1/2}(1)}$$

Now from Pathria Appendix D Eq(8)

$$g_\nu(1) = \lim_{a \rightarrow 0} \frac{\Gamma(1-\nu)}{a^{1-\nu}}$$

$$\text{So } \frac{g_{-1/2}(1)}{g_{3/2}(1)} = \lim_{a \rightarrow 0} \frac{\Gamma(3/2)}{a^{3/2}} \left( \frac{a^{1/2}}{\Gamma(1/2)} \right)^3 = \frac{\Gamma(3/2)}{[\Gamma(1/2)]^3}$$

$$= \frac{\frac{1}{2} \pi^{1/2}}{\pi^{3/2}} = \frac{1}{2\pi}$$

$$\text{since } \Gamma(1/2) = \sqrt{\pi}$$

$$\Gamma(3/2) = \frac{1}{2} \sqrt{\pi}$$


$$\frac{d}{dT} \left( \frac{C_V}{Nk_B} \right) = \frac{45}{8} \frac{1.341}{2.612} \frac{1}{T_c} - \frac{27}{8} \left( \frac{1}{2\pi} \right)^2 \frac{1}{T_c}$$

$$= \frac{2.89}{T_c} - \frac{3.66}{T_c} = -\frac{0.77}{T_c}$$

$$\boxed{\frac{d}{dT} \left( \frac{C_V}{Nk_B} \right) = -\frac{0.77}{T_c}, \quad T = T_c^+}$$

slope of  $C_V$   
is discontinuous at  
 $T_c$ .

$C_V$  has a cusp at  $T_c$



## Entropy

For single species gas we had for Gibbs free energy

$$G = N\mu$$

$$\text{Also } G = E - TS + pV$$

(since  $G$  is Legendre transform of  $E$  with respect to  $S$  and  $V$ )

$$\Rightarrow N\mu = E - TS + pV$$

$$\text{or } S = \frac{E + pV - N\mu}{T}$$

$$\frac{S}{Nk_B} = \frac{E + pV}{Nk_B T} - \frac{\mu}{k_B T}$$

$$\text{we had earlier } E = \frac{3}{2} pV \Rightarrow pV = \frac{2}{3} E$$

$$\frac{S}{Nk_B} = \frac{5}{3} \frac{E}{N} \frac{1}{k_B T} - \frac{\mu}{k_B T}$$

$$z = e^{M/k_B T}, \quad z = 1 \text{ for } T < T_c$$

we had earlier  $\frac{E}{N} = \frac{3}{2} \frac{k_B T}{\lambda^3} g_{5/2}(z)$

and  $n = \frac{1}{v} = \frac{g_{3/2}(z)}{\lambda^3}$  for  $T > T_c$

$$\Rightarrow \frac{S}{N k_B} = \frac{5}{2} \frac{v}{\lambda^3} g_{5/2}(z) - \ln z = \begin{cases} \frac{5}{2} \frac{g_{5/2}(z)}{g_{3/2}(z)} - \ln z, & T > T_c \\ \frac{5}{2} \frac{v}{\lambda^3} g_{5/2}(1), & T \leq T_c \end{cases}$$

Note: For  $T \leq T_c$  we had that the density of the "normal" a density  $n_0 = n - \frac{g_{3/2}(1)}{\lambda^3}$  in

the condensate, and a density  $\frac{g_{3/2}(1)}{\lambda^3}$  in the "normal" state (i.e. the density of excited particles)  $\equiv n_n$

$$\Rightarrow \text{for } T \leq T_c, \quad \frac{S}{N k_B} = \frac{5}{2} \left( \frac{n_n}{n} \right) \frac{g_{5/2}(1)}{g_{3/2}(1)} \rightarrow 0 \text{ as } T \rightarrow 0$$

We can imagine that each normal particle carries

entropy  $\frac{5}{2} k_B \frac{g_{5/2}(1)}{g_{3/2}(1)}$ . The entropy at  $T < T_c$  /per particle

is just the ~~for~~ above entropy per "normal" particle times the fraction of normal particles.

$\Rightarrow$  normal particles carry the entropy  
condensate has zero entropy

entropy difference per particle between normal state

and condensed state is  $\frac{\Delta S}{N} = \frac{5}{2} k_B \frac{g_{5/2}(1)}{g_{3/2}(1)}$

latent heat of condensation

$$L = T \Delta S = \frac{5}{2} k_B T \frac{g_{5/2}(1)}{g_{3/2}(1)}$$

energy released upon converting one normal particle to one condensate particle.

⇒ mixed phase is like coexistence region of a 1st order phase transition (like water ↔ ice ~~3~~ - need to remove energy to turn water to ice)

⇒ "two fluid" model of mixed region