


Liouville's theorem

The concept of density matrix will soon be expanded beyond the particular example of the microcanonical ensemble. It can also be generalized to non-equilibrium situations. $\rho(q_i, p_i, t)$
We therefore want to see what general condition ρ must satisfy in order that $\frac{\partial \rho}{\partial t} = 0$, i.e. steady-state

Consider an initial density ρ of points in phase space. As the systems represented by these initial points evolve in time, their trajectories give the density $\rho(t)$ at later times. Think of the points in ρ like particles in a fluid.



The probability density ρ must obey a local conservation equation

$$\frac{\partial \rho}{\partial t} + \vec{\nabla} \cdot (\rho \vec{v}) = 0$$

Where \vec{v} is the "velocity" vector of the probability "current" $\rho \vec{v}$, that tells how the points in ρ flow in phase space.

\vec{v} is the $6N$ dimensional vector $(\dot{q}_1, \dots, \dot{q}_N, \dot{p}_1, \dots, \dot{p}_N)$

and

$$\begin{aligned} \vec{\nabla} \cdot (\rho \vec{v}) &= \sum_{i=1}^{3N} \left[\frac{\partial}{\partial q_i} (\rho \dot{q}_i) + \frac{\partial}{\partial p_i} (\rho \dot{p}_i) \right] \\ &= \sum_{i=1}^{3N} \left[\frac{\partial \rho}{\partial q_i} \dot{q}_i + \rho \frac{\partial \dot{q}_i}{\partial q_i} + \frac{\partial \rho}{\partial p_i} \dot{p}_i + \rho \frac{\partial \dot{p}_i}{\partial p_i} \right] \end{aligned}$$

$$\vec{\nabla} \cdot (p \vec{v}) = \sum_i \left[\frac{\partial p}{\partial g_i} \dot{g}_i + \frac{\partial p}{\partial p_i} \dot{p}_i \right] + p \left[\frac{\partial \dot{g}_i}{\partial g_i} + \frac{\partial \dot{p}_i}{\partial p_i} \right]$$

use $\dot{g}_i = \frac{\partial H}{\partial p_i}$ $\dot{p}_i = -\frac{\partial H}{\partial g_i}$

$$\Rightarrow \frac{\partial \dot{g}_i}{\partial g_i} = \frac{\partial H}{\partial p_i \partial g_i} \quad \frac{\partial \dot{p}_i}{\partial p_i} = -\frac{\partial H}{\partial g_i \partial p_i}$$

so $\frac{\partial \dot{g}_i}{\partial g_i} + \frac{\partial \dot{p}_i}{\partial p_i} = 0$

and

$$\vec{\nabla} \cdot (p \vec{v}) = \sum_i \left\{ \frac{\partial p}{\partial g_i} \frac{\partial H}{\partial p_i} - \frac{\partial p}{\partial p_i} \frac{\partial H}{\partial g_i} \right\}$$

$$= [p, H] \quad \leftarrow \text{defines the "Poisson bracket"}$$

so

$$\frac{\partial p}{\partial t} + [p, H] = 0$$

$$\approx \frac{\partial p}{\partial t} + \sum_{i=1}^{3N} \left\{ \frac{\partial p}{\partial g_i} \frac{dg_i}{dt} + \frac{\partial p}{\partial p_i} \frac{dp_i}{dt} \right\} = \frac{dp}{dt} = 0$$

↑
total time derivative

$\frac{dp}{dt}$ is also called the convective derivative. It is how p changes in time if one moves along with "particles" (ie the systems on their trajectories)

$\frac{dp}{dt} = 0 \Rightarrow$ density in phase space is ~~invariant~~ ^{constant} in time as it flows - like an incompressible fluid.

density of points at t_0 = density at t

Equilibrium requires a stronger condition, namely $\frac{\partial \rho}{\partial t} = 0$,
 so that ensemble averages will not vary in time.

$$\frac{\partial \rho}{\partial t} = 0 \Rightarrow [\rho, H] = 0$$

$$[\rho, H] = \sum_i \left[\frac{\partial \rho}{\partial q_i} \frac{\partial H}{\partial p_i} - \frac{\partial \rho}{\partial p_i} \frac{\partial H}{\partial q_i} \right]$$

we see that $[\rho, H] = 0$ if $\rho(q_i, p_i)$ depends
 on q_i, p_i only via the function $H[q_i, p_i]$, i.e.

$$\rho = \rho(H(q_i, p_i)). \text{ Then } \frac{\partial \rho}{\partial q_i} = \frac{\partial \rho}{\partial H} \frac{\partial H}{\partial q_i}, \quad \frac{\partial \rho}{\partial p_i} = \frac{\partial \rho}{\partial H} \frac{\partial H}{\partial p_i}$$

$$[\rho, H] = \sum_i \left[\frac{\partial \rho}{\partial H} \frac{\partial H}{\partial q_i} \frac{\partial H}{\partial p_i} - \frac{\partial \rho}{\partial H} \frac{\partial H}{\partial p_i} \frac{\partial H}{\partial q_i} \right] = 0$$

so $\rho(q_i, p_i)$ must be constant on constant
 energy surfaces, if ρ is to describe equilibrium.

We already saw one example

microcanonical ensemble $\rho(q_i, p_i) \sim \delta(H(q_i, p_i) - E)$

another choice later will be

canonical ensemble $\rho(q_i, p_i) \sim e^{-H(q_i, p_i)/k_B T}$

Microcanonical Ensemble and Entropy

We saw that the microcanonical ensemble, at energy E , assigned equal weight to all systems on the surface in phase space of constant energy $H[q_i, p_i] = E$.

To count the number of such states on the energy surface we define the "density of states"

$$g(E) \equiv \frac{\int dq_i dp_i \delta(H[q_i, p_i] - E)}{h^{3N}}$$

where h is a constant with units of $q_i p_i$.

h^{3N} represents the volume of phase space occupied by one "state". Classically, h is totally arbitrary so our thermodynamic results should not depend on it. Quantum mechanically, we will see that h turns out to be Planck's constant.

At this stage, the factor $\frac{1}{h^{3N}}$ is introduced so that $g(E)$ has the units of $1/\text{energy}$.

We can now define the number of states in a shell of thickness Δ about the energy surface E .

$$\Omega(E, V, N) = \int_{E-\frac{\Delta}{2}}^{E+\frac{\Delta}{2}} dE' g(E')$$


Ω is a pure number. Again, Δ is arbitrary, but

$$\frac{E}{N} \ll \Delta \ll E$$

assumed to be ~~small~~. It can be thought of as representing the finite accuracy with which one knows E . Our thermodynamic results should not depend on Δ . [both h and Δ are introduced so that ~~the~~ ~~dimension~~ of Ω is a dimensionless pure number that we can think of as being the ~~total~~ number of microscopic states occupied in the microcanonical ensemble at E]

Example: Compute Ω for the ideal gas of non interacting particles, confined to a volume V

$$H = \sum_i \frac{p_i^2}{2m}$$



$$\begin{aligned} g(E) &= \int \frac{d^3r_i}{h^{3N}} \int d^3p_i \delta\left(\sum_i \frac{p_i^2}{2m} - E\right) \\ &= \frac{V^N}{h^{3N}} \int d^3p_i \delta\left(\sum_i \frac{p_i^2}{2m} - E\right) \end{aligned}$$

The surface of constant energy is just the surface of a sphere in $3N$ dimensional momentum space given by the coords $p_{1x}, p_{1y}, p_{1z}, \dots, p_{Nx}, p_{Ny}, p_{Nz}$. The radius of the sphere is $\sqrt{2mE}$.

Let $P \equiv \sqrt{\sum_i p_i^2}$ be the length of the momentum vector in the $3N$ dimensional momentum space.

$$\text{Then } \prod_{i=1}^N d^3 p_i = dP P^{3N-1} d\Omega_{3N}$$

↖ differential solid angle
in $3N$ dimensional space

$$g(E) = \frac{V^N}{h^{3N}} \int d\Omega_{3N} \int_0^{\infty} dP P^{3N-1} \delta\left(\frac{P^2}{2m} - E\right)$$

$$= \frac{V^N}{h^{3N}} S_{3N} \int_0^{\infty} dP P^{3N-1} \frac{\delta(P - \sqrt{2mE})}{(P/m)}$$

↖ area of unit
sphere in $3N$ -dim space

↖ from
converting the
 δ -function

$$= \frac{V^N}{h^{3N}} S_{3N} m (2mE)^{\frac{3N-2}{2}}$$

From Appendix C of Pathria (eqn C.76) or elsewhere,
one has the area of unit sphere in d -dim space

$$S_d = \frac{2\pi^{d/2}}{\Gamma(d/2)}$$

where $\Gamma(n) = (n-1)!$

for integer n

Γ is the Gamma function

$$\text{So } S_{3N} = \frac{2\pi^{3N/2}}{\left(\frac{3N}{2}-1\right)!}$$

$$g(E) = \frac{V^N}{h^{3N}} \frac{2\pi^{3N/2}}{\left(\frac{3N}{2}-1\right)!} m \frac{(2mE)^{\frac{3N}{2}}}{2mE}$$

$$g(E) = \frac{V^N (2\pi m E)^{3N/2}}{h^{3N} \left(\frac{3N}{2} - 1\right)!} \frac{1}{E}$$

$$\Omega(E) = \int_{E-\frac{\Delta}{2}}^{E+\frac{\Delta}{2}} dE' g(E') \approx g(E) \Delta$$

$$\Omega(E) = \frac{V^N (2\pi m E)^{3N/2}}{h^{3N} \left(\frac{3N}{2} - 1\right)!} \frac{\Delta}{E}$$

For large N , $\Omega(E)$ is a very rapidly increasing function of E ! $\sim E^{\frac{3N}{2}-1}$

We will now argue that $\Omega(E)$ is related to the entropy of the system.

Consider two subsystems separated by a wall

E_1	E_2
V_1	V_2
N_1	N_2

$$E_T = E_1 + E_2 \quad \text{energy conserved}$$

let $g_1(E_1)$ is density of states of system 1 with energy E_1
 $g_2(E_2)$ is density of states of system 2 with energy E_2

Now suppose the wall is thermally conducting so that energy can be transferred between the two systems. $\Rightarrow E_1$ can vary but $E_T = E_1 + E_2$ is fixed. What will be the value of E_1 when the system comes to equilibrium?