

pressure - separate out ground state from sum as we saw we needed to do in computing N/V

$$\begin{aligned} \frac{P}{k_B T} &= \frac{1}{V} \ln Z = -\frac{1}{V} \sum_{\vec{k}} \ln (1 - z e^{-\beta E(\vec{k})}) \\ &\approx -\frac{1}{V} \ln (1 - z) - \frac{4\pi}{(2\pi)^3} \int_0^{\infty} dk k^2 \ln (1 - z e^{-\beta \hbar^2 k^2 / 2m}) \\ &\quad \uparrow \quad \quad \quad \uparrow \\ &\quad \vec{k}=0 \text{ ground state} \quad \quad \text{all other } |\vec{k}| > 0 \text{ states} \\ &= \frac{1}{V} \ln \left(\frac{1}{1-z} \right) + \frac{g_{5/2}(z)}{\lambda^3} \quad \lambda = \left(\frac{h^2}{2\pi m k_B T} \right)^{1/2} \end{aligned}$$

where $g_{5/2}(z) \equiv \frac{1}{\Gamma(5/2)} \int_0^{\infty} dy \frac{y^{3/2}}{z^{-1} e^y - 1}$ as derived when we began our discussion of quantum gases

also recall the number of bosons occupying the ground state is

$$n(0) = \frac{1}{z^{-1} e^{\beta E(0)} - 1} = \frac{1}{z^{-1} - 1} = \frac{z}{1-z}$$

$$\text{So } n(0) + 1 = \frac{z}{1-z} + 1 = \frac{1}{1-z}$$

$$\frac{P}{k_B T} = \frac{\ln(n(0) + 1)}{V} + \frac{g_{5/2}(z)}{\lambda^3}$$

In the thermodynamic limit of $V \rightarrow \infty$, the first term always vanishes as $n(0) \leq N = nV$ and $\lim_{V \rightarrow \infty} \left[\frac{\ln(nV)}{V} \right] = 0$


So the condensate does not contribute to the pressure.

This is not surprising as particles in the condensate have $\vec{k}=0$ and hence carry no momentum. In the kinetic theory of gases, one sees that pressure arises from particles with finite momentum $|\vec{p}| > 0$ hitting the walls of the container

So $\frac{p}{k_B T} = \frac{g_{5/2}(z)}{\lambda^3} = g_{5/2}(z) \left(\frac{2\pi m k_B T}{h^2} \right)^{3/2}$

$p = g_{5/2}(z(T)) \left(\frac{2\pi m}{h^2} \right)^{3/2} (k_B T)^{5/2}$ ← equation of state

↑ for a system of fixed density n , z must be chosen to be a function of T that gives the desired density n .




Note $g_{5/2}(z=1) = \zeta(5/2) = 1.342$ is finite

In thermodynamic limit of $V \rightarrow \infty$, $z=1$ for $T \leq T_c(m)$

↑ critical temperature depends on the system's fixed density

$\Rightarrow p = g_{5/2}(1) \left(\frac{2\pi m}{h^2} \right)^{3/2} (k_B T)^{5/2}$ for $T \leq T_c$

Note: for $T \leq T_c$, the pressure $p \propto T^{5/2}$ is independent of the system density!



p vs T curves at constant density n


recall $T_c(m) \sim m^{2/3}$

$T_c(m) = \left(\frac{m}{2.16} \right)^{2/3} \frac{h^2}{2\pi m k_B}$

Define $m_c(T) = 2.612 \left(\frac{2\pi m k_B T}{h^2} \right)^{3/2}$ inverse of $T_c(m)$

$m_c(T)$ is the critical density at a given T
 — a system with $m > m_c(T)$ will be in a
 Bose condensed mixed state at temperature T .

phase diagram in p - T plane




Can also consider the transition in terms of
 p and $v = \frac{V}{N} = \frac{1}{m}$ for various fixed T .

At the transition $p \propto T_c(m)^{5/2}$, $T_c(m) \propto m^{2/3}$

\Rightarrow at the transition $p \propto (m^{2/3})^{5/2} = m^{5/3} = v^{-5/3}$
 below the transition p is independent of
 density and hence independent of v .

For fixed T , the transition occurs when density m
 exceeds $m_c(T)$, or when v drops below $v_c(T) = \frac{1}{m_c(T)}$
 $v_c(T) \sim T^{-3/2}$

curves of p vs v at constant T



Thermodynamic functions

Earlier we found $\frac{E}{V} = \frac{3}{2} p$

$$\Rightarrow \frac{E}{N} = \frac{3}{2} p \frac{V}{N} = \frac{3}{2} p v = \frac{3}{2} \frac{k_B T v}{\lambda^3} g_{5/2}(z)$$

In above we regard $\frac{E}{N}$ as a function of either v or z . That is we either determine v for a given z, T or we determine z needed for a given v, T (Recall $z = e^{\beta \mu}$, $v = \frac{V}{N}$ and N and μ are conjugate variables)

$z=1$ in mixed state
 $z < 1$ in normal state

specific heat

$$\frac{C_V}{N k_B} = \frac{1}{k_B} \left(\frac{\partial (E/N)}{\partial T} \right)_{v, N} = \frac{3}{2} v \left\{ \frac{d}{dT} \left(\frac{T}{\lambda^3} \right) g_{5/2}(z) + \frac{T}{\lambda^3} \frac{\partial g_{5/2}(z)}{\partial z} \left\{ \frac{dz}{dT} \right\} \right.$$

For $T \leq T_c$, $z = 1$ so $\frac{dz}{dT} = 0$ and only 1st term remains

$$\frac{T}{\lambda^3} \propto T^{5/2} \text{ so } \frac{d}{dT} \left(\frac{T}{\lambda^3} \right) = \frac{5}{2} \left(\frac{T}{\lambda^3} \right) \frac{1}{T} = \frac{5}{2} \frac{1}{\lambda^3}$$

\swarrow $z=1$ here for all $T \leq T_c$

$$\Rightarrow \frac{C_V}{Nk_B} = \frac{3}{2} \nu \left(\frac{5}{2} \frac{1}{\lambda^3} \right) g_{5/2}(1) = \frac{15}{4} g_{5/2}(1) \frac{\nu}{\lambda^3}$$

$$= \frac{15}{4} g_{5/2}(1) \nu \left(\frac{2\pi m k_B T}{h^2} \right)^{3/2}$$

Note, at T_c , $n = \frac{g_{3/2}(1)}{\lambda_c^3}$, and $\nu = \frac{1}{n}$

$$\frac{C_V(T_c)}{Nk_B} = \frac{15}{4} \frac{g_{5/2}(1)}{g_{3/2}(1)} = \frac{15}{4} \frac{1.341}{2.612} = 1.925 \leftarrow \text{this is larger than the classical ideal gas value of } \frac{3}{2}$$

So $\boxed{\frac{C_V}{Nk_B} = 1.925 \left(\frac{T}{T_c} \right)^{3/2} \quad T \leq T_c}$

For $T \geq T_c$, z varies with T and we need to evaluate the 2nd term as well

1st term gives $\frac{15}{4} g_{5/2}(z(T)) \frac{\nu}{\lambda^3}$ \swarrow here z depends on T for $T > T_c$

2nd term: from Pathria Appendix D Eq(10),

$$z \frac{d}{dz} [g_\nu(z)] = g_{\nu-1}(z)$$

$$\Rightarrow \frac{d g_{5/2}}{dz} \frac{dz}{dT} = g_{3/2} \frac{1}{z} \frac{dz}{dT}$$

To find $\frac{1}{z} \frac{dz}{dT}$ consider our earlier result for the density when $T > T_c$:

$$m = \frac{g_{3/2}(z)}{\lambda^3} \quad \leftarrow \text{determines } z(T) \text{ for fixed } m$$

$$\text{for } m \text{ fixed} \Rightarrow 0 = \frac{dm}{dT} = \frac{d}{dT} \left(\frac{1}{\lambda^3} \right) g_{3/2} + \frac{1}{\lambda^3} \frac{dg_{3/2}}{dz} \frac{dz}{dT}$$

$$0 = \frac{3}{2} \frac{1}{\lambda^3 T} g_{3/2} + \frac{1}{\lambda^3} g_{1/2} \frac{1}{z} \frac{dz}{dT}$$

$$\Rightarrow \frac{1}{z} \frac{dz}{dT} = -\frac{3}{2} \frac{g_{3/2}}{g_{1/2}} \frac{1}{T}$$

$$\frac{C_V}{Nk_B} = \frac{15}{4} g_{5/2}(z) \frac{v}{\lambda^3} + \frac{3}{2} v \frac{T}{\lambda^3} g_{3/2}(z) \left(-\frac{3}{2} \right) \frac{g_{3/2}(z)}{g_{1/2}(z)} \frac{1}{T}$$

$$\text{use } m = \frac{1}{v} = \frac{g_{3/2}(z)}{\lambda^3} \Rightarrow \frac{v}{\lambda^3} = \frac{1}{g_{3/2}(z)}$$

$$\boxed{\frac{C_V}{Nk_B} = \frac{15}{4} \frac{g_{5/2}(z)}{g_{3/2}(z)} - \frac{9}{4} \frac{g_{3/2}(z)}{g_{1/2}(z)} \quad T > T_c}$$

$$\text{Note } g_{1/2}(1) = \sum_{e=1}^{\infty} \frac{1}{e^{1/2}} \rightarrow \infty$$

So as $T \rightarrow T_c^+$ from above, and $z \rightarrow 1$

$$\frac{C_V}{Nk_B}(T_c^+) = \frac{15}{4} \frac{g_{5/2}(1)}{g_{3/2}(1)} - \frac{9}{4} \frac{g_{3/2}(1)}{\infty} = \frac{15}{4} \frac{1.341}{2.612} = 1.925$$

\Rightarrow C_V is continuous at T_c

Finally we want to show that although C_V is continuous at T_c , $\frac{dC_V}{dT}$ is discontinuous

For $T \leq T_c$ $\frac{C_V}{Nk_B} = 1.925 \left(\frac{T}{T_c} \right)^{3/2}$

$$\frac{d}{dT} \left(\frac{C_V}{Nk_B} \right) = \frac{3}{2} (1.925) \left(\frac{T}{T_c} \right)^{1/2} \frac{1}{T_c} = 2.89 \left(\frac{T}{T_c} \right)^{1/2} \frac{1}{T_c}$$

so slope at T_c^- (just below T_c)

\therefore $\frac{d}{dT} \left(\frac{C_V}{Nk_B} \right) = \frac{2.89}{T_c}$, $T = T_c^-$

For $T > T_c$

$$\frac{C_V}{Nk_B} = \frac{15}{4} \frac{g_{5/2}(z)}{g_{3/2}(z)} - \frac{9}{4} \frac{g_{3/2}(z)}{g_{1/2}(z)}$$

$$\frac{d}{dT} \left(\frac{C_V}{Nk_B} \right) = \frac{15}{4} \frac{g_{3/2} \frac{dg_{5/2}}{dz} \frac{dz}{dT} - g_{5/2} \frac{dg_{3/2}}{dz} \frac{dz}{dT}}{(g_{3/2}(z))^2}$$

$$- \frac{9}{4} \frac{g_{1/2} \frac{dg_{3/2}}{dz} \frac{dz}{dT} - g_{3/2} \frac{dg_{1/2}}{dz} \frac{dz}{dT}}{(g_{1/2}(z))^2}$$

$$= \frac{1}{z} \frac{dz}{dT} \left\{ \frac{15}{4} \left(\frac{g_{3/2}^2 - g_{5/2} g_{1/2}}{g_{3/2}^2} \right) - \frac{9}{4} \left(\frac{g_{1/2}^2 - g_{3/2} g_{-1/2}}{g_{1/2}^2} \right) \right\}$$

use $\frac{1}{z} \frac{dz}{dT} = -\frac{3}{2} \frac{g_{3/2}}{g_{1/2}} \frac{1}{T}$ as found earlier

$$\frac{d}{dT} \left(\frac{C_V}{Nk_B} \right) = -\frac{3}{8T} \frac{g_{3/2}}{g_{1/2}} \left\{ 15 \left(1 - \frac{g_{5/2} g_{1/2}}{g_{3/2}^2} \right) - 9 \left(1 - \frac{g_{3/2} g_{-1/2}}{g_{1/2}^2} \right) \right\}$$

Now as $T \rightarrow T_c^+$ from above, $z \rightarrow 1$, we have
 $g_{5/2}(1)$ and $g_{3/2}(1)$ are finite, but $g_{1/2}(1)$ and
 $g_{-1/2}(1) \rightarrow \infty$

\Rightarrow at T_c^+

$$\frac{d}{dT} \left(\frac{C_V}{Nk_B} \right) = \frac{45}{8T_c} \frac{g_{5/2}(1)}{g_{3/2}(1)} - \frac{27}{8T_c} \frac{g_{3/2}^2(1) g_{-1/2}(1)}{g_{1/2}^3(1)}$$

Now from Pathria Appendix D Eq (8)

$$g_\nu(1) = \lim_{a \rightarrow 0} \frac{\Gamma(1-\nu)}{a^{1-\nu}}$$

$$\text{So } \frac{g_{-1/2}(1)}{g_{3/2}^3(1)} = \lim_{a \rightarrow 0} \frac{\Gamma(3/2)}{a^{3/2}} \left(\frac{a^{1/2}}{\Gamma(1/2)} \right)^3 = \frac{\Gamma(3/2)}{[\Gamma(1/2)]^3}$$

$$= \frac{\frac{1}{2} \pi^{1/2}}{\pi^{3/2}} = \frac{1}{2\pi}$$

$$\text{since } \Gamma(1/2) = \sqrt{\pi} \\ \Gamma(3/2) = \frac{1}{2} \sqrt{\pi}$$


$$\frac{d}{dT} \left(\frac{C_V}{Nk_B} \right) = \frac{45}{8} \frac{1.341}{2.612} \frac{1}{T_c} - \frac{27}{8} \frac{(2.612)^2}{2\pi} \frac{1}{T_c}$$

$$= \frac{2.89}{T_c} - \frac{3.66}{T_c} = -\frac{0.77}{T_c}$$

$$\boxed{\frac{d}{dT} \left(\frac{C_V}{Nk_B} \right) = -\frac{0.77}{T_c}, \quad T = T_c^+}$$

slope of C_V
is discontinuous at
 T_c .

C_V has a cusp at T_c



goes to classical $\frac{3}{2}$ as $T \rightarrow \infty$

$$\frac{dC_V}{dT} > 0 \quad \text{for } T = T_c^-$$

$$\frac{dC_V}{dT} < 0 \quad \text{for } T = T_c^+$$

Entropy

For single species gas we had for Gibbs free energy

$$G = N\mu$$

$$\text{Also } G = E - TS + pV$$

(since G is Legendre transform of E with respect to S and V)

$$\Rightarrow N\mu = E - TS + pV$$

$$\text{or } S = \frac{E + pV - N\mu}{T}$$

$$\frac{S}{Nk_B} = \frac{E + pV}{Nk_B T} - \frac{\mu}{k_B T}$$

we had earlier $E = \frac{3}{2} pV \Rightarrow pV = \frac{2}{3} E$

$$\frac{S}{Nk_B} = \frac{5}{3} \frac{E}{N} \frac{1}{k_B T} - \frac{\mu}{k_B T}$$

$$z = e^{M/k_B T}, \quad z = 1 \text{ for } T < T_c$$

We had earlier $\frac{E}{N} = \frac{3}{2} \frac{k_B T}{\lambda^3} g_{5/2}(z)$

and $n = \frac{1}{v} = \frac{g_{3/2}(z)}{\lambda^3}$ for $T > T_c$

$$\Rightarrow \frac{S}{N k_B} = \frac{5}{2} \frac{v}{\lambda^3} g_{5/2}(z) - \ln z = \begin{cases} \frac{5}{2} \frac{g_{5/2}(z)}{g_{3/2}(z)} - \ln z, & T > T_c \\ \frac{5}{2} \frac{v}{\lambda^3} g_{5/2}(1), & T \leq T_c \end{cases}$$

Note: For $T \leq T_c$ we had that the density of the ~~normal~~ a density $m_0 = n - \frac{g_{3/2}(1)}{\lambda^3}$ in

the condensate, and a density $\frac{g_{3/2}(1)}{\lambda^3}$ in the 'normal' state (i.e. the density of excited particles) $\frac{g_{3/2}(1)}{\lambda^3} \equiv m_n$

$$\Rightarrow \text{for } T \leq T_c, \quad \frac{S}{N k_B} = \frac{5}{2} \left(\frac{m_n}{m} \right) \frac{g_{5/2}(1)}{g_{3/2}(1)} \rightarrow 0 \text{ as } T \rightarrow 0$$

We can imagine that each normal particle carries

entropy $\frac{5}{2} k_B \frac{g_{5/2}(1)}{g_{3/2}(1)}$

The entropy at $T < T_c$ /per particle

is just the ~~frac~~ above entropy per 'normal' particle times the fraction of normal particles.

\Rightarrow normal particles carry the entropy
condensate has zero entropy

entropy difference per particle between normal state and condensed state is $\frac{\Delta S}{N} = \frac{5}{2} k_B \frac{g_{5/2}(1)}{g_{3/2}(1)}$

latent heat of condensation

$$L = T \Delta S = \frac{5}{2} k_B T \frac{g_{5/2}(1)}{g_{3/2}(1)}$$

energy released upon converting one normal particle to one condensate particle.

⇒ mixed phase is like coexistence region of a 1st order phase transition (like water ↔ ice ~~?~~ - need to remove energy to turn water to ice)

⇒ "two fluid" model of mixed region

Bose-Einstein Condensation in laser cooled gases

Gases of alkali atoms Li, Na, K, Rb, Cs

- all have a single s-electron in outermost shell.
- important for efficiency of laser cooling
- use isotopes such that total intrinsic spin of all electrons and nucleons add up to an integer T_0
 \Rightarrow atoms are bosons
- all have a net magnetic moment - used to confine dilute gas of atoms in a "magnetic trap"
- use "laser cooling" to get very low temperatures in low density gases, to try and see BEC

magnetic trap \rightarrow effective harmonic potential for atoms

$$V(r) = \frac{1}{2} m \omega_0^2 r^2 \quad \omega_0 \approx 2\pi \times 100 \text{ Hz}$$

1995 - 10^3 atoms with $T_0 \sim 100 \text{ nK}$

1999 - 10^8 atoms with $T_0 \sim \mu\text{K}$ gas size \sim many microns

How was BEC in these systems observed?

energy levels of ideal (non-interacting) bosons in harmonic trap

$$E(n_x, n_y, n_z) = (n_x + n_y + n_z + 3/2) \hbar \omega_0$$

n_x, n_y, n_z integers

ground state condensate wavefunction

$$\psi_0(r) \sim e^{-r^2/2a^2} \quad \text{with} \quad a = \left(\frac{\hbar}{m\omega_0}\right)^{1/2}$$

$a \sim 1 \mu\text{m}$ for current traps

\Rightarrow condensate has spatial extent $\sim a$

The spatial extent of the n^{th} excited energy level is roughly

$$m\omega_0^2 \langle r^2 \rangle \sim E(n) \approx n\hbar\omega_0$$

$$\Rightarrow \langle r^2 \rangle \sim \frac{n\hbar}{m\omega_0} \quad \text{or} \quad \sqrt{\langle r^2 \rangle} = \left(\frac{n\hbar}{m\omega_0} \right)^{1/2}$$

For $k_B T \gg \hbar\omega_0$, the atoms are excited up to level $n \sim \frac{k_B T}{\hbar\omega_0}$

\Rightarrow spatial extent of the normal component of the gas is

$$R \sim \left(\frac{n\hbar}{m\omega_0} \right)^{1/2} \sim \left(\frac{\hbar k_B T}{\hbar m \omega_0^2} \right)^{1/2} = \left(\frac{k_B T}{m\omega_0^2} \right)^{1/2}$$

$$R \sim a \left(\frac{k_B T}{\hbar\omega_0} \right)^{1/2} \gg a$$

If T_c is the BEC transition temperature, then for $T \gg T_c$ one sees a more or less uniform cloud of atoms with radius $R \sim a \left(\frac{k_B T}{\hbar\omega_0} \right)^{1/2} \gg a$.
But when one cools to $T < T_c$, one now has a finite fraction of the atoms condensed in the ground state, \Rightarrow superimposed on the atomic cloud of radius R , one sees the growth of a sharp peak in density at the center of cloud - this peak has a radius $a \ll R$.

To find T_c , (use $z=1$ at $T \leq T_c$)
for $T \leq T_c$

$$n = n_0 + \int_0^\infty dn_x \int_0^\infty dn_y \int_0^\infty dn_z \frac{1}{e^{(n_x + n_y + n_z) \hbar \omega_0 / k_B T} - 1}$$

$$= n_0 + \left(\frac{k_B T}{\hbar \omega_0} \right)^3 \int_0^\infty dx \int_0^\infty dy \int_0^\infty dz \frac{1}{e^{(x+y+z)} - 1}$$

$$= n_0 + \left(\frac{k_B T}{\hbar \omega_0} \right)^3 \zeta(3)$$

at T_c , $n_0 = 0 \Rightarrow k_B T_c = \hbar \omega_0 \left(\frac{n}{\zeta(3)} \right)^{1/3}$

condensate density $n_0(T) = n \left(1 - \left(\frac{T}{T_c} \right)^3 \right)$

↑
different from ideal free gas due to presence of magnetic trapping potential

$$\zeta(3) = 1 + \frac{1}{2^3} + \frac{1}{3^3} + \frac{1}{4^3} + \dots$$