

Lindhard Dielectric Function - fixes Thomas-Fermi at large \bar{k} .

Consider a potential $U(\vec{r})$ applied to the electron gas. (for an electrostatic potential $U = -eV^{tot}$)

To compute the change in electron density $\delta n(\vec{r})$ we could compute the effect of U on electron eigenstates, and then use these new eigenstates to compute δn , summing over all occupied eigenstates

Using ~~stationary~~ Rayleigh-Schrodinger stationary perturbation theory, to lowest order in U the eigenstates become

$$|\psi_k\rangle = |k\rangle + \sum_{k'} \frac{|k'\rangle \langle k'|U|k\rangle}{E_k - E_{k'}}$$

where $|k\rangle$ is the unperturbed plane wave eigenstate with energy $E_k = \hbar^2 k^2 / 2m$, and $|\psi_k\rangle$ is the new eigenstate resulting from the perturbation U .

The electron density as a function of position for the state $|\psi_k\rangle$ is

$$|\langle r | \psi_k \rangle|^2$$

so the change in electron density due to the perturbation U is

$$|\langle r | \psi_k \rangle|^2 - |\langle r | k \rangle|^2$$

$$= \langle \psi_k | r \rangle \langle r | \psi_k \rangle - \langle k | r \rangle \langle r | k \rangle$$

$$= \left[\langle r|k \rangle + \sum_{k'} \frac{\langle r|k' \rangle \langle k'|U|k \rangle}{\epsilon_k - \epsilon_{k'}} \right] \left[\langle k|r \rangle + \sum_{k'} \frac{\langle k'|r \rangle \langle r|U|k' \rangle}{\epsilon_k - \epsilon_{k'}} \right] - \langle k|r \rangle \langle r|k \rangle$$

To linear order in U this gives

$$= \sum_{k'} \left\{ \frac{\langle r|k \rangle \langle k'|r \rangle \langle k'|U|k \rangle}{\epsilon_k - \epsilon_{k'}} + \frac{\langle k|r \rangle \langle r|k' \rangle \langle k'|U|k \rangle}{\epsilon_k - \epsilon_{k'}} \right\}$$

Now $\langle r|k \rangle = \frac{e^{i\vec{k}\cdot\vec{r}}}{\sqrt{V}}$ $V = \text{volume}$

$$\langle k|r \rangle = \frac{e^{-i\vec{k}\cdot\vec{r}}}{\sqrt{V}}$$

$$\langle k'|U|k \rangle = \int \frac{d^3r}{V} e^{-i\vec{k}'\cdot\vec{r}} U(\vec{r}) e^{i\vec{k}\cdot\vec{r}}$$

$$= \frac{1}{V} \int d^3r e^{-i(\vec{k}'-\vec{k})\cdot\vec{r}} U(\vec{r})$$

$$= \frac{1}{V} U_{\vec{k}'-\vec{k}} \quad \text{Fourier transf of } U(\vec{r})$$

So above is

$$= \frac{1}{V^2} \sum_{k'} \left\{ \frac{e^{-i(\vec{k}'-\vec{k})\cdot\vec{r}} U_{\vec{k}-\vec{k}'}}{\epsilon_k - \epsilon_{k'}} + \frac{e^{-i(\vec{k}-\vec{k}')\cdot\vec{r}} U_{\vec{k}'-\vec{k}}}{\epsilon_k - \epsilon_{k'}} \right\}$$

The total induced electron density δn is obtained by summing over all occupied states
 spin degeneracy

$$\delta n(\vec{r}) = 2 \sum_{\vec{k}} f_{\vec{k}} \frac{1}{V} \sum_{\vec{k}'} \left\{ e^{-i(\vec{k}' - \vec{k}) \cdot \vec{r}} \frac{U_{\vec{k} - \vec{k}'}}{\epsilon_{\vec{k}} - \epsilon_{\vec{k}'}} + e^{-i(\vec{k} - \vec{k}') \cdot \vec{r}} \frac{U_{\vec{k}' - \vec{k}}}{\epsilon_{\vec{k}} - \epsilon_{\vec{k}'}} \right\}$$

where $f_{\vec{k}}$ is the Fermi occupation function $\frac{1}{e^{(\epsilon_{\vec{k}} - \mu)/k_B T} + 1}$

Fourier transform to get $\delta n(\vec{q})$

$$\begin{aligned} \delta n(\vec{q}) &= \int d^3r e^{-i\vec{q} \cdot \vec{r}} \delta n(\vec{r}) \\ &= \frac{1}{V} \sum_{\vec{k}, \vec{k}'} f_{\vec{k}} \left\{ \frac{V \delta_{\vec{q}, \vec{k} - \vec{k}'}}{\epsilon_{\vec{k}} - \epsilon_{\vec{k}'}} U_{\vec{k} - \vec{k}'} + \frac{V \delta_{\vec{q}, \vec{k}' - \vec{k}}}{\epsilon_{\vec{k}} - \epsilon_{\vec{k}'}} U_{\vec{k}' - \vec{k}} \right\} \end{aligned}$$

where the integrals over the plane wave factors give the $V \delta_{\vec{q}, \vec{k} - \vec{k}'}$ terms. Now use the δ 's to do the sum on \vec{k}' .

$$\delta n(\vec{q}) = \frac{2}{V} \sum_{\vec{k}} f_{\vec{k}} \left\{ \frac{U_{\vec{q}}}{\epsilon_{\vec{k}} - \epsilon_{\vec{k} - \vec{q}}} + \frac{U_{\vec{q}}}{\epsilon_{\vec{k}} - \epsilon_{\vec{k} + \vec{q}}} \right\}$$

So

$$\frac{\delta n(\vec{q})}{U_{\vec{q}}} = \frac{2}{V} \sum_{\vec{k}} f_{\vec{k}} \left\{ \frac{1}{\epsilon_{\vec{k}} - \epsilon_{\vec{k} - \vec{q}}} + \frac{1}{\epsilon_{\vec{k}} - \epsilon_{\vec{k} + \vec{q}}} \right\}$$

Now make substitution $\vec{k}' = \vec{k} - \vec{q}$ in first summation term to get

$$\frac{\delta m(q)}{U_q} = \frac{2}{V} \sum_{\vec{k}} \frac{f_{\vec{k}+\vec{q}} - f_{\vec{k}}}{\epsilon_{\vec{k}+\vec{q}} - \epsilon_{\vec{k}}}$$

$$\frac{\delta m(q)}{U_q} = \int \frac{d^3k}{4\pi^3} \frac{f_{\vec{k}+\vec{q}} - f_{\vec{k}}}{\epsilon_{\vec{k}+\vec{q}} - \epsilon_{\vec{k}}}$$

For electrostatic potential, $U_q = -eV_{\vec{q}}^{\text{tot}}$,
and $\delta p = -e\delta m$, so

~~$$\frac{\delta p}{V_{\text{tot}}}$$~~

$$\frac{\delta p(q)}{V_{\text{tot}}(q)} = \frac{-e\delta m_q}{U_q/(-e)} = e^2 \frac{\delta m_q}{U_q}$$

$$= e^2 \int \frac{d^3k}{4\pi^3} \frac{f_{\vec{k}+\vec{q}} - f_{\vec{k}}}{\epsilon_{\vec{k}+\vec{q}} - \epsilon_{\vec{k}}}$$

For small q , $f_{\vec{k}+\vec{q}} - f_{\vec{k}} \approx \frac{\partial f}{\partial \epsilon} \frac{\partial \epsilon}{\partial \vec{q}} \cdot \vec{q}$

$$\epsilon_{\vec{k}+\vec{q}} - \epsilon_{\vec{k}} \approx \frac{\partial \epsilon}{\partial \vec{q}} \cdot \vec{q}$$

$$\frac{\delta p}{V_{\text{tot}}} = e^2 \int \frac{d^3k}{4\pi^3} \frac{\partial f}{\partial \epsilon} = e^2 \int d\epsilon g(\epsilon) \frac{\partial f}{\partial \epsilon}$$

as $T \rightarrow 0$ $\frac{\partial f}{\partial \epsilon} \rightarrow -\delta(\epsilon - \epsilon_F)$

$$\frac{\delta p}{V_{\text{tot}}} = -e^2 g(\epsilon_F), \text{ so } \epsilon(\vec{q}) = 1 - \frac{4\pi}{q^2} \frac{\delta p}{V_{\text{tot}}} = 1 + \frac{4\pi e^2}{q^2} g(\epsilon_F)$$

Same as Thomas-Fermi result

Friedel oscillations + Kohn anomaly

Lindhard dielectric function at bigger q

$$\epsilon(q) = 1 + \frac{4\pi e^2}{q^2} \sum_k \frac{f_k - f_{k+q}}{\epsilon_{k+q} - \epsilon_k}$$

$$\epsilon_{k+q} - \epsilon_k = \left(\frac{q^2 + 2\vec{k} \cdot \vec{q}}{2m} \right) \hbar^2$$


$$\epsilon_k = \frac{\hbar^2 k^2}{2m}$$

$$\epsilon(q) = 1 + \frac{4\pi e^2}{q^2} \frac{2m}{\hbar^2} \int_R \frac{d^3k}{(2\pi)^3} \frac{1}{q^2 + 2\vec{k} \cdot \vec{q}}$$


spin up
or down
x 2 x 2
[k+q full
k empty]

Region of k-space
such that [k full
k+q empty]

The region R is such that $|\vec{k}| < k_F$ and $|\vec{k} + \vec{q}| > k_F$
We can depict it graphically as



as q increases, region R also increases until $q \geq 2k_F$



The integral $\epsilon(q) = 1 + \frac{4\pi e^2}{q^2} \frac{g m}{\hbar^2} \int_R \frac{d^3 k}{(2\pi)^3} \frac{1}{q^2 + 2\vec{k} \cdot \vec{q}}$
 can be done explicitly and one gets

$$\epsilon(q) = 1 + \frac{4\pi e^2}{q^2} g(\epsilon_F) \left[\frac{1}{2} + \frac{1-x^2}{4x} \ln \left| \frac{1+x}{1-x} \right| \right]$$


where $x = q/2k_F$

as $x \rightarrow 0$, $[\dots] = 1$ and we get back Thomas-Fermi result
 at $x = 1$, i.e. $q = 2k_F$, $\epsilon(q)$ has a logarithmic singularity

If we formally write $\epsilon(q) = 1 + \frac{k_0^2(q)}{q^2}$

to define a q dependent screening length $k_0(q)$
 then

$$\frac{k_0^2(q)}{k_0^2(0)} = [\dots]$$



as q increases the effective screening length $1/k_0$ increases.
 Screening is less effective at small length scales than in Thomas Fermi approx

If you take the Fourier transf of $\frac{4\pi Q}{q^2 \epsilon(q)}$
 to get real space potential of a point charge,


the singularity at $q = 2k_F$ gives rise to a piece

$$\sim \frac{1}{r^3} \cos(2k_F r)$$

decays more slowly than T-F and oscillates in sign, electron is alternatively attracted and repelled by the charge Q with a period $\frac{\pi}{k_F}$


These are known as "Friedel" or "Ruderman-Kittel" oscillations. These are important for giving the "RKKY" interaction between magnetic impurities, that is the origin of "spin glasses". - see homework problem for details.

The origin of the singularity at $q = 2k_F$ is understood more physically in terms of the behavior of the region of integration R.



As q increases, the region R increases, until $q = 2k_F$. When $q > 2k_F$, R is the entire Fermi sphere and no longer changes as q increases further. This singularity in the volume R gives rise to the singularity in $\epsilon(q)$ at $q = 2k_F$

This is true also for more generally shaped Fermi surfaces - $\epsilon(\vec{q})$ will be singular for any \vec{q} that displaces the Fermi surfaces so they touch at a tangential point.



Kohn effect: a phonon (ion lattice vibration) at ~~such~~ a wavevector \vec{q} sets up an electrostatic potential with wavevector \vec{q} .

If \vec{q} is just such a critical \vec{q} as above, where $\epsilon(\vec{q})$ has a singularity, the screened ion-ion interaction will be proportional to $1/\epsilon(\vec{q})$, and also have a singularity. Since the phonon frequency $\omega(\vec{q})$ is determined by the ion-ion interaction, we expect to see $\omega(\vec{q})$ have a weak singularity at the above critical \vec{q} 's.

RKKY Interaction and Spin Glasses

In our discussion of the Lindhard dielectric function we saw that:

If there is a potential energy $U(\vec{r})$ that couples to the electron density, i.e. the perturbation in the Hamiltonian is

$$\sum_i U(\vec{r}_i) = \int d^3r U(\vec{r}) n(\vec{r})$$

with $n(\vec{r}) \equiv \sum_i \delta(\vec{r} - \vec{r}_i)$ is the electron density then U induces a change in electron density $\delta n(\vec{r})$ given, in Fourier transform space, by

$$\delta n(\vec{q}) = \chi(\vec{q}) U(\vec{q})$$

$$\text{with } \chi(\vec{q}) \equiv \frac{2}{V} \sum_{\vec{k}} \frac{f_{\vec{k}+\vec{q}} - f_{\vec{k}}}{\epsilon_{\vec{k}+\vec{q}} - \epsilon_{\vec{k}}} = 2 \int \frac{d^3k}{(2\pi)^3} \frac{f_{\vec{k}+\vec{q}} - f_{\vec{k}}}{\epsilon_{\vec{k}+\vec{q}} - \epsilon_{\vec{k}}}$$

where $f_{\vec{k}}$ is the Fermi occupation function for the free electron state with wave vector \vec{k} and energy $\epsilon_{\vec{k}}$.

Consider a magnetic impurity with spin \vec{S}_0 located at position \vec{R}_0 . We will assume the interaction of \vec{S}_0 with the conduction electrons is via a local spin-spin interaction.

$$\delta H = -J\mu_B \vec{S}_0 \cdot \sum_i \vec{s}_i |\psi_i(\vec{R}_0)|^2$$

\uparrow spin of electron i \uparrow probability for electron i to be at position \vec{R}_0

$$= J \vec{S}_0 \cdot \vec{m}(\vec{R}_0)$$

\uparrow magnetization density of electrons

Let us take the direction of \vec{S}_0 to be \hat{z} . Then

$$\delta H = -J\mu_B S_0 [m_{\uparrow}(\vec{R}_0) - m_{\downarrow}(\vec{R}_0)]$$

\uparrow density of \uparrow electrons \uparrow density of \downarrow electrons

$$= \delta H_{\uparrow} + \delta H_{\downarrow}$$

$$\delta H_{\uparrow} \equiv -J\mu_B S_0 m_{\uparrow}(\vec{R}_0) = \int d^3r U_{\uparrow}(\vec{r}) m_{\uparrow}(\vec{r})$$

$$\delta H_{\downarrow} = +J\mu_B S_0 m_{\downarrow}(\vec{R}_0) = \int d^3r U_{\downarrow}(\vec{r}) m_{\downarrow}(\vec{r})$$

$$\text{where } \begin{cases} U_{\uparrow}(\vec{r}) = -J\mu_B S_0 \delta(\vec{r} - \vec{R}_0) \\ U_{\downarrow}(\vec{r}) = +J\mu_B S_0 \delta(\vec{r} - \vec{R}_0) \end{cases}$$

We then have that U_{\uparrow} and U_{\downarrow} induce perturbations δm_{\uparrow} and δm_{\downarrow} in the spin \uparrow and spin \downarrow electron densities.

$$\delta m_{\uparrow}(\vec{q}) = \frac{1}{2} \chi(\vec{q}) U_{\uparrow}(\vec{q})$$

$$\delta m_{\downarrow}(\vec{q}) = \frac{1}{2} \chi(\vec{q}) U_{\downarrow}(\vec{q}) = -\frac{1}{2} \chi(\vec{q}) U_{\uparrow}(\vec{q})$$

\uparrow
factor of $\frac{1}{2}$ since m_{\uparrow} and m_{\downarrow} are both $\frac{1}{2}$ of total density $m = m_{\uparrow} + m_{\downarrow}$

induced electron magnetization is then

$$\begin{aligned} m_z(\vec{q}) &= -\mu_B [\delta m_{\uparrow}(\vec{r}) - \delta m_{\downarrow}(\vec{r})] \\ &= -\mu_B \chi(\vec{q}) U_{\uparrow}(\vec{q}) \end{aligned}$$

$$\text{Now } U_{\uparrow}(\vec{r}) = -J\mu_B S_0 \delta(\vec{r} - \vec{R}_0)$$

$$\begin{aligned} \text{so } U_{\uparrow}(\vec{q}) &= \int d^3\vec{r} e^{-i\vec{q}\cdot\vec{r}} U_{\uparrow}(\vec{r}) \\ &= -J\mu_B S_0 e^{-i\vec{q}\cdot\vec{R}_0} \end{aligned}$$

$$\begin{aligned} \Rightarrow m_z(\vec{r}) &= + \int \frac{d^3q}{(2\pi)^3} J\mu_B^2 S_0 \chi_q e^{-i\vec{q}\cdot\vec{R}_0} e^{i\vec{q}\cdot\vec{r}} \\ &= +J\mu_B^2 S_0 \int \frac{d^3q}{(2\pi)^3} \chi(\vec{q}) e^{i\vec{q}\cdot(\vec{r} - \vec{R}_0)} \end{aligned}$$

$$m_z(\vec{r}) = +J\mu_B^2 S_0 \chi(\vec{r} - \vec{R}_0)$$

\uparrow
Fourier transform of $\chi(\vec{q})$ evaluated at position $\vec{r} - \vec{R}_0$