

Monkeys to hominids

Evans 2002

2.5 5 7.5

Pe


World N

e in DNA

+18

3

2



- Distinct process from steady rate of increased difference in junk DNA.
- Most evident in parts of genes called human accelerated regions (HARs), of which 55 have been noted.


13 June 2011


Astronomy 106, Summer 2011


Monkeys to hominids (continued)

- HARs were discovered in 2006 by Katie Pollard (UCSF), as one of the first huge achievements of the new science of genomics.
- □ HAR1, chromosome 20, for example:
 - Present in reptiles onward.
 - Base-pair difference between chimpanzees and chickens: 2.
 - Base-pair difference between chimpanzees and humans: 18.


13 June 2011 Astronomy 106, Summer 2011



































- □ Raw meat is very difficult, Wikimedia time consuming and Commons energetically expensive to eat and on top of that not all of it can be digested.
- □ *H. erectus*'s smaller teeth and gut point to the development of using fire to **cook** meat
- Gathering around a fire for protection, warmth and cooking was a natural place for the growth of language, social skills and relationships

13 June 2011

Astronomy 106, Summer 2011

Mid-lecture Break

□ Homework #4 due Thursday. Recitation tomorrow.



13 June 2011

12

11

Genetic diversity of Africa

From *H. erectus* and onward, the adaptability and organizational skills permitted by larger brain size – especially tool- and speech-making capabilities – allowed hominids to spread past the bounds of their East African homeland.

□ Indeed, they had to, for dietary reasons.

- Grasslands spread and where prey go predators followBig herbivores like gorillas and chimps can live (in
- forests) at population densities of about 1 per km².
- But hominids, who preferred meat, could no longer digest leaves and grasses, and weren't so efficient at hunting yet, could only live at densities no greater than about 0.1 per km².

13 June 2011	Astronomy 106, Summer 2011	13

Genetic diversity of Africa (continued)


□ Thus competition among the hominids ...

- more advanced ones forced less advanced ones to inhospitable places,
- or killed them off and ate them.
- □ and their overspreading of temperate Africa south of the Sahara and the Sudd, a huge swamp along the Nile.
- □ By roughly 100,000 years ago, *H. sapiens* had occupied all of this part of Africa and eliminated its competition, leaving the apes and monkeys to their trees.
- □ Still dominating the hominid gene pool and having 0.1-1 Myr of differentiation, this process led to a notable diversity of races among *H. sapiens* that remains the greatest, even now that the species is worldwide.

13 June 2011 Astronomy 106, Summer 2011

14

Genetic diversity of Africa (continued)




Out of Africa

Of course, there was some leakage from the African gene pool, as hominids explored for better food sources.

- Presumably the best exit has always been to endure the passage of the Sudd and follow the Nile; this is consistent with the ages of archaic human fossils, which tend to be older the closer they are to the isthmus of Suez.
- □ There have been at least three escapes: by *H. erectus.*, by *H. heidelbergensis/neanderthalensis*, and by *H sapiens*.
- A small band of *H. erectus* escaped Africa somewhere around 1 Myr ago, rather early in the life of the species. This band multiplied rapidly and spread throughout the temperate regions of Africa, Europe and Asia.
 Thus "Java man" and "Peking man."

13 June 2011	Astronomy 106, Summer 2011	

16



Out of Africa (continued)

About 130,000 years ago, near the beginning of the most recent ice age (Wisconsin-Weichsel-Wurm), a small band of *H. neanderthalensis* left Africa and flourished by hunting ice-age megafauna, such as mastodons and mammoths, on the treeless tundra south of the ice sheets.

- □ They seem quickly to have displaced *H. erectus* in Europe and the Near East.
- Their remains, and the products of the Mousterian industry, comprise oldest finds of the Old Stone Age, or lower Paleolithic Age, in Europe.
 - This distinction means less to the rest of the world than it does to Europe, of course. Things in back home in Africa were developing more smoothly.

13 June 2011 Astronomy 106, Summer 2011 18

Ranges of H. neanderthalensis and (W) ice sheets 13 June 2011 19 Astronomy 106, Summer 2011



Out of Africa (continued)

Then, about 80,000 years ago, with the W ice age still in progress, a small group of *H. sapiens* left Africa via the Nile valley, and similarly to its predecessors began to reproduce, flourish, and expand all over the world.

- □ Also like its predecessors, this group represented a tiny selection of the gene pool, and reproduced very rapidly as it soon took over all the habitats.
- □ Since its numbers grew large so much faster than the time over which H. sapiens had evolved, the Out Of Africa portion of the species is genetically very homogeneous. Norwegians, Chinese and Australian Aborigines are very close genetically, and all are distinctively different from all of the remaining African races.

13 June 2011

Astronomy 106, Summer 2011

20

21

The Middle Paleolithic

H. sapiens brought with them many new improvements, for which the remains grace the Eurasian Middle Paleolithic:

even better tools (the Aurignacian industry), including composite structures made of wood or bone and pebbles sharpened to an edge. Neanderthals tried to follow suit here (the Chatelperronian industry), but it was too little too late.

new ideas, like

- boats. With these, H. sapiens colonized Australia about 40,000 years ago.
- (subprimate) animal friends. Dogs begin to appear around the campfires in the middle paleolithic. Astronomy 106, Summer 2011

13 June 2011

The beginnings of the Upper Paleolithic

By about the end of the W ice age 18,000 years ago, *H. sapiens* had done away with *H. neanderthalensis* and *H. erectus*.

- □ Despite many claims that these species interbred, especially in East Asia, this appears to be a dubious proposition. There is essentially no genetic legacy from *H*. *neanderthalensis* and *H*. *erectus* that *H*. *sapiens* didn't have when they left Africa.
- □ *H. sapiens* began to leave the first remains of cultural development, in the form of the famous cave paintings of such locations as Lascaux (France, at right) and Altamira (Spain).



22

13 June 2011 Astronomy 106, Summer 2011

What was *H. sapiens* brain-size advantage over *H. erectus*?

A. No advantage: brains roughly the same size.

B. H. sapiens brains are 50% larger than H. erectus.

C. H. sapiens brains are about twice the size of H. erectus.

D. *H. sapiens* brains are five times larger than *H. erectus*.

14 June 2011

Astronomy 106, Summer 2011

23

24

What was *H. sapiens* brain-size advantage over *H. neanderthalensis*?

A. No advantage: brains roughly the same size.

B. H. sapiens brains are 50% larger than Neanderthals.

- C. H. sapiens brains are about twice the size of Neanderthals.
- D. H. sapiens brains are five times larger than Neanderthals.

14 June 2011 Astronomy 106, Summer 2011

The beginnings of the Upper Paleolithic

- Within the next few thousand years they occupied and adapted to all of the habitable land of Europe, Asia and Australia, and followed the edges of the ice sheets as they retreated north.
- The adaptation in various climes, and the lack of communication over large distances, led during this time to differentiation of habits of "culture" as well as habits of dining.
 - The most notable of these is the origin of the basic families of language, since those left the most indelible mark.

13 June 2011	Astronomy 106, Summer 2011	25

