

1. (a) Show that, for a *blackbody* which occupies a small solid angle $\Delta\Omega$, at low temperatures the relation between temperature and $B - V$ color is something like

$$T \approx \frac{7400 \text{ K}}{(B - V) + 1.4}$$

(Hint: Use the Wien approximation and assume the properties of the B and V filters given in Lecture 3.)

- (b) This relation is a very poor approximation at high temperatures. Why?
2. Suppose that two stars with masses m_1 and m_2 are in circular orbits about their common center of mass with radii r_1 and r_2 , respectively. The orbital period is P .
- (a) Show that $r_1 = m_2 r / (m_1 + m_2)$ and $r_2 = m_1 r / (m_1 + m_2)$, where $r = r_1 + r_2$ is the distance between the two stars.
- (b) The speeds of the stars in orbit are constant and given by $v_1 = \Omega r_1$ and $v_2 = \Omega r_2$, respectively, where $\Omega = 2\pi/P$ is the angular speed of revolution of either star about the center of mass. Show that Newton's second law, applied to either star, leads to

$$m_1 + m_2 = \frac{\Omega^2 r^3}{G}$$

This should look familiar: it is Kepler's third law, in a slightly different form than usual.

- (c) Show that if the orbital axis (the line through the center of mass, perpendicular to the plane of the orbits) is inclined by an angle i with respect to our line of sight, then the maximum Doppler velocities that will be observed for the two stars relative to the Doppler velocity of their center of mass are $v_{1r} = \Omega r_1 \sin i$ and $v_{2r} = \Omega r_2 \sin i$, respectively.
- (d) Eliminate r_1 and r_2 from the previous expressions to show that

$$r = \frac{v_{1r} + v_{2r}}{\Omega \sin i} \quad \text{and} \quad \frac{m_2}{m_1} = \frac{v_{1r}}{v_{2r}}$$

Note that the right-hand side of each expression contains only observables. With these, and the results of part b, measurements can be used to obtain the masses and separations of the stars.

- (e) Suppose that a certain binary is eclipsing, and you have measured the period to be 11 days and the radial velocities to be $v_{1r} = 75 \text{ km/s}$ and $v_{2r} = 100 \text{ km/s}$. What are the masses (in M_\odot) and separation (in R_\odot) of the two stars?
3. *Single-line spectroscopic binaries and the "mass function."* Consider the binary star system from the previous problem.

- (a) From the equations you derived in the previous problem, show that the sum of the stellar masses is given in terms of the radial velocities by

$$m_1 + m_2 = \frac{P}{2\pi G} \frac{(v_{1r} + v_{2r})^3}{\sin^3 i}$$

- (b) Next, show that

$$\frac{m_2^3}{(m_1 + m_2)^2} \sin^3 i = \frac{P}{2\pi G} v_{1r}^3 \quad (1)$$

In the right hand side, only the observable quantities P and v_{1r} appear. This quantity, $f(m_1, m_2) = P v_{1r}^3 / 2\pi G$, is called the *mass function*.

- (c) Show that the left hand side of Eqn. 1 is always less than m_2 ; in other words, m_2 is always greater than the mass function.

Note: Equation 1 is useful for single-line spectroscopic binaries: those in which one of the stars (1) is much brighter than the other, so that only its spectral lines are seen. In particular, the lower limit to the mass of the unseen companion, m_2 , obtained in this manner has been useful in the identification of black holes in binary stellar systems, as we will see in a couple of weeks.