

Giant Molecular Clouds

Observed properties
Dynamical State
Formation

September 20, 2022

University of Rochester

The components of the ISM

Diffuse ISM neutral atomic clouds
embedded in ionized
medium

Dense ISM neutral molecular
clouds

Ionized nebulae HII regions,
planetary nebulae,
SNRs



ρ Ophiuchi cloud complex (NASA/WISE).


The components of the ISM: Dense ISM

Dense ISM: H, neutral and mostly molecular instead of atomic, in the form of clouds with densities $n_{\text{H}_2} = 10 - 10^6 \text{ cm}^{-3}$. Temperature is $10 - 100 \text{ K}$, mass is $10^3 - 10^6 M_{\odot}$.

- ▶ As much mass (total in Galaxy) as the diffuse ISM, $10^9 - 10^{10} M_{\odot}$, but volume is small in comparison.
- ▶ **Molecular cloud complexes** are usually physically connected to complexes of diffuse atomic clouds.
- ▶ The visual extinction through a molecular cloud is $\gg 1$.
- ▶ Best spectra line tracers are rotational lines of CO. H_2 radiates too poorly and is excited too inefficiently to be an effective tracer.
- ▶ So far, 233 molecular species have been detected in interstellar clouds. Smallest molecule: H_2 ; largest: C_{70} fullerene.

Giant molecular clouds

Taurus molecular cloud (*Herschel*)



Molecular clouds: Internal structure

- ▶ Molecular clouds are **clumpy**; they tend to consist of denser clumps in a range of sizes and masses, blending into a less-dense background.
- ▶ Molecular clouds are **cold**; they tend to have $T < 20$ K.
- ▶ Molecular clouds are **turbulent**: the random internal velocities are typically ~ 1 km/s, much larger than the average molecular speeds in a quiescent gas in equilibrium at the same temperature.
- ▶ Molecular clouds (and clumps) generally **rotate** slowly.
- ▶ Molecular clouds are **magnetized**: they are threaded by the same magnetic flux present when the material was diffuse and atomic. Now it has been compressed to a much smaller size, and the fields are correspondingly larger.


Mass distribution in GMCs

$$\frac{dN}{d \ln M} = N_u \left(\frac{M}{M_u} \right)^{\zeta} \quad M \lesssim M_u$$

where $0.3 < \zeta < 0.9$ and
 $M_u \sim 5 \times 10^6 M_{\odot}$.

In GMC cores, $0.9 < \zeta < 1.5$ with
 $M \gtrsim 1 M_{\odot}$.

GMC mass distribution in the inner MW
(Rosolowsky, 2005)



GMC scaling relations

In the range $10^2 M_{\odot} \lesssim M \lesssim 10^6 M_{\odot}$,

$$M \propto R^2$$

and

$$\Delta v \propto R^{1/2} \propto \rho^{-1/2}$$

Molecular clouds & star formation

- ▶ Clumps get massive enough and cold enough that the gas pressure cannot hold up their weight, and they collapse.
 - ▶ As clumps collapse, they heat in their cores.
 - ▶ If they are sufficiently massive and collapse to a small enough scale, temperatures can reach the fusion ignition point and a star is formed.
- ▶ The clumps are constantly being rearranged, compressed, or distended by turbulence.
- ▶ Collapse does not happen with spherical symmetry; often collapse is easier along the axis of rotation because of centrifugal forces.

Jeans criterion & Jeans mass

Critical wavelength (or critical wavenumber) between a stable and unstable equilibrium in a gas cloud supported by internal gas pressure:

$$\lambda_J = c_x \left(\frac{\pi}{G\rho_0} \right)^{1/2} \quad k_J = \frac{2\pi}{\lambda_J}$$

Any perturbation with $\lambda > \lambda_J$ (or $k < k_J$) is unstable and will result in a runaway gravitational collapse of the gas cloud.

Critical mass

The corresponding critical mass contained within a sphere with diameter λ_J is the **Jeans mass**.

$$M_J = \frac{\pi^{5/2}}{6} \frac{c_s^3}{(G^3 \rho_0)^{1/2}} \simeq 40 M_\odot \left(\frac{c_s}{0.2 \text{ km/s}} \right)^3 \left(\frac{n_{H_2}}{100 \text{ cm}^{-3}} \right)^{-1/2}$$

A GMC will collapse under its own gravity if its mass exceeds the Jeans mass.

For an isothermal sphere in pressure equilibrium with its surroundings, the critical mass is the **Bonner-Ebert mass**,

$$M_{BE} = 1.182 \frac{c_s^3}{(G^3 \rho_0)^{1/2}}$$

Since both GMCs and molecular clumps have masses exceeding both the Jeans mass and the Bonner-Ebert mass, they should collapse under their own gravity *if* they are supported by nothing more than thermal pressure.

Free-fall time & non-thermal pressures

The timescale within which they will collapse is equal to the **free-fall time**,

$$t_{\text{ff}} = \sqrt{\frac{3\pi}{32G\rho_0}} \simeq 3.6 \times 10^6 \text{ yr} \left(\frac{n_{\text{H}_2}}{100 \text{ cm}^{-3}} \right)^{-1/2}$$

Turbulence can provide additional pressure support if mean square velocity, σ , is high enough. Replace sound speed with effective sound speed,

$$c_{s,\text{eff}}^2 = c_s^2 + \sigma^2$$

Magnetic fields are also a potential non-thermal pressure source. If a cloud's potential energy is equal to the magnetic energy, then the critical mass with a uniform magnetic field is

$$M_{\Phi} \equiv \frac{5^{3/2}}{48\pi^2} \frac{B^3}{G^{3/2}\rho_0^2} \simeq 1.6 \times 10^5 M_{\odot} \left(\frac{n_{\text{H}_2}}{100 \text{ cm}^{-3}} \right)^{-2} \left(\frac{B}{30\mu\text{G}} \right)^3$$