

# Galactic star formation

The formation of massive stars

Empirical star formation laws

The IMF

The formation of Pop III stars

September 27, 2022

University of Rochester

# Massive star formation

Nuclear fusion can ignite before accretion process is complete, possibly significantly altering gas accretion due to resulting radiation pressure.

Balancing a star's gravitational force with the force from radiation pressure defines the **Eddington luminosity**,

$$L_E = \frac{4\pi cGM_*}{\kappa_d}$$

where  $\kappa_d \equiv$  the dust opacity per unit mass.

# Kennicutt-Schmidt Law

**Schmidt Law** (1959) relates  $\dot{\Sigma}_*$  and the gas surface density:

$$\dot{\Sigma}_* \propto \sigma_{\text{gas}}^N$$

where  $N \sim 1 - 2$ .


This was extended by Kennicutt (1998) to also include starburst galaxies, finding

$$\dot{\Sigma}_* = (2.5 \pm 0.7) \times 10^{-4} \left( \frac{\Sigma_{\text{gas}}}{M_{\odot} \text{pc}^{-2}} \right)^{1.4 \pm 0.15} M_{\odot} / \text{yr} / \text{kpc}^2$$

where  $\Sigma_{\text{gas}} = \Sigma_{\text{HI}} + \Sigma_{\text{H}_2}$ .

# Kennicutt-Schmidt Law


The Kennicutt-Schmidt law implies that a galaxy's SFR is controlled by the self-gravity of the gas.



# Kennicutt-Schmidt Law

Kennicutt (1998) also observed a similar correlation between  $\dot{\Sigma}_*$  and  $\Sigma_{\text{gas}}/\tau_{\text{dyn}}$ :

$$\dot{\Sigma}_* \approx 0.017 \Sigma_{\text{gas}} \Omega$$




# SF thresholds

Possible explanations for lack of star formation in outer disk:



**Gravitational instability** If the gas surface density is too low, then the gas will be stable to gravitational collapse.

**Shear instability** Shear from differential rotation prevents gravitational perturbations from growing.

**Ability to form  $H_2$**  If a region's pressure is too low, molecular H will not form.



# Local SF laws



# IMF

The Initial Mass Function,  $\phi(m) dm$ , describes the relative number of stars formed with masses  $m \pm 0.5dm$ . It is assumed to be continuous, and can be normalized so that

$$\int_{m_\ell}^{m_u} m\phi(m) dm = 1M_\odot$$

Typically,  $m_\ell \simeq 0.08M_\odot$  and  $m_u \simeq 100M_\odot$ .

For a total mass  $M_*$  of newly formed stars,

$$dN(m) = M_*[M_\odot]\phi(m) dm \quad dM(m) = M_*[M_\odot]m\phi(m) dm$$

The logarithm IMF is also often used, where  $\xi(m) d \log m = \phi(m) dm$ . The slopes of the IMF are

$$b(m) \equiv -\frac{d \log \phi}{d \log m} \quad \beta(m) \equiv -\frac{d \log \xi}{d \log m} = b - 1$$



# Constructing the IMF from observations

To define the observed IMF, one must estimate the stellar mass of a collection of stars within some volume from their luminosity.

Along the way, this will also produce the luminosity function,  $\Phi(M_r)$ , the number density of stars as a function of absolute magnitude.


If the IMF is time-independent, then the luminosity function is

$$\Phi_0(M_r) = \Phi(M_r) \frac{\int_0^{t_0} \psi(t) dt}{\int_{t_0 - \tau_{\text{MS}}(M_r)}^{t_0} \psi(t) dt}$$

where  $\psi(t)$  is the star formation history in the given volume. The IMF is then

$$\phi(m) \propto \frac{dM_r}{dm} \Phi_0[M_r(m)]$$

# Various IMFs from observations



# (Non-)universality of the IMF

Our knowledge of the IMF and the processes involved is too incomplete to definitively state whether or not the IMF is universal. Possible reasons to expect it to be non-universal include

**Gas-phase metallicity** The radiative cooling rate of gas depends on the chemical composition. Low metallicity regions could produce top-heavy IMFs.

**Ambient radiation** FIR photons can raise both the minimum temperature and the Jeans mass in star-forming clouds, resulting in a top-heavy IMF.