

# Clumps With Bounded Magnetic Fields, Their Interaction With Shocks, and the Effect of Magnetic Thermal Conduction



Shule Li, Adam Frank, Eric Blackman

*University of Rochester, Department of Physics and Astronomy. Rochester, New York 14627*

## Introduction

Problems involving magnetized clouds and clumps, especially their interaction with shocks are common in astrophysical environments and have been a topic of research in the past decade. The magnetic field structure, whether aligned with the shock or perpendicular to the shock, can have profound influence on the shocked behavior and evolution of the clump. In this presentation we review some basic results of the shocked MHD clumps by past simulations, as well as movies of preliminary 3D simulations produced by our parallel MHD code AstroBEAR. We will also discuss future directions of numerical simulations on such topic.

## Outline

1. Introduction
2. Previous work
3. Simulations of MHD clumps with external field
4. Future Directions



## Meaningful Quantities

Density Contrast:  $\chi = \frac{\rho_{clump}}{\rho_{amb}}$


Sonic Mach number:  $M = \frac{u_{wind}}{c_{s,amb}}$

Alfvenic Mach number:  $M_A = \frac{u_{wind}}{v_a}$


Magnetic Beta:  $\beta = \frac{2}{\gamma} \frac{c_{s,amb}^2}{v_a^2}$

Clump crushing time:  $\tau_{cc} = \frac{\chi^{1/2} r_{clump}}{u_{wind}}$

Jones, T.W., Ryu, Dongsu, Tregillis, I.L.  
1996 ApJ, 473, 365



Density log plot at  $2\tau_{cc}$  (top) and  $6\tau_{cc}$  (bottom). Left:  $\beta = 4$ . Right:  $\beta = 256$ .



Field line plot at  $2\tau_{cc}$  (top) and  $6\tau_{cc}$  (bottom). Left:  $\beta = 4$ . Right:  $\beta = 256$ .

### Perpendicular field evolution:

The magnetic field is stretched and wrapped around the clump, which effectively confines the clump and prevents its fragmentation, even for moderately strong field  $\beta = 4$ . The clump embedded in the stretched field is compressed, but then, because of the strong confining effect of the field develops a streamlined profile and is not strongly eroded.

The magnetic pressure at the nose of the clump increases drastically, which acts as a shock absorber.

At later stage, the stretched field around the clump edge has  $\beta < 1$  even for moderately strong initial field condition. This protects the bullet from further disruption.



**Jones, T.W., Ryu, Dongsu, Tregillis, I.L.  
1996 ApJ, 473, 365**


**Aligned field evolution:**

No significant difference in terms of density evolution even for  $\beta = 1$ .


The criterion for the magnetic field removing K-H instabilities at the clump boundaries is:  $\beta < 1$  along the boundary. The criterion for the magnetic field to stabilize R-T instabilities is roughly:  $\beta < \chi/M$ .

But there is no place that the magnetic field becomes energetically dominant; that is, almost everywhere  $\beta \gg 1$ .


Magnetic fields stretched over the top of the bullet can have a significant stabilizing influence that improves the flow coherence.



Density log plot at  $2\tau_{cc}$  (top) and  $6\tau_{cc}$  (bottom). Left:  $\beta = 4$ . Right:  $\beta = 256$ .




Field line plot at  $2\tau_{cc}$  (top) and  $6\tau_{cc}$  (bottom). Left:  $\beta = 4$ . Right:  $\beta = 256$ .



3D density contour at for weak and strong aligned field.

**Min-Su Shin, James M. Stone, Gregory F. Snyder, 2008 ApJ, 680, 336**




3D density contour (left) and magnetic pressure (right) at for weak aligned field, perpendicular field, and oblique shock cases, at  $4\tau_{cc}$ .



**Cunningham, Andrew J.; Frank, Adam; Varnière, Peggy;  
Mitran, Sorin; Jones, Thomas W. 2009 ApJS, 182, 519**

2D simulations done by  
AstroBEAR 1.0. The  
logarithm of the density  
distribution is presented in  
gray-scale, red lines  
delineate the magnetic field  
lines and blue lines show  
regions of AMR-enhanced  
resolution.



### 3D Contour of log Density


I'll show movies of 3D contour plot of log density here, for  $\beta = 2$  and  $\beta=0.5$ , with aligned and perpendicular field cases. So there are four movies here.




## **Magnetic Pressure and Field Line Structure**

I'll show movies of 2D cut of magnetic pressure and field lines here, for the same four cases on page 8.

## Future Work I: Contained field cases



The density evolution of contained field case remains similar to that of the hydro case in terms of morphology. The strong contained field does have a resistance that holds the clump material together at  $t = 2\tau_{cc}$ , the effect is weak comparing to the global field case of a greater  $\beta$ .



In the weak bounded field (top), field lines compressed and then unfold. The toroidal component becomes dominating. In the strong contained field case (bottom), the fields retain more of original configuration and provide some restriction of clump deformation.



## Future Work II: Multiphysics processes



Flux limited anisotropic thermal conduction

Microphysical Resistivity

Viscosity

I may show movies of MHD clump with resistivity here.

The Current Sheet Reconnection Outflow Test



## Summary