

Potential from charge moving with constant velocity

$$\vec{r}_0(t) = \vec{v}t$$

find retarded time $t' = t - \frac{|\vec{r} - \vec{r}_0(t')|}{c} \Rightarrow c(t - t') = |\vec{r} - \vec{r}_0(t')|$

$$\rightarrow c^2(t - t')^2 = |\vec{r} - \vec{r}_0(t')|^2 = r^2 + v^2 t'^2 - 2\vec{r} \cdot \vec{v} t'$$

$$c^2 t^2 + c^2 t'^2 - 2c^2 t t' = r^2 + v^2 t'^2 - 2\vec{r} \cdot \vec{v} t'$$

quadratic equation in t'

$$\rightarrow (c^2 - v^2)t'^2 - 2(c^2 t - \vec{r} \cdot \vec{v})t' + c^2 t^2 - r^2 = 0$$

quadratic formula

$$t' = \frac{(c^2 t - \vec{r} \cdot \vec{v}) \pm \sqrt{(c^2 t - \vec{r} \cdot \vec{v})^2 - (c^2 t^2 - r^2)(c^2 - v^2)}}{(c^2 - v^2)} \quad (*)$$

to get sign correct, note that when $v=0$, we want

$$t' = t - \frac{r}{c}$$

Apply to above:

$$t' = \frac{c^2 t \pm \sqrt{c^4 t^2 - c^4 t^2 + c^2 r^2}}{c^2}$$

$$= t \pm \frac{r}{c} \Rightarrow \underline{\underline{\text{take } \ominus \text{ sign}}}$$

scalar potential $V(\vec{r}, t) = \frac{q}{4\pi\epsilon_0} \frac{1}{|\vec{r} - \vec{r}_0(t')| (1 - \frac{1}{c} \hat{n}(t') \cdot \vec{v}(t'))}$

here $\vec{v}(t') = \vec{v}$ constant

$$\underbrace{|\vec{r} - \vec{r}_0(t')|}_{= c(t - t')} \left(1 - \frac{1}{c} \hat{n}(t') \cdot \vec{v}\right) = \underbrace{|\vec{r} - \vec{v}t'|}_{= c(t - t')} - \frac{1}{c} (\vec{r} - \vec{v}t') \cdot \vec{v}$$

$$= c(t - t') - \frac{1}{c} \vec{r} \cdot \vec{v} + \frac{v^2 t'}{c} = ct - \frac{\vec{r} \cdot \vec{v}}{c} - c(1 - \frac{v^2}{c^2})t'$$

$$= \frac{1}{c} \left[c^2 t - \vec{r} \cdot \vec{v} - (c^2 - v^2) t' \right] \quad \text{insert (*) for } t'$$

$$= \frac{1}{c} \left[\underbrace{c^2 t - \vec{r} \cdot \vec{v} - (c^2 t - \vec{r} \cdot \vec{v})}_{\text{cancel}} + \sqrt{(c^2 t^2 - \vec{r} \cdot \vec{v})^2 - (c^2 t - \vec{r} \cdot \vec{v})(c^2 - v^2)} \right]$$

$$= \frac{1}{c} \sqrt{(c^2 t - \vec{r} \cdot \vec{v})^2 - (c^2 t^2 - r^2)(c^2 - v^2)}$$

simplify the square root

$$= \frac{1}{c} \sqrt{c^4 t^2 + (\vec{r} \cdot \vec{v})^2 - 2c^2 t \vec{r} \cdot \vec{v} - c^4 t^2 + r^2 c^2 + c^2 v^2 t^2 - r^2 v^2}$$

$$= \frac{1}{c} \sqrt{c^2 (\vec{r} - \vec{v}t)^2 + (\vec{r} \cdot \vec{v})^2 - r^2 v^2}$$

$$= \sqrt{(\vec{r} - \vec{v}t)^2 + \frac{(\vec{r} \cdot \vec{v})^2}{c^2} - \frac{r^2 v^2}{c^2}}$$

$$V(\vec{r}, t) = \frac{q}{4\pi\epsilon_0} \frac{1}{\sqrt{(\vec{r} - \vec{v}t)^2 + \frac{(\vec{r} \cdot \vec{v})^2}{c^2} - \frac{r^2 v^2}{c^2}}}$$

Note, for $(\frac{v}{c})^2 \ll 1$

$$V(\vec{r}, t) \approx \frac{q}{4\pi\epsilon_0} \frac{1}{\sqrt{(\vec{r} - \vec{v}t)^2}} = \frac{q}{4\pi\epsilon_0} \frac{1}{|\vec{r} - \vec{r}_0(t)|^2}$$

"quasi static" limit
looks like instantaneous
Coulomb potential

vector potential

$$\vec{A}(\vec{r}, t) = \frac{\vec{v}}{c^2} V(\vec{r}, t)$$

fields from charges moving with constant $\vec{v}$

$$\vec{E} = -\vec{\nabla}V - \frac{\partial \vec{A}}{\partial t} = -\vec{\nabla}V - \frac{\vec{v}}{c^2} \frac{\partial V}{\partial t} \quad \text{as } \vec{A} = \frac{\vec{v}}{c^2} V$$

$$-\vec{\nabla}V = \frac{q}{4\pi\epsilon_0} \frac{(\frac{1}{2})}{(\quad)^{3/2}} \left[\vec{\nabla}(\vec{r}-\vec{v}t)^2 + \frac{1}{c^2} \vec{\nabla}(\vec{r}\cdot\vec{v})^2 - \frac{v^2}{c^2} \vec{\nabla}r^2 \right]$$

$$= \frac{q}{4\pi\epsilon_0} \frac{1}{(\quad)^{3/2}} \left[\vec{r}-\vec{v}t + \frac{\vec{v}}{c^2} (\vec{r}\cdot\vec{v}) - \frac{v^2}{c^2} \vec{r} \right]$$

$$-\frac{\partial V}{\partial t} = \frac{q}{4\pi\epsilon_0} \frac{(\frac{1}{2})}{(\quad)^{3/2}} \left[\frac{\partial}{\partial t} (\vec{r}-\vec{v}t)^2 \right] = \frac{q}{4\pi\epsilon_0} \frac{(-1)}{(\quad)^{3/2}} \left[(\vec{r}-\vec{v}t) \cdot \vec{v} \right]$$

$$\vec{E} = -\vec{\nabla}V - \frac{\vec{v}}{c^2} \frac{\partial V}{\partial t} = \frac{q}{4\pi\epsilon_0} \frac{1}{(\quad)^{3/2}} \left[\vec{r}-\vec{v}t + \frac{\vec{v}}{c^2} (\vec{r}\cdot\vec{v}) - \frac{v^2}{c^2} \vec{r} - \frac{\vec{v}}{c^2} (\vec{r}\cdot\vec{v} - v^2 t) \right]$$



$$\boxed{\vec{E}(\vec{r},t) = \frac{q}{4\pi\epsilon_0} \frac{(\vec{r}-\vec{v}t)(1-v^2/c^2)}{((\vec{r}-\vec{v}t)^2 + (\frac{\vec{r}\cdot\vec{v}}{c})^2 - (\frac{rv}{c})^2)^{3/2}}}$$

$$\vec{B} = \vec{\nabla} \times \vec{A} = \vec{\nabla} \times \left(\frac{\vec{v}}{c^2} V \right) = (\vec{\nabla} V) \times \frac{\vec{v}}{c^2} = (-\vec{E} - \frac{\partial \vec{A}}{\partial t}) \times \frac{\vec{v}}{c^2}$$

$$= -\vec{E} \times \frac{\vec{v}}{c^2} - \frac{\partial}{\partial t} \left(\frac{\vec{v}}{c^2} V \right) \times \frac{\vec{v}}{c^2} = \frac{\vec{v}}{c^2} \times \vec{E} - \frac{\partial V}{\partial t} \underbrace{\frac{\vec{v}}{c^2} \times \frac{\vec{v}}{c^2}}_{=0}$$

$$\boxed{\vec{B}(\vec{r},t) = \frac{\vec{v}}{c^2} \times \vec{E}(\vec{r},t)}$$

Note: The factor $1-v^2/c^2$ reminds one of special relativity!



for $v^2 \ll c^2$

$$\vec{E}(\vec{r}, t) \approx \frac{q}{4\pi\epsilon_0} \frac{(\vec{r} - \vec{v}t)}{|\vec{r} - \vec{v}t|^3} \quad \text{instantaneous Coulomb field}$$

$$\begin{aligned} \vec{B}(\vec{r}, t) &= \frac{\vec{v}}{c^2} \times \vec{E} = \frac{1}{4\pi\epsilon_0 c^2} \frac{q\vec{v} \times (\vec{r} - \vec{v}t)}{|\vec{r} - \vec{v}t|^3} \\ &= \frac{\mu_0}{4\pi} \int d^3r' \frac{\vec{j}(\vec{r}') \times (\vec{r} - \vec{r}')}{|\vec{r} - \vec{r}'|^3} \end{aligned}$$

$$\text{with } \vec{j}(\vec{r}') = q\vec{v} \delta(\vec{r}' - \vec{v}t)$$

so $\vec{B}$ looks just like Biot-Savart law, even though for moving charge, $\vec{v} \cdot \vec{j} \neq 0$.

Since a charge moving with constant $\vec{v}$ can always be considered to be at rest in another ~~the~~ frame of reference, the formulas above for $\vec{E}$ and $\vec{B}$ for charge moving with $\vec{v}$, will tell us how $\vec{E}$ and $\vec{B}$ transform under a Lorentz transformation between two frames in relative uniform motion. ~~This~~ In fact, the Lorentz transf so defined, was first discovered as a law of electrodynamics, before the theory of special relativity was discovered.

For a pt charge in state of arbitrary motion ($\vec{v} \neq 0$)
one finds (see text)

$$\vec{E}(\vec{r}, t) = \frac{q}{4\pi\epsilon_0} \frac{|\vec{r} - \vec{r}_0(t')|}{\left[|\vec{r} - \vec{r}_0(t')| \cdot [c\hat{m}(t') - \vec{v}(t')] \right]^3} \left[[c\hat{m}(t') - \vec{v}(t')] (c^2 - v^2) + (\vec{r} - \vec{r}_0(t')) \times ([c\hat{m}(t') - \vec{v}(t')] \times \vec{a}(t')) \right]$$

where $t' = t - \frac{|\vec{r} - \vec{r}_0(t')|}{c}$ is retarded time

$$\hat{m}(t') = \frac{\vec{r} - \vec{r}_0(t')}{|\vec{r} - \vec{r}_0(t')|} \quad \vec{v} = \frac{d\vec{r}_0}{dt}, \quad \vec{a} = \frac{d^2\vec{r}_0}{dt^2}$$

first term $\sim \frac{1}{r^2}$ is "generalized Coulomb field" or "velocity field"

second term $\sim \frac{1}{r}$ is "acceleration field"

$$\vec{B}(\vec{r}, t) = \frac{\hat{m}(t')}{c} \times \vec{E}(\vec{r}, t)$$

$\vec{B}$ always $\perp \vec{E}$, and $\vec{B} \perp \hat{m}$, vector from observer at $\vec{r}$ to retarded point $\vec{r}_0(t')$.

Can use above to write down total force between two point charges, in arbitrary states of motion (see text)

Can use to construct $\vec{S}$ and get power radiated by an accelerating point charge - in non-relativistic limit, one recovers Larmor's formula - see text