

## Electrons in a crystal

When we discussed X-ray scattering we saw that the condition of constructive interference required that an incident wavevector  $\vec{k}$  could only scatter into wavevectors  $\vec{k}' = \vec{k} - \vec{K}$  where  $\vec{K}$  was in the R.L.

For electrons in a crystal we expect similar behavior since our argument depended only on the wave nature of the scattering.

So a free electron plane wave state  $e^{i\vec{k}\cdot\vec{r}}$  will get scattered into states  $e^{i\vec{k}'\cdot\vec{r}}$  with  $\vec{k}' = \vec{k} - \vec{K}$ . We therefore expect the eigenstates of an electron in the periodic conic potential of the crystal will be a mixture of these scattered states

$$\begin{aligned}\psi_k(\vec{r}) &= \sum_{\vec{K}} e^{i(\vec{k}-\vec{K})\cdot\vec{r}} c_{\vec{k}-\vec{K}} \\ &= e^{i\vec{k}\cdot\vec{r}} \sum_{\vec{K}} e^{-i\vec{K}\cdot\vec{r}} c_{\vec{k}-\vec{K}} \\ &= e^{i\vec{k}\cdot\vec{r}} u_k(\vec{r}) \quad \text{where } u_k(\vec{r}) = \sum_{\vec{K}} e^{-i\vec{K}\cdot\vec{r}} c_{\vec{k}-\vec{K}}\end{aligned}$$

$u_k(\vec{r})$  is periodic on the Bravais lattice:

$$\begin{aligned}u(\vec{r} + \vec{R}) &= \sum_{\vec{K}} e^{-i\vec{K}\cdot(\vec{r} + \vec{R})} c_{\vec{k}-\vec{K}} = \sum_{\vec{K}} e^{-i\vec{K}\cdot\vec{r}} e^{-i\vec{K}\cdot\vec{R}} c_{\vec{k}-\vec{K}} \\ &= \sum_{\vec{K}} e^{-i\vec{K}\cdot\vec{r}} c_{\vec{k}-\vec{K}} = u_k(\vec{r}) \quad \text{since } e^{-i\vec{K}\cdot\vec{R}} = 1 \\ &\quad \text{for all } \vec{R} \text{ in R.L.}\end{aligned}$$

This then yields Bloch's Theorem which we prove more rigorously later.

The energy eigenstates of an electron in a potential with the periodicity of a Bravais lattice ~~can~~ can be written in the form

$$\psi_k(\vec{r}) = e^{i\vec{k} \cdot \vec{r}} u_k(\vec{r})$$

where  $u_k(\vec{r} + \vec{R}) = u_k(\vec{r})$  is periodic on the B.L.

An equivalent alternative statement of Bloch's Theorem is

$$\psi_k(\vec{r} + \vec{R}) = e^{i\vec{k} \cdot \vec{R}} \psi_k(\vec{r})$$

If the wavevector  $\vec{k}$  lies exactly on the Bragg plane bisected by R.L. vector  $\vec{R}$ , then the two free electron plane wave states  $\vec{k}$  and  $\vec{k}' = \vec{k} - \vec{R}$  are degenerate in energy and so we expect they must mix equally in forming the new eigenstates in the presence of the periodic potential. Since we start with two states  $\vec{k}, \vec{k}'$  we should wind up with two new states. If they mix equally, their relative coefficients should just be a phase factor  $e^{i\phi}$ . Hence we expect the two eigenstates to have the form

$$\psi_1(\vec{r}) = e^{i\vec{k} \cdot \vec{r}} + e^{i\phi_1} e^{i\vec{k}' \cdot \vec{r}}$$

$$\psi_2(\vec{r}) = e^{i\vec{k} \cdot \vec{r}} + e^{i\phi_2} e^{i\vec{k}' \cdot \vec{r}}$$

The new eigenstates  $\psi_1$  and  $\psi_2$  should be orthogonal

$$0 = \langle \psi_1 | \psi_2 \rangle = \int d^3r \psi_1^*(\vec{r}) \psi_2(\vec{r})$$


$$= \int d^3r \left[ e^{-i\vec{k}\cdot\vec{r}} + e^{-i\phi_1} e^{-i\vec{k}'\cdot\vec{r}} \right] \left[ e^{i\vec{k}\cdot\vec{r}} + e^{i\phi_2} e^{i\vec{k}'\cdot\vec{r}} \right]$$

$$= \int d^3r \left[ 1 + e^{i(\phi_2 - \phi_1)} + e^{-i\phi_1} e^{i(\vec{k} - \vec{k}')\cdot\vec{r}} + e^{i\phi_2} e^{i(\vec{k}' - \vec{k})\cdot\vec{r}} \right]$$

these two terms integrate to zero - proof later

$$= V \left[ 1 + e^{i(\phi_2 - \phi_1)} \right]$$

$$= 0 \text{ only if } \phi_2 - \phi_1 = \pi$$



$$\vec{k}' = \vec{k} - \vec{k}_\perp$$

write  $\vec{k} = \vec{k}_\parallel + \vec{k}_\perp$   
then  $\vec{k}' = -\vec{k}_\parallel + \vec{k}_\perp$

$$\text{So } \psi_1(\vec{r}) = e^{i\phi_1/2} e^{i\vec{k}_\perp \cdot \vec{r}} \left\{ e^{-i\phi_1/2} e^{i\vec{k}_\parallel \cdot \vec{r}} + e^{i\phi_2/2} e^{-i\vec{k}_\parallel \cdot \vec{r}} \right\}$$

$$= e^{i\phi_1/2} e^{i\vec{k}_\perp \cdot \vec{r}} 2 \cos(\vec{k}_\parallel \cdot \vec{r} - \phi_1/2)$$

Similarly

$$\psi_2(\vec{r}) = e^{i\phi_2/2} e^{i\vec{k}_\perp \cdot \vec{r}} 2 \cos(\vec{k}_\parallel \cdot \vec{r} - \phi_2/2)$$

$$= e^{i\phi_2/2} e^{i\vec{k}_\perp \cdot \vec{r}} 2 \cos(\vec{k}_\parallel \cdot \vec{r} - \frac{\phi_1}{2} - \frac{\pi}{2})$$

$$= e^{i\phi_2/2} e^{i\vec{k}_\perp \cdot \vec{r}} 2 \sin(\vec{k}_\parallel \cdot \vec{r} - \frac{\phi_1}{2})$$

So

$$\left. \begin{aligned} |\psi_1(\vec{r})|^2 &\sim \cos^2(\vec{k}_{||} \cdot \vec{r} - \phi/2) \\ |\psi_2(\vec{r})|^2 &\sim \sin^2(\vec{k}_{||} \cdot \vec{r} - \phi/2) \end{aligned} \right\} \text{probability density}$$


reflection from Bragg plane gives standing waves in direction  $\parallel$  to  $\vec{k}$

Since  $\vec{k}_{||} = \frac{\vec{k}}{2}$  we have  $\vec{k}_{||} \cdot \vec{R} = \frac{\vec{k} \cdot \vec{R}}{2} = \pi n$   
where  $n$  is an integer

$\Rightarrow |\psi_1(\vec{r})|^2$  and  $|\psi_2(\vec{r})|^2$  have the periodicity of the B.L.

$$\begin{aligned} \propto |\psi_1(\vec{r} + \vec{R})|^2 &\approx \cos^2(\vec{k}_{||} \cdot \vec{r} + \pi n - \phi/2) = \cos^2(\vec{k}_{||} \cdot \vec{r} - \phi/2) \\ &= |\psi_1(\vec{r})|^2 \end{aligned}$$

The only difference between  $\psi_1$  and  $\psi_2$  is their relative phase shift of  $\frac{\pi}{2}$ , which shifts the location of the peaks of  $|\psi_1|^2$  and  $|\psi_2|^2$  relative to the positions of the ions. By symmetry we expect one of these, say  $\psi_1$ , will have its peaks positioned at the same sites as the ions, while the other,  $\psi_2$ , will have its peaks positioned in between the sites of the ions. For example, if  $\vec{k}$  is the smallest non-zero R.L. vector,  $|\vec{k}| = \frac{2\pi}{a}$  with  $a$  = separation between ions, we expect,




The spatial periodicity of  $\psi_1$  and  $\psi_2$  is  $\lambda = \frac{2\pi}{k_{11}} = \frac{2\pi}{(K/2)}$   
 $= \frac{2\pi}{\pi/a} = 2a$ . The spatial periodicity of  $|\psi_1|^2$  and  $|\psi_2|^2$  is therefore  $\frac{1}{2}\lambda = a$ . Since  $\psi_1$  and  $\psi_2$  are  $\frac{\pi}{2}$  out of phase, then  $|\psi_1|^2$  and  $|\psi_2|^2$  are  $\pi$  out of phase, hence their relative peaks are separated by  $a/2$  as in the diagram above.

Now when one turns on the attractive ionic potential, the state  $\psi_1$  will have a lower energy than the state  $\psi_2$  since  $\psi_1$  has its peaks located at the ions, where the interaction is strongest, while  $\psi_2$  has its peaks in between the ions, where the interaction is weakest.

Hence we expect that the two states  $\psi_1$  and  $\psi_2$ , which were degenerate in energy before turning on the ionic potential, now split in energy, with a finite energy gap between them,  $\epsilon_1 < \epsilon_2$ .

⇒ Ionic potential cause a gap to appear in the energy dispersion relation  $E(\vec{k})$  whenever  $\vec{k}$  crosses a Bragg plane. Since the Brillouin zones are formed by the intersections of Bragg planes, gaps in  $E(\vec{k})$  open as  $\vec{k}$  crosses the boundary of any Brillouin zone.


For a 1-dimensional B.L. of lattice constant  $a$ . We expect the dispersion relation to look as in the sketch below:



$R = ma$   $m$  integer  
 $K = nb$ ,  $b = \frac{2\pi}{a}$ ,  $n$  integer  
 gap open in  $E(k)$   
 everytime  $k$  crosses  
 the boundary of a  
 Brillouin Zone


Bragg Planes are at  
 $\frac{K}{2} = \left(\frac{2\pi n}{a}\right)\left(\frac{1}{2}\right) = \frac{\pi n}{a}$   
 for  $n$  integers


For a 2-dimensional square B.L. of lattice constant  $a$



Bragg Planes divide  $k$ -space into Brillouin Zones

1st BZ 

2nd BZ 

3rd BZ 

whenever  $k$  crosses the surface of a BZ, there is a discrete jump in the energy  $E(k)$

Each BZ is a primitive cell of the R.L.

Each  $\vec{k}$  in  $k$ -space can be written as

$\vec{k} = \vec{g} + \vec{K}$  with  $\vec{K}$  a R.L. vector and  $\vec{g}$  a vector in the 1<sup>st</sup> BZ.  $\vec{g}$  is unique

$\Rightarrow$  each  $n^{\text{th}}$  BZ may be mapped onto the 1<sup>st</sup> BZ by translating its pieces by appropriate R.L. vectors  $\vec{K}$

It is customary to label the eigenstates and eigenvalues by the  $\vec{g}$  and by discrete index  $n$ .  $\vec{g}$  is called the "crystal momentum" and  $n$  the "band index." The state  $(\vec{g}, n)$  corresponds to the free electron state in the  $n^{\text{th}}$  BZ with wave vector  $\vec{k} = \vec{g} + \vec{K}$  ( $\vec{K}$  is the RL vector that translates  $\vec{g}$  into the  $n^{\text{th}}$  BZ)

The wavefunctions  $\psi_{\vec{g},n}$  and energies  $E_n(\vec{g})$  are called the band structure

## Born-von Karman boundary conditions and Fourier transforms for a Bravais lattice

We generalize the idea of periodic (or Born-von Karman) boundary conditions to electron states on a Bravais lattice.

B.L. vectors  $\vec{R} = n_1 \vec{a}_1 + n_2 \vec{a}_2 + n_3 \vec{a}_3$   
 $n_1, n_2, n_3$  integers.

For a BL of finite size we have  $0 \leq n_i < N_i$   
 $N = N_1 N_2 N_3$  is total number of points in the BL.  
Total volume of this finite BL is

$$V = \underbrace{\vec{a}_1 \cdot (\vec{a}_2 \times \vec{a}_3)}_{\text{volume of primitive cell}} N$$

We want our electron wave functions to be periodic on such a finite BL

$$\psi(\vec{r} + N_i \vec{a}_i) = \psi(\vec{r})$$

As we saw earlier for free electrons, this imposes a constraint on the wave vectors  $\vec{k}$  that can appear in the Fourier transform of  $\psi(\vec{r})$ .

Write  $\psi(\vec{r})$  in terms of its Fourier transform

$$\psi(\vec{r}) = \sum_{\vec{k}} e^{i \vec{k} \cdot \vec{r}} c_{\vec{k}}$$

↑ Fourier coefficients for  $\psi(\vec{r})$

Then

$$\begin{aligned}\psi(\vec{r} + N_i \vec{a}_i) &= \sum_{\vec{k}} e^{i\vec{k} \cdot (\vec{r} + N_i \vec{a}_i)} c_{\vec{k}} \\ &= e^{i N_i \vec{k} \cdot \vec{a}_i} \sum_{\vec{k}} e^{i\vec{k} \cdot \vec{r}} c_{\vec{k}} \\ &= e^{i N_i \vec{k} \cdot \vec{a}_i} \psi(\vec{r})\end{aligned}$$

$\Rightarrow$  must have  $e^{i N_i \vec{k} \cdot \vec{a}_i} = 1$  for all  $\vec{k}$   
that appear in Fourier transform of  $\psi(\vec{r})$

Write  $\vec{k}$  in terms of the primitive vectors of the RL

$$\vec{k} = x_1 \vec{b}_1 + x_2 \vec{b}_2 + x_3 \vec{b}_3 \quad x_i \text{ are not in general integers}$$

$$\text{Then } e^{i N_i \vec{k} \cdot \vec{a}_i} = e^{i N_i 2\pi x_i} = 1$$

$$\Rightarrow x_i N_i = m_i \text{ integer}$$

$$x_i = \frac{m_i}{N_i}$$


So the allowed wavevectors are

$$\boxed{\vec{k} = \frac{m_1}{N_1} \vec{b}_1 + \frac{m_2}{N_2} \vec{b}_2 + \frac{m_3}{N_3} \vec{b}_3}$$

where  $m_1, m_2, m_3$  are integers.

This is the Born - von Karman boundary conditions

2/3



volume per allowed wavevector in  $\vec{k}$ -space is the volume of the parallelepiped formed by the vectors  $\frac{\vec{b}_i}{N_i}$

$$= \frac{\vec{b}_1 \cdot (\vec{b}_2 \times \vec{b}_3)}{N_1 N_2 N_3} = \frac{(2\pi)^3}{N (\vec{a}_1 \cdot (\vec{a}_2 \times \vec{a}_3))}$$

$$= \frac{(2\pi)^3}{vN} = \frac{(2\pi)^3}{V}$$

$$V = vN \\ = \text{total volume BL}$$

where  $v = \vec{a}_1 \cdot (\vec{a}_2 \times \vec{a}_3)$  is the volume of the primitive cell of the B.L. and  $\frac{(2\pi)^3}{v}$  is the volume of the primitive cell of the R.L.

Note: volume per allowed wavevector  $\frac{(2\pi)^3}{v}$  is same as we had for free electrons.

Number of allowed  $\vec{k}$  values in any primitive cell of the R.L. is

$$\frac{\left[ \frac{(2\pi)^3}{v} \right]}{\left[ \frac{(2\pi)^3}{vN} \right]} = N \text{ number of cells in BL}$$

Number of allowed  $\vec{k}$  values in any primitive cell of the R.L., for example the 1<sup>st</sup> BZ, is  $N$