


monopole: $l=0$ term

$$\phi^{(0)}(\vec{r}) = \frac{1}{r} \int d^3r' f(r') \quad P_0(\cos\theta) = 1$$

$$= \frac{q}{r} \quad \text{where } q = \int d^3r' f(r') \text{ is } \underline{\text{total charge}}$$

dipole: $l=1$ term

$$\phi^{(1)}(\vec{r}) = \frac{1}{r^2} \int d^3r' f(\vec{r}') r' P_1(\cos\theta)$$



$$= \frac{1}{r^2} \int d^3r' f(\vec{r}') r' \cos\theta$$

$$\text{Now } \hat{r} \cdot \hat{r}' = r r' \cos\theta \Rightarrow \hat{r} \cdot \vec{r}' = r' \cos\theta$$

$$\phi^{(1)}(\vec{r}) = \frac{1}{r^2} \hat{r} \cdot \int d^3r' f(\vec{r}') \vec{r}'$$

$$= \frac{\vec{p} \cdot \hat{r}}{r^2} \quad \text{where } \vec{p} = \int d^3r' f(\vec{r}') \vec{r}'$$

is the dipole moment

For a set of point charges q_i at \vec{r}_i ,

$$\vec{p} = \sum_i q_i \vec{r}_i$$

quadrupole: $l=2$ term

$$\begin{aligned}\phi^{(2)}(\vec{r}) &= \frac{1}{r^3} \int d^3r' \rho(\vec{r}') r'^2 P_2(\cos\theta) \\ &= \frac{1}{r^3} \int d^3r' \rho(\vec{r}') r'^2 \frac{1}{2} (3\cos^2\theta - 1)\end{aligned}$$

use $\cos\theta = \hat{r}' \cdot \hat{r}$

$$\begin{aligned}\phi^{(2)}(\vec{r}) &= \frac{1}{r^3} \int d^3r' \rho(\vec{r}') \frac{1}{2} (3(\hat{r}' \cdot \hat{r})^2 - (r')^2) \\ &= \frac{1}{r^3} \hat{r} \cdot \left[\int d^3r' \rho(\vec{r}') \frac{1}{2} (3\hat{r}'\hat{r}' - (r')^2 \overset{\leftrightarrow}{\mathbf{I}}) \right] \cdot \hat{r}\end{aligned}$$

where $\overset{\leftrightarrow}{\mathbf{I}}$ is the identity tensor such that for any two vectors \vec{v} and \vec{u} , $\vec{u} \cdot \overset{\leftrightarrow}{\mathbf{I}} \cdot \vec{v} = \vec{u} \cdot \vec{v}$.

and $\hat{r}'\hat{r}'$ is the tensor such that for any two vectors \vec{v} and \vec{u} , $\vec{u} \cdot [\hat{r}'\hat{r}'] \cdot \vec{v} = (\vec{u} \cdot \hat{r}')(\hat{r}' \cdot \vec{v})$.

Define quadrupole tensor $\overset{\leftrightarrow}{\mathbf{Q}} \equiv \int d^3r' \rho(\vec{r}') (3\hat{r}'\hat{r}' - (r')^2 \overset{\leftrightarrow}{\mathbf{I}})$

$$\phi^{(2)}(\vec{r}) = \frac{1}{r^3} \frac{1}{2} \hat{r} \cdot \overset{\leftrightarrow}{\mathbf{Q}} \cdot \hat{r}$$

So to lowest three terms

$$\phi(\vec{r}) = \frac{q}{r} + \frac{\vec{p} \cdot \hat{r}}{r^2} + \frac{\hat{r} \cdot \overset{\leftrightarrow}{\mathbf{Q}} \cdot \hat{r}}{2r^3} + \dots$$

defined in terms of the moments q , \vec{p} , $\overset{\leftrightarrow}{\mathbf{Q}}$ of the charge distribution.

Note, the moments q , \vec{P} , \vec{Q} do not depend on the observation point \vec{r} - we can calculate them once and then use them to get $\phi(\vec{r})$ at all \vec{r} .

monopole: $q = \int d^3r \rho(\vec{r})$ scalar integral

dipole $\vec{P} = \int d^3r \rho(\vec{r}) \vec{r}$ vector integral
 $\hat{e}_1 \equiv \hat{x}, \hat{e}_2 \equiv \hat{y}, \hat{e}_3 \equiv \hat{z}$

if we pick a coordinate system, we have to do 3 integrations to get the three components of \vec{P}

$$\hat{e}_i \cdot \vec{P} = P_i = \int d^3r \rho(\vec{r}) r_i$$

quadrupole $\vec{Q} = \int d^3r \rho(\vec{r}) (3\vec{r}\vec{r} - r^2 \vec{I})$ tensor integral

if we pick a coord system x, y, z then


\vec{Q} is a matrix with components $\hat{e}_1 \equiv \hat{x}, \hat{e}_2 \equiv \hat{y}, \hat{e}_3 \equiv \hat{z}$

$$\hat{e}_i \cdot \vec{Q} \cdot \hat{e}_j = Q_{ij} = \int d^3r \rho(\vec{r}) [3r_i r_j - r^2 \delta_{ij}]$$


There are 9 elements of the 3×3 matrix Q_{ij} , but $Q_{ij} = Q_{ji}$ is symmetric so there are only 6 independent elements to compute.

General method

$$\phi(\vec{r}) = \sum_{l=0}^{\infty} \frac{1}{r^{l+1}} \int d^3r' \rho(\vec{r}') (r')^l P_l(\cos\theta)$$



in above, θ is angle between \vec{r} and \hat{z}
 if we think of θ as the spherical coord θ ,
 then in effect, above is choosing \vec{r} to be on
 \hat{z} axis. We would like a representation in
 which \vec{r} is positioned arbitrarily with respect
 to the axes used in describing ρ



Use the addition theorem for spherical harmonics
 - see Jackson 3.6 for discussion + proof

$$P_l(\cos\delta) = \frac{4\pi}{2l+1} \sum_{m=-l}^l Y_{lm}^*(\theta', \phi') Y_{lm}(\theta, \phi)$$

where (θ, ϕ) are the angles of \hat{r} , (θ', ϕ') are
 the angles of \hat{r}' , and δ is the angle
 between \hat{r} and \hat{r}' , i.e. $\cos\delta = \hat{r} \cdot \hat{r}'$

$$\cos\theta = \hat{z} \cdot \hat{r}$$

$$\cos\theta' = \hat{z} \cdot \hat{r}'$$

\Rightarrow

$$\phi(\vec{r}) = \sum_{l=0}^{\infty} \frac{1}{r^{l+1}} \frac{4\pi}{2l+1} \sum_{m=-l}^l \int d^3r' \rho(\vec{r}') (r')^l Y_{lm}^*(\theta', \phi') Y_{lm}(\theta, \phi)$$

Define the moment

$$f_{lm} \equiv \int d^3r' \rho(\vec{r}') (r')^l Y_{lm}^*(\theta', \phi')$$

independent of observation point

Then

$$\phi(\vec{r}) = 4\pi \sum_{l=0}^{\infty} \sum_{m=-l}^l \frac{f_{lm} Y_{lm}(\theta, \phi)}{(2l+1) r^{l+1}}$$

see Jackson eqn (4.4), (4.5), (4.6) to relate f_{lm} to q , \vec{p} , \vec{Q} .

$$\phi(\vec{r}) = \frac{q}{r} + \frac{\vec{p} \cdot \hat{r}}{r^2} + \frac{\hat{r} \cdot \vec{Q} \cdot \hat{r}}{2r^3}$$

electric field $\vec{E} = -\vec{\nabla}\phi = -\frac{\partial\phi}{\partial r} \hat{r} + \frac{1}{r} \frac{\partial\phi}{\partial\theta} \hat{\theta} + \frac{1}{r\sin\theta} \frac{\partial\phi}{\partial\phi} \hat{\phi}$

For the monopole term $\vec{E} = \frac{q}{r^2} \hat{r}$

For the dipole term, choose \vec{p} along \hat{z} axis so

$$\phi(\vec{r}) = \frac{p \cos\theta}{r^2}$$

$$\vec{E} = \frac{2p \cos\theta}{r^3} \hat{r} + \frac{p \sin\theta}{r^3} \hat{\theta}$$

$$\vec{E} = \frac{p}{r^3} (2\cos\theta \hat{r} + \sin\theta \hat{\theta})$$

note

$$p \cos\theta \hat{r} = (\vec{p} \cdot \hat{r}) \hat{r}$$

$$p \sin\theta \hat{\theta} = -(\vec{p} \cdot \hat{\theta}) \hat{\theta}$$


Now $\vec{p} = (\vec{p} \cdot \hat{r}) \hat{r} + (\vec{p} \cdot \hat{\theta}) \hat{\theta}$

$$\Rightarrow -(\vec{p} \cdot \hat{\theta}) \hat{\theta} = (\vec{p} \cdot \hat{r}) \hat{r} - \vec{p}$$

so

$$\vec{E} = \frac{1}{r^3} \left[2(\vec{p} \cdot \hat{r}) \hat{r} + (\vec{p} \cdot \hat{r}) \hat{r} - \vec{p} \right]$$

$$= \frac{1}{r^3} \left[3(\vec{p} \cdot \hat{r}) \hat{r} - \vec{p} \right] \quad \text{expresses } \vec{E} \text{ in coord}$$



$$\vec{E} = \frac{1}{r^3} [3(\vec{p} \cdot \hat{r})\hat{r} - \vec{p}]$$


expresses \vec{E} of dipole
in coord free form

Origin of coordinates

The definition of the multipole moments depends on
the choice of origin of the coordinates

Suppose transform to $\vec{r}' = \vec{r} - \vec{d}$

In the \vec{r}' coord system



$$\tilde{q} = \int d^3r' \rho(r') = \int d^3r \rho(r) = q$$

monopole does not depend on choice of origin

$$\tilde{\vec{p}} = \int d^3r' \rho(r') \vec{r}' = \int d^3r \rho(\vec{r} - \vec{d})$$

$$= \int d^3r \rho \vec{r} - \vec{d} \int d^3r \rho$$


$$\tilde{\vec{p}} = \vec{p} - \vec{d}q \quad \tilde{\vec{p}} = \vec{p} \text{ only if } q=0!$$

if $q \neq 0$, then $\tilde{\vec{p}} \neq \vec{p}$

~~One could~~ \Rightarrow If $q \neq 0$, one could always choose
an origin of coords for which $\vec{p} = 0!$

For HW you will show that $\tilde{\vec{p}} = \vec{p}$ only if both
 $q=0$ and $\vec{p}=0$.

Example two charges q_1 at \vec{r}_1 and q_2 at \vec{r}_2
 $q_1 + q_2 = q \neq 0$



monopole $q_1 + q_2 = q$
 dipole $\vec{p} = q_1 \vec{r}_1 + q_2 \vec{r}_2$
 quadrupole $\vec{Q} = (3\vec{r}_1 \vec{r}_1 - r_1^2 \vec{I}) q_1$
 $+ (3\vec{r}_2 \vec{r}_2 - r_2^2 \vec{I}) q_2$

We can make the dipole moment vanish by shifting to a new coord system $\vec{r}' = \vec{r} - \vec{d}$ where $\vec{d} = \frac{\vec{p}}{q}$

$$\vec{r}' = \vec{r} - \frac{q_1 \vec{r}_1 + q_2 \vec{r}_2}{q_1 + q_2} = \frac{q_1 (\vec{r} - \vec{r}_1) + q_2 (\vec{r} - \vec{r}_2)}{q_1 + q_2}$$

positions of q_1, q_2 in new coords are

$$\vec{r}'_1 = \frac{q_2}{q_1 + q_2} (\vec{r}_1 - \vec{r}_2)$$

$$\vec{r}'_2 = \frac{-q_1}{q_1 + q_2} (\vec{r}_1 - \vec{r}_2)$$

origin of new coord system is at

$$\vec{r}' = 0 \Rightarrow \vec{r} = \frac{q_1 \vec{r}_1 + q_2 \vec{r}_2}{q_1 + q_2}$$

lies along vector from \vec{r}_2 to \vec{r}_1

"center of charge"

for many charges q_i at positions \vec{r}_i , the origin that makes dipole moment vanish is at

$$\vec{r} = \frac{\sum_i q_i \vec{r}_i}{\sum_i q_i}$$

In this coord system

$$\vec{p}' = q_1 \vec{r}_1' + q_2 \vec{r}_2' = \frac{q_1 q_2}{q_1 + q_2} (\vec{r}_1 - \vec{r}_2) - \frac{q_2 q_1}{q_1 + q_2} (\vec{r}_1 - \vec{r}_2) \\ = 0 \quad \text{as it must be!}$$

Quadrupole moment in the coord system in which $\vec{p}' = 0$
the quadrupole tensor is

$$\vec{Q}' = [3\vec{r}_1' \vec{r}_1' - (r_1')^2 \vec{I}] q_1 + [3\vec{r}_2' \vec{r}_2' - (r_2')^2 \vec{I}] q_2$$

let us choose ~~coord~~ spherical coordinates with origin at O'
and \hat{z} axis aligned along $\vec{r}_1 - \vec{r}_2$, so that

$$\vec{r}_1 - \vec{r}_2 = s \hat{z} \quad \text{where } s = |\vec{r}_1 - \vec{r}_2| \text{ is separation} \\ \text{between the charges}$$

$$\text{then } \vec{r}_1' = \frac{q_2}{q_1 + q_2} s \hat{z}$$

$$\vec{r}_2' = \frac{-q_1}{q_1 + q_2} s \hat{z}$$

$$\vec{Q}' = \left(\frac{q_2}{q_1 + q_2} \right)^2 q_1 [3s^2 \hat{z} \hat{z} - s^2 \vec{I}] \\ + \left(\frac{-q_1}{q_1 + q_2} \right)^2 q_2 [3s^2 \hat{z} \hat{z} - s^2 \vec{I}]$$

$$\vec{Q}' = \frac{q_2^2 q_1 + q_1^2 q_2}{(q_1 + q_2)^2} s^2 [3 \hat{z} \hat{z} - \vec{I}]$$

$$= \frac{q_1 q_2}{q_1 + q_2} s^2 [3 \hat{z} \hat{z} - \vec{I}]$$

$$Q'_{ij} = \frac{q_1 q_2}{q_1 + q_2} s^2 \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 2 \end{pmatrix}$$

in xyz coord system

$$\text{as } \hat{z} \hat{z} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$\vec{I} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

The contribution of quadrupole to the potential is

$$\phi_{\text{quad}} = \frac{1}{2} \frac{\hat{r} \cdot \vec{Q}' \cdot \hat{r}}{r^3}$$

$$\hat{r} = \begin{pmatrix} \sin \theta \cos \varphi \\ \sin \theta \sin \varphi \\ \cos \theta \end{pmatrix}$$

with origin at O' this becomes

in xyz coords

$$\phi_{\text{quad}} = \frac{s^2}{2r^3} \frac{q_1 q_2}{q_1 + q_2} (\sin \theta \cos \varphi, \sin \theta \sin \varphi, \cos \theta) \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 2 \end{pmatrix} \begin{pmatrix} \sin \theta \cos \varphi \\ \sin \theta \sin \varphi \\ \cos \theta \end{pmatrix}$$

do matrix multiplications

$$\phi_{\text{quad}} = \frac{s^2}{2r^3} \frac{q_1 q_2}{q_1 + q_2} (2 \cos^2 \theta - \sin^2 \theta)$$

independent of φ as it must be due to azimuthal symmetry

Example

simple charge configs


• $q \Rightarrow$ monopole is leading term

$+q \quad -q \Rightarrow$ monopole $= 0 \Rightarrow$ dipole is leading term
 \vec{p} is indep of origin

$+q \quad -q$ \Rightarrow monopole $= 0 \Rightarrow$ total dipole is
sum of dipoles of individual neutral pairs

$\vec{p} = 0$

leading term is quadrupole



when monopole $= 0$ and dipole $= 0$,
quadrupole is indep of origin.
 \rightarrow total quadrupole is sum of
quadrupoles of individual
clusters with $q = 0$ and $\vec{p} = 0$

$$Q = Q_1 + Q_2$$

$$\text{with } Q_2 = -Q_1$$

$\Rightarrow Q = 0$ leading term is octopole