

Helmholtz Theorem

$$\text{Suppose } \left. \begin{aligned} \vec{\nabla} \cdot \vec{E}(\vec{r}) &= f(\vec{r}) \\ \vec{\nabla} \times \vec{E}(\vec{r}) &= \vec{g}(\vec{r}) \end{aligned} \right\} \text{ for } \vec{r} \text{ in a volume } V$$
$$\vec{E}(\vec{r}) = \vec{h}(\vec{r}) \quad \text{for } \vec{r} \text{ on surface } S \text{ of vol } V$$

Then if we know $f(\vec{r})$, $\vec{g}(\vec{r})$ and $\vec{h}(\vec{r})$, that information uniquely determines the vector function $\vec{E}(\vec{r})$

Proof:

Suppose we had two different solutions $\vec{E}(\vec{r})$ and $\vec{E}'(\vec{r})$
Then define

$$\vec{G}(\vec{r}) = \vec{E}(\vec{r}) - \vec{E}'(\vec{r})$$

$\vec{G}$ must satisfy

$$\left. \begin{aligned} \vec{\nabla} \cdot \vec{G} &= 0 \\ \vec{\nabla} \times \vec{G} &= 0 \end{aligned} \right\} \text{ for all } \vec{r} \text{ in } V$$

$$\vec{G} = 0 \quad \text{for all } \vec{r} \text{ on } S$$

Now $\vec{\nabla} \times \vec{G} = 0$ implies we can find a scalar function ϕ such that $\vec{G} = \vec{\nabla} \phi$. Then
 $\vec{\nabla} \cdot \vec{G} = 0 \Rightarrow \nabla^2 \phi = 0$ for all $\vec{r}$ in V .

A function ϕ that satisfies $\nabla^2 \phi = 0$ within a region V is said to be a harmonic function on V .

An important property of harmonic functions is that the value at a position $\vec{r}$, is equal to the average of the values on the surface of a sphere centered at $\vec{r}$.

$$\phi(\vec{r}) = \frac{1}{4\pi R^2} \oint_{\Sigma} d\vec{a}' \phi(\vec{r}')$$

Σ $\nwarrow$ surface of sphere of radius R centered at $\vec{r}$.

From this property we can conclude that a harmonic function on V can have no local maximum or minimum within the volume V . All maxima and minima must lie on surface S of V .

Proof: Just consider a small sphere centered on $\vec{r}$ that fits within the volume V . If $\vec{r}$ was a max., then for $\vec{r}'$ on surface of sphere, $\phi(\vec{r}') < \phi(\vec{r})$. But then we would have $\phi(\vec{r}) < \frac{1}{4\pi R^2} \oint da' \phi(\vec{r}')$ in violation of the above property of harmonic functions.

Back to our function $\vec{G}(\vec{r})$. We have

$$\vec{\nabla} \cdot \vec{G} = 0, \quad \vec{G} = \vec{\nabla} \phi \Rightarrow \nabla^2 \phi = 0 \text{ in } V$$

$$\vec{G} = \vec{\nabla} \phi = 0 \text{ on surface } S \text{ of } V \Rightarrow \phi = \text{constant on } S.$$

All max and min of ϕ must be on surface S

$$\Rightarrow \phi_{\max} = \phi_{\min} = \text{constant},$$

$$\Rightarrow \phi = \text{constant throughout volume } V$$

$$\Rightarrow \vec{\nabla} \phi = \vec{G} = 0 \text{ throughout } V$$

$$\Rightarrow \vec{E} = \vec{E}' \text{ for all } \vec{r} \text{ in } V$$

$\Rightarrow$ solution is unique!

Magnetostatics

Loentz Force

a charge q , in motion with velocity $\vec{v}$, feels the force

$$\vec{F} = q(\vec{E} + k_4 \vec{v} \times \vec{B}) \quad \leftarrow \text{Loentz force}$$

$\vec{B}$ is the magnetic field at the position of the charge.
 k_4 is a universal constant.

Just as the constant k_1 fixed the units of charge q , the constant k_4 can be viewed as fixing the units of B magnetic field. By choosing the units of q and B appropriately, we are free to choose any values for k_1 and k_4 .

Magnetic field $\vec{B}$ is generated by moving charge.
 A charge q' with velocity $\vec{v}'$ ($v' \ll c$) located at the origin $\vec{r}' = 0$ produces a magnetic field at position $\vec{r}$,

holds only non relativistically $\rightarrow \vec{B}(\vec{r}) = k_5 q' \frac{\vec{v}' \times \vec{r}}{r^3} = \frac{k_5}{k_1} \vec{v}' \times \vec{E}(\vec{r})$

k_5 is a universal constant. we will see that it cannot be chosen independently of k_1 and k_4 .
 (since k_1 fixed units of q , and k_4 fixed units of $\vec{B}$, there are no further new quantities whose units could be adjoined to allow us to fix k_5 arbitrarily)

(11)

The force on a charge q at position $\vec{r}$, moving with velocity $\vec{v}$, due to a charge q' at the origin moving with velocity $\vec{v}'$ is, in non-relativistic limit ($v, v' \ll c$),

$$\vec{F} = k_1 q q' \frac{\vec{r}}{r^3} + k_4 k_5 q q' \vec{v} \times \left(\frac{\vec{v}' \times \vec{r}}{r^3} \right)$$

↑
Coulomb force

↑
magnetic analog of Coulomb force

The magnetic part is just the point charge equivalent of the Biot-Savart law for the force between current carrying wires. If we regard $q\vec{v} = \vec{I}$ as the current of charge q , and $q'\vec{v}' = \vec{I}'$ as the current of charge q' , then the magnetic force is $k_4 k_5 \vec{I} \times (\vec{I}' \times \frac{\vec{r}}{r^3})$ which is the Biot-Savart Law.

Rewrite above force as

$$\vec{F} = k_1 \left(1 + \frac{k_4 k_5}{k_1} \vec{v} \times \vec{v}' \times \right) \frac{\vec{r}}{r^3} q q'$$

we see that $\left(\frac{k_4 k_5}{k_1} \right)$ has units of $(\text{velocity})^{-2}$

it must be independent of whatever convention one used to choose the units of q or B (ie independent of choices for k_1 and k_4). Experimentally it is found that

$$\left(\frac{k_4 k_5}{k_1} \right) = \frac{1}{c^2}$$

c = speed of light in vacuum

Continuum current density

For charges q_i at positions $\vec{r}_i(t)$ with $\vec{v}_i = \frac{d\vec{r}_i}{dt}$
we define the current density

$$\vec{j}(\vec{r}, t) = \sum_i q_i \vec{v}_i(t) \delta(\vec{r} - \vec{r}_i(t))$$

units of $\vec{j}$ are $(\text{charge}) \left(\frac{\text{length}}{\text{time}} \right) \left(\frac{1}{\text{length}^3} \right) = \left(\frac{\text{charge}}{\text{area} \cdot \text{time}} \right)$

charge per unit area per unit time

For a surface S'

$$\int_{S'} da \hat{n} \cdot \vec{j} = I \quad \text{current (charge per unit time) passing through surface } S'$$

Charge Conservation

vol V bounded by surface S'

$$\frac{d}{dt} \int_V d^3r \rho(\vec{r}, t) = - \oint_{S'} da \hat{n} \cdot \vec{j}$$

rate of change of total charge in V = $(-)$ charge flowing out of V through S' per unit time

$$\text{use } \oint_{S'} da \hat{n} \cdot \vec{j} = \int_V d^3r \vec{\nabla} \cdot \vec{j} = - \int_V d^3r \frac{\partial \rho(\vec{r}, t)}{\partial t}$$

$\Rightarrow$ local charge conservation

$$\boxed{\frac{\partial \rho}{\partial t} + \vec{\nabla} \cdot \vec{j} = 0}$$

A static situation has $\frac{\partial \mathcal{L}}{\partial t} = 0$

$\Rightarrow$ magnetostatics is defined by the condition $\vec{\nabla} \cdot \vec{j} = 0$

Differential formulation of Biot-Savart

For a set of charges q_i at $\vec{r}_i$ we have

$$\vec{B}(\vec{r}) = \sum_i k_5 q_i \vec{v}_i \times \frac{(\vec{r} - \vec{r}_i)}{|\vec{r} - \vec{r}_i|^3}$$

$$= k_5 \int d^3r' \vec{j}(\vec{r}') \times \frac{(\vec{r} - \vec{r}')}{|\vec{r} - \vec{r}'|^3}$$

$$= k_5 \int d^3r' \vec{j}(\vec{r}') \times \vec{\nabla} \left(\frac{-1}{|\vec{r} - \vec{r}'|} \right)$$

$$\vec{B}(\vec{r}) = k_5 \vec{\nabla} \times \left[\int d^3r' \frac{\vec{j}(\vec{r}')}{|\vec{r} - \vec{r}'|} \right]$$

where we used $\vec{\nabla} \times (\vec{A} \phi) = -\vec{A} \times \vec{\nabla} \phi$ when $\vec{A}$ is indep of $\vec{r}$

$\Rightarrow \boxed{\vec{\nabla} \cdot \vec{B} = 0}$ since $\vec{\nabla} \cdot (\vec{\nabla} \times \vec{A}) = 0$ for any vector function $\vec{A}$
integral form $\oint d\vec{a} \cdot \vec{B} = 0$

$$\vec{\nabla} \times \vec{B} = k_5 \vec{\nabla} \times \left(\vec{\nabla} \times \left(\int d^3r' \frac{\vec{j}(\vec{r}')}{|\vec{r} - \vec{r}'|} \right) \right)$$

$$\text{use } \vec{\nabla} \times (\vec{\nabla} \times \vec{A}) = \vec{\nabla} (\vec{\nabla} \cdot \vec{A}) - \nabla^2 \vec{A}$$

$$\vec{\nabla} \times \vec{B} = k_5 \vec{\nabla} \left[\int d^3r' \vec{\nabla} \cdot \left(\frac{\vec{J}(\vec{r}')}{|\vec{r} - \vec{r}'|} \right) \right] \\ - k_5 \int d^3r' \vec{J}(\vec{r}') \nabla^2 \left(\frac{1}{|\vec{r} - \vec{r}'|} \right)$$

in the 2nd term, $\nabla^2 \left(\frac{1}{|\vec{r} - \vec{r}'|} \right) = -4\pi \delta(\vec{r} - \vec{r}')$

in the 1st term, $\vec{\nabla} \cdot \frac{\vec{J}(\vec{r}')}{|\vec{r} - \vec{r}'|} = \vec{J}(\vec{r}') \cdot \vec{\nabla} \left(\frac{1}{|\vec{r} - \vec{r}'|} \right) \\ = -\vec{J}(\vec{r}') \cdot \vec{\nabla}' \left(\frac{1}{|\vec{r} - \vec{r}'|} \right)$

since $\vec{\nabla} = -\vec{\nabla}'$

So $\int d^3r' \vec{\nabla} \cdot \left(\frac{\vec{J}(\vec{r}')}{|\vec{r} - \vec{r}'|} \right) = - \int d^3r' \vec{J}(\vec{r}') \cdot \vec{\nabla}' \left(\frac{1}{|\vec{r} - \vec{r}'|} \right) \\ = \int d^3r' \left(\vec{\nabla}' \cdot \vec{J}(\vec{r}') \right) \left(\frac{1}{|\vec{r} - \vec{r}'|} \right)$

integrate by parts

surface term $\rightarrow 0$ as

we take surface $\rightarrow \infty$

since $\vec{J} \rightarrow 0$ as $r \rightarrow \infty$

But for magnetostatics $\vec{\nabla} \cdot \vec{J} = 0 \Rightarrow$ only 2nd term remains

Then, for magnetostatics

$$\vec{\nabla} \times \vec{B} = 4\pi k_5 \vec{J} \quad \text{Ampere's law}$$

integral form $\oint_C d\vec{l} \cdot \vec{B} = 4\pi k_5 \int_S da \hat{n} \cdot \vec{J}$

$\uparrow$ curve bounding surface $\uparrow$

Although above diff eqs were derived starting from a "non-relativistic"