

Note: Lorentz gauge condition does not uniquely determine  $\vec{A}$  and  $\phi$ . If one constructs  $\vec{A}$  and  $\phi$  obeying Lorentz gauge condition, and then constructs

$$\vec{A}' = \vec{A} + \vec{\nabla}\chi$$

$$\phi' = \phi - \frac{1}{c} \frac{\partial \chi}{\partial t}$$

then  $\vec{A}'$  and  $\phi'$  will also be in Lorentz gauge provided  $\square^2 \chi = 0$  (proof left to reader)

## 2) Coulomb Gauge

gauge constraint: require  $\vec{\nabla} \cdot \vec{A} = 0$

if  $\vec{A}$  is in the Coulomb Gauge, then

$\vec{A}' = \vec{A} + \vec{\nabla}\chi$  will also be in Coulomb gauge provided  $\nabla^2 \chi = 0$ .

Then Gauss' law becomes

$$\nabla^2 \phi + \frac{1}{c} \frac{\partial}{\partial t} (\vec{\nabla} \cdot \vec{A}) = -4\pi\rho$$

$$\Rightarrow \boxed{\nabla^2 \phi = -4\pi\rho} \quad \text{same as electrostatics!}$$

$$\Rightarrow \phi(\vec{r}, t) = \int d^3r' \frac{\rho(\vec{r}', t)}{|\vec{r} - \vec{r}'|}$$

no matter what motion the source  $\rho(\vec{r}, t)$  has!  $\phi$  is given by the instantaneous Coulomb potential even though electromagnetic fields have a finite velocity of propagation  $c$ !

Ampere's Law becomes:

$$-\nabla^2 A + \frac{1}{c^2} \frac{\partial^2 \vec{A}}{\partial t^2} = \frac{4\pi}{c} \vec{j} - \vec{\nabla} (\vec{\nabla} \cdot \vec{A} + \frac{1}{c} \frac{\partial \phi}{\partial t})$$

$$\Rightarrow \nabla^2 A = \frac{4\pi}{c} \vec{j} - \frac{1}{c} \vec{\nabla} \left( \frac{\partial \phi}{\partial t} \right)$$

$$\begin{aligned} \text{where } \vec{\nabla} \left( \frac{\partial \phi}{\partial t} \right) &= \vec{\nabla} \left[ \int d^3r' \frac{\partial \rho}{\partial t} \frac{1}{|\vec{r} - \vec{r}'|} \right] \\ &= - \vec{\nabla} \left[ \int d^3r' \frac{\vec{\nabla}' \cdot \vec{j}(\vec{r}', t)}{|\vec{r} - \vec{r}'|} \right] \quad \text{by continuity eqn.} \end{aligned}$$

To see the meaning of this term, recall - any vector function  $\vec{j}$  can be written as the sum of a curlfree and a divergenceless part

$$\vec{j} = \vec{j}_{||} + \vec{j}_{\perp} \quad \text{where} \quad \begin{aligned} \vec{\nabla} \times \vec{j}_{||} &= 0 & \text{curlfree} \\ \vec{\nabla} \cdot \vec{j}_{\perp} &= 0 & \text{divergenceless} \end{aligned}$$

where

$$\vec{j}_{||}(\vec{r}) = -\frac{1}{4\pi} \vec{\nabla} \int d^3r' \frac{\vec{\nabla}' \cdot \vec{j}(\vec{r}')}{|\vec{r} - \vec{r}'|} \quad \text{longitudinal part}$$

$$\begin{aligned} \vec{j}_{\perp}(\vec{r}) &= \cancel{\frac{1}{4\pi} \vec{\nabla} \times \int d^3r' \frac{\vec{\nabla}' \times \vec{j}(\vec{r}')}{|\vec{r} - \vec{r}'|}} \quad \text{transverse part} \\ &= \frac{1}{4\pi} \vec{\nabla} \times \int d^3r' \frac{\vec{\nabla}' \times \vec{j}(\vec{r}')}{|\vec{r} - \vec{r}'|} \end{aligned}$$

$$\text{So } \vec{\nabla} \left( \frac{\partial \phi}{\partial t} \right) = 4\pi \vec{j}_{||}, \text{ and}$$

$$\nabla^2 A = \frac{4\pi}{c} \vec{j} - \frac{4\pi}{c} \vec{j}_{||} = \frac{4\pi}{c} \vec{j}_{\perp}$$

Returning to Ampere's law we see that the term

$$\vec{\nabla} \left( \frac{\partial \phi}{\partial t} \right) = -\vec{\nabla} \int d^3r' \left[ \frac{\vec{\nabla}' \cdot \vec{j}(r', t)}{|\vec{r} - \vec{r}'|} \right]$$

$$= 4\pi \vec{j}_{||}(\vec{r}, t)$$

So Ampere's law becomes

$$\square^2 \vec{A} = \frac{4\pi}{c} \vec{j} - \frac{4\pi}{c} \vec{j}_{||}$$

$$\boxed{\square^2 \vec{A} = \frac{4\pi}{c} \vec{j}_{\perp}}$$

In Coulomb gauge, only the transverse part of  $\vec{j}$  serves as a source for  $\vec{A}$ .

$\vec{A}$  describes the transverse modes, i.e. the EM radiation (recall in EM waves, the fields are always  $\perp$  direction of propagation)

$\phi$  describes the longitudinal modes

Coulomb gauge is not Lorentz invariant - if  $\vec{\nabla} \cdot \vec{A} = 0$  in one inertial reference frame, in general  $\vec{\nabla} \cdot \vec{A} \neq 0$  in another.

In Coulomb gauge, if  $\rho = 0$ , then  $\phi = 0$  and

$$\vec{E} = -\frac{1}{c} \frac{\partial \vec{A}}{\partial t}$$

## Transverse + Longitudinal Parts of vector functions

To prove the preceding claim,  $\vec{f} = \vec{f}_{\parallel} + \vec{f}_{\perp}$ , where  $\vec{\nabla} \times \vec{f}_{\parallel} = 0$  and  $\vec{\nabla} \cdot \vec{f}_{\perp} = 0$ , we first desire to prove Helmholtz Theorem.

Helmholtz Theorem: For a vector function  $\vec{f}(\vec{r})$  if one knows the divergence and curl of  $\vec{f}$  then one can ~~uniquely~~ uniquely determine  $\vec{f}$  itself. That is, if

$$\vec{\nabla} \cdot \vec{f} = 4\pi D(\vec{r}) \quad \text{where } D(\vec{r}) \text{ is a known scalar function}$$

$$\vec{\nabla} \times \vec{f} = 4\pi \vec{C}(\vec{r}) \quad \text{where } \vec{C}(\vec{r}) \text{ is a known vector function}$$

~~Then one can solve for~~

And if well defined boundary conditions on  $\vec{f}$  are known (here we will assume  $\vec{f}(\vec{r}) \rightarrow 0$  as  $\vec{r} \rightarrow \infty$ ) then there is a unique solution for  $\vec{f}(\vec{r})$ .

We prove this by construction!

Assume a solution of the form

$$\vec{f} = -\vec{\nabla}\phi + \vec{\nabla} \times \vec{W} \quad \text{where } \phi \text{ is a scalar and } \vec{W} \text{ a vector}$$

Now we show that we can find such a solution

First consider

$$\vec{\nabla} \cdot \vec{f} = -\nabla^2 \varphi + \vec{\nabla} \cdot (\vec{\nabla} \times \vec{W}) = -\nabla^2 \varphi + 0 = 4\pi D(\vec{r})$$

So  $-\nabla^2 \varphi = 4\pi D(\vec{r})$  This is just Poisson's equation we saw in electrostatics  
Solution when  $\varphi(\vec{r}) \rightarrow 0$  as  $r \rightarrow \infty$  is given by

$$\boxed{\varphi(\vec{r}) = \int d^3r' \frac{D(\vec{r}')}{|\vec{r} - \vec{r}'|}}$$

Coulomb-like  
integral solution

Now consider

$$\begin{aligned} \vec{\nabla} \times \vec{f} &= -\vec{\nabla} \times \vec{\nabla} \varphi + \vec{\nabla} \times (\vec{\nabla} \times \vec{W}) = 0 - \nabla^2 \vec{W} + \vec{\nabla} (\vec{\nabla} \cdot \vec{W}) \\ &= 4\pi \vec{C}(\vec{r}) \end{aligned}$$

Choose a gauge in which  $\vec{\nabla} \cdot \vec{W} = 0$  (just like Coulomb gauge in magnetostatics)

$$\text{Then } -\nabla^2 \vec{W} = 4\pi \vec{C}(\vec{r})$$

$$\boxed{\vec{W}(\vec{r}) = \int d^3r' \frac{\vec{C}(\vec{r}')}{|\vec{r} - \vec{r}'|}}$$

just like solution  
for vector pot  $\vec{A}$   
in magnetostatics

So we have constructed a solution

$$\vec{f}(\vec{r}) = -\vec{\nabla} \varphi + \vec{\nabla} \times \vec{W}$$

$$= -\vec{\nabla} \int d^3r' \frac{D(\vec{r}')}{|\vec{r} - \vec{r}'|} + \vec{\nabla} \times \int d^3r' \frac{\vec{C}(\vec{r}')}{|\vec{r} - \vec{r}'|}$$

$$\text{where } \vec{\nabla} \cdot \vec{f} = 4\pi D \quad \text{and} \quad \vec{\nabla} \times \vec{f} = 4\pi \vec{C}$$

Note: For above solution to be well defined, the integrals must converge. They will converge if the "sources"  $D(\vec{r})$  and  $\vec{C}(\vec{r})$  are sufficiently "localized" in space, i.e.  $D(\vec{r}) \rightarrow 0$ ,  $\vec{C}(\vec{r}) \rightarrow 0$  sufficiently fast as  $\vec{r} \rightarrow \infty$ .

Now we show that the above solution is unique.

Suppose there was another solution  $\vec{g}$  such that

$$\vec{\nabla} \cdot \vec{g} = 4\pi D \quad \text{and} \quad \vec{\nabla} \times \vec{g} = 4\pi \vec{C}$$

Consider  $\vec{h} \equiv \vec{f} - \vec{g}$  then

$$\vec{\nabla} \cdot \vec{h} = 0 \quad \text{and} \quad \vec{\nabla} \times \vec{h} = 0$$

Can show that only such  $\vec{h}$  that also has  $\vec{h}(\vec{r}) \rightarrow 0$  as  $\vec{r} \rightarrow \infty$  is  $\vec{h} \equiv 0$ , so  $\vec{g} = \vec{f}$  and solution is unique.

As a consequence of Helmholtz Theorem we have also shown the following

- Any vector function  $\vec{F}$  can be written as terms of a scalar and vector potential

$$\vec{F} = -\vec{\nabla} \phi + \vec{\nabla} \times \vec{W}$$

or equivalently

② Any vector function  $\vec{F}$  can be written in terms of a curl free and a divergenceless part

$$\vec{F} = \vec{F}_{||} + \vec{F}_{\perp} \quad \text{where} \quad \vec{\nabla} \times \vec{F}_{||} = 0 \quad \text{curlfree}$$

$$\vec{\nabla} \cdot \vec{F}_{\perp} = 0 \quad \text{divergenceless}$$

$$\text{where} \begin{cases} \vec{F}_{||}(\vec{r}) = -\vec{\nabla} \phi(\vec{r}) = -\vec{\nabla} \int \frac{d^3 r'}{4\pi} \frac{[\vec{\nabla}' \cdot \vec{F}(\vec{r}')] }{|\vec{r} - \vec{r}'|} \\ \vec{F}_{\perp}(\vec{r}) = \vec{\nabla} \times \vec{W}(\vec{r}) = \vec{\nabla} \times \int \frac{d^3 r'}{4\pi} \frac{[\vec{\nabla}' \times \vec{F}(\vec{r}')] }{|\vec{r} - \vec{r}'|} \end{cases}$$

where in above we used  $\bar{\phi}(\vec{r}') = \frac{1}{4\pi} \vec{\nabla}' \cdot \vec{F}(\vec{r}')$

$$\bar{C}(\vec{r}') = \frac{1}{4\pi} \vec{\nabla}' \times \vec{F}(\vec{r}')$$

~~where~~  $\vec{F}_{||}$  is called the longitudinal part of  $\vec{F}$

$\vec{F}_{\perp}$  is called the transverse part of  $\vec{F}$

to understand the reason for these names, we need to consider the Fourier transform

Above can be generalized to situations where  $\vec{F}$  satisfies other boundary conditions, say has a specified value on a given boundary surface. One just replaces  $\frac{1}{|\vec{r} - \vec{r}'|}$  by the appropriate Green's function — see more to come!

## Discussion regarding Fourier transforms

$$\vec{f}(\vec{r}) = \int_{-\infty}^{\infty} \frac{d^3k}{(2\pi)^3} e^{i\vec{k}\cdot\vec{r}} \vec{f}(\vec{k}) \quad \text{Fourier transf}$$

$$\vec{f}(\vec{k}) = \int_{-\infty}^{\infty} d^3r e^{-i\vec{k}\cdot\vec{r}} \vec{f}(\vec{r}) \quad \text{inverse transf}$$

Some special cases well worth remembering

### ① Transform of Dirac function

$$\delta_{\vec{r}_0}(\vec{k}) \equiv \int d^3r e^{-i\vec{k}\cdot\vec{r}} \delta(\vec{r}-\vec{r}_0) = e^{-i\vec{k}\cdot\vec{r}_0}$$

$$\Rightarrow \delta(\vec{r}-\vec{r}_0) = \int_{-\infty}^{\infty} \frac{d^3k}{(2\pi)^3} e^{i\vec{k}\cdot\vec{r}} \delta_{\vec{r}_0}(\vec{k})$$

$$\boxed{\delta(\vec{r}-\vec{r}_0) = \int \frac{d^3k}{(2\pi)^3} e^{i\vec{k}_0(\vec{r}-\vec{r}_0)}}$$

or letting  $\vec{r} \leftrightarrow \vec{k}$  in the above

$$\delta(\vec{k}-\vec{k}_0) = \int \frac{d^3r}{(2\pi)^3} e^{i\vec{r}\cdot(\vec{k}-\vec{k}_0)}$$

### ② Transform of Coulomb potential $\frac{1}{|\vec{r}-\vec{r}'|}$

We know

$$\nabla^2 \left( \frac{1}{|\vec{r}-\vec{r}'|} \right) = -4\pi \delta(\vec{r}-\vec{r}')$$

Suppose  $f(\vec{k}) \equiv \int d^3r e^{-i\vec{k}\cdot\vec{r}} \frac{1}{|\vec{r}-\vec{r}'|}$  is the

Fourier transf of  $\frac{1}{|\vec{r}-\vec{r}'|}$

$$\text{Substitute } \left\{ \begin{aligned} \frac{1}{|\vec{r}-\vec{r}'|} &= \int \frac{d^3k}{(2\pi)^3} e^{i\vec{k}\cdot\vec{r}} f(\vec{k}) \\ \delta(\vec{r}-\vec{r}') &= \int \frac{d^3k}{(2\pi)^3} e^{i\vec{k}\cdot(\vec{r}-\vec{r}')} \end{aligned} \right.$$

into above Poisson equation

$$\nabla^2 \int \frac{d^3k}{(2\pi)^3} e^{i\vec{k}\cdot\vec{r}} f(\vec{k}) = -4\pi \int \frac{d^3k}{(2\pi)^3} e^{i\vec{k}\cdot(\vec{r}-\vec{r}')} f(\vec{k})$$

↑  
operates only on  $\vec{r}$   
so move inside integral

$$\nabla^2 e^{i\vec{k}\cdot\vec{r}} = \vec{\nabla} \cdot (\vec{\nabla} e^{i\vec{k}\cdot\vec{r}})$$

$$\textcircled{1} \quad \vec{\nabla} e^{i\vec{k}\cdot\vec{r}} = \sum_{i=1}^3 \hat{x}_i \frac{\partial}{\partial x_i} e^{i\vec{k}\cdot\vec{r}} = \sum_{i=1}^3 \hat{x}_i i k_i e^{i\vec{k}\cdot\vec{r}} \\ = i\vec{k} e^{i\vec{k}\cdot\vec{r}} \quad \text{where } \hat{x}_1, \hat{x}_2, \hat{x}_3 = \hat{x}, \hat{y}, \hat{z}$$

$$\textcircled{2} \quad \vec{\nabla} \cdot (i\vec{k} e^{i\vec{k}\cdot\vec{r}}) = (i\vec{k}) \cdot (i\vec{k}) e^{i\vec{k}\cdot\vec{r}} = -k^2 e^{i\vec{k}\cdot\vec{r}}$$

$$\text{so } \nabla^2 e^{i\vec{k}\cdot\vec{r}} = -k^2 e^{i\vec{k}\cdot\vec{r}}$$

Poisson equation gives

$$\int \frac{d^3k}{(2\pi)^3} e^{i\vec{k}\cdot\vec{r}} (-k^2) f(\vec{k}) = -4\pi \int \frac{d^3k}{(2\pi)^3} e^{i\vec{k}\cdot\vec{r}} e^{-i\vec{k}\cdot\vec{r}'}$$

$$\int \frac{d^3k}{(2\pi)^3} e^{i\vec{k}\cdot\vec{r}} [-k^2 f(\vec{k})] = \int \frac{d^3k}{(2\pi)^3} e^{i\vec{k}\cdot\vec{r}} [-4\pi e^{-i\vec{k}\cdot\vec{r}'}]$$

As is true for Fourier series, so it is true for Fourier transforms: If two functions are equal, then their Fourier transforms are equal.

$$\Rightarrow -k^2 f(\vec{k}) = -4\pi e^{-i\vec{k}\cdot\vec{r}'}$$

$$f(\vec{k}) = \frac{4\pi}{k^2} e^{-i\vec{k}\cdot\vec{r}'}$$

$\Rightarrow$  is the Fourier transform of  $\frac{1}{|\vec{r}-\vec{r}'|}$

---