

Spherical Coordinates

$$\nabla^2 \phi = \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial \phi}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial \phi}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 \phi}{\partial \varphi^2} = 0$$

$$\phi(r, \theta, \varphi) = R(r) \Theta(\theta) \Phi(\varphi)$$

$$r^2 \nabla^2 \phi = \Theta \Phi \frac{d}{dr} \left(r^2 \frac{dR}{dr} \right) + \frac{R \Phi}{\sin \theta} \frac{d}{d\theta} \left(\sin \theta \frac{d\Theta}{d\theta} \right) + \frac{R \Theta}{\sin^2 \theta} \frac{d^2 \Phi}{d\varphi^2} = 0$$

$$\frac{r^2 \sin^2 \theta}{\Phi} \nabla^2 \phi = \underbrace{\frac{\sin^2 \theta}{R} \frac{d}{dr} \left(r^2 \frac{dR}{dr} \right) + \frac{\sin \theta}{\Theta} \frac{d}{d\theta} \left(\sin \theta \frac{d\Theta}{d\theta} \right)}_{\substack{\text{depends only on } r \text{ and } \theta \\ = -\text{const}}} + \underbrace{\frac{1}{\Phi} \frac{d^2 \Phi}{d\varphi^2}}_{\substack{\text{depends only on } \varphi \\ = \text{const}}} = 0$$

take $\frac{1}{\Phi} \frac{d^2 \Phi}{d\varphi^2} = -m^2$

$$\Rightarrow \boxed{\Phi = e^{\pm i m \varphi}} \quad m \text{ integer for } 2\pi \text{ periodicity in } \varphi$$

$$\Rightarrow \frac{\sin^2 \theta}{R} \frac{d}{dr} \left(r^2 \frac{dR}{dr} \right) + \frac{\sin \theta}{\Theta} \frac{d}{d\theta} \left(\sin \theta \frac{d\Theta}{d\theta} \right) = m^2$$

$$\underbrace{\frac{1}{R} \frac{d}{dr} \left(r^2 \frac{dR}{dr} \right)}_{\substack{\text{depends only on } r \\ = \text{const}}} + \underbrace{\frac{1}{\Theta \sin \theta} \frac{d}{d\theta} \left(\sin \theta \frac{d\Theta}{d\theta} \right) - \frac{m^2}{\sin^2 \theta}}_{\substack{\text{depends only on } \theta \\ = -\text{const}}} = 0$$

call the const $l(l+1)$

For R

$$\frac{1}{r} \frac{d}{dr} \left(r^2 \frac{dR}{dr} \right) - l(l+1) = 0$$

Solutions are of the form $R(r) = a_l r^l + b_l r^{-(l+1)}$
substitute in to verify

$$\begin{aligned} \frac{d}{dr} \left(r^2 \frac{dR}{dr} \right) &= \frac{d}{dr} \left(r^2 (l a_l r^{l-1} - (l+1) b_l r^{-l-2}) \right) \\ &= \frac{d}{dr} (l a_l r^{l+1} - (l+1) b_l r^{-l}) \\ &= l(l+1) a_l r^l + l(l+1) b_l r^{-(l+1)} = l(l+1) R \end{aligned}$$

For Θ :

$$\frac{1}{\sin \theta} \frac{d}{d\theta} \left(\sin \theta \frac{d\Theta}{d\theta} \right) - \frac{m^2}{\sin^2 \theta} = -l(l+1)$$

$$\text{let } x = \cos \theta$$

$$dx = -\sin \theta d\theta$$

$$d\theta = \frac{-dx}{\sin \theta}$$

above becomes

$$0 \leq \theta \leq \pi$$

solutions for $-1 \leq x \leq 1$

correspond to $l \geq 0$ integers

$$\frac{d}{dx} \left[(1-x^2) \frac{d\Theta}{dx} \right] + \left[l(l+1) - \frac{m^2}{1-x^2} \right] \Theta = 0$$

Called generalized Legendre Equation - solutions are called the associated Legendre functions.

ordinary Legendre polynomials are solutions
for $m=0$

For the special case $m=0$, i.e. the solution has azimuthal symmetry and Φ does not depend on the angle φ (i.e. rotational symmetry about $\hat{z}$ axis),

we want the solutions to

$$\frac{d}{dx} \left[(1-x^2) \frac{d\Theta}{dx} \right] + \ell(\ell+1) \Theta = 0$$

The solutions are known as the Legendre polynomials, $P_\ell(x)$.

They are given, for ℓ integer, by

$$P_\ell(x) = \frac{1}{2^\ell \ell!} \left(\frac{d}{dx} \right)^\ell (x^2-1)^\ell \quad \text{Rodriguez's formula}$$

The lowest ℓ polynomials are

$$P_0(x) = 1$$

$$P_2(x) = \frac{1}{2}(3x^2-1)$$

$$P_1(x) = x$$

$$P_3(x) = \frac{1}{2}(5x^3-3x)$$

In general, $P_\ell(x)$ is a polynomial of order ℓ with only even powers if ℓ is even, and only odd powers if ℓ is odd. $\Rightarrow P_\ell(x) \begin{cases} \text{even in } x & \text{for } \ell \text{ even} \\ \text{odd in } x & \text{for } \ell \text{ odd} \end{cases}$

$P_\ell(x)$ is normalized so that $P_\ell(1) = 1$

Note: Legendre polynomials are only for integer $l \geq 0$.
what about solutions for non integer l ?

The $P_l(x)$ give one solution for each integer l .
But $P_l(x)$ are defined by a 2nd order differential equation - shouldn't there be a 2nd independent solution for each l ?

It turns out that these "2nd" solutions, as well as solutions for non integer l , all blow up at either $x = -1$ or $x = 1$, i.e. at $\theta = 0$ or $\theta = \pi$.

They therefore are physically unacceptable and we do not need to consider them. See Jackson B.2

The Legendre polynomials are orthogonal and form a complete set of basis functions on the interval $-1 \leq x \leq 1$.

$$\int_{-1}^1 dx P_l(x) P_m(x) = \int_0^\pi d\theta \sin\theta P_l(\cos\theta) P_m(\cos\theta) = \begin{cases} 0 & l \neq m \\ \frac{2}{2l+1} & l = m \end{cases}$$

$\Rightarrow$ we can expand any function $f(\theta)$, $0 \leq \theta \leq \pi$, as a linear combination of the $P_l(\cos\theta)$.
This is the reason they are useful for solving problems of Laplace's eqn with spherical boundary surfaces.

For $m \neq 0$, the solutions to (see Jackson 3.5)

$$\frac{d}{dx} \left[(1-x^2) \frac{d\Theta}{dx} \right] + \left[l(l+1) - \frac{m^2}{1-x^2} \right] \Theta = 0$$

are the associated Legendre functions $P_l^m(x)$.

For $P_l^m(x)$ to be finite in interval $-1 \leq x \leq 1$

one again finds that l must be integer $l \geq 0$, and integer m must satisfy $|m| \leq l$, i.e. $m = -l, -(l-1), \dots, 0, \dots, (l-1), l$.

For each l and m there is only one such non divergent solution.

It is typical to combine the solutions $P_l^m(\cos\theta)$ to the θ -part of the equation with the $\Phi_m(\varphi) = e^{im\varphi}$ solutions to the φ -part of the equation to define the spherical harmonics


$$Y_{lm}(\theta, \varphi) \equiv \sqrt{\frac{2l+1}{4\pi} \frac{(l-m)!}{(l+m)!}} P_l^m(\cos\theta) e^{im\varphi}$$

The Y_{lm} are orthogonal

$$\int_0^{2\pi} d\varphi \int_0^\pi d\theta \sin\theta Y_{l'm'}^*(\theta, \varphi) Y_{lm}(\theta, \varphi) = \delta_{ll'} \delta_{mm'}$$

and are a complete set of basis functions for expanding any function $f(\theta, \varphi)$ defined on the surface of a sphere.

Behavior of fields near conical hole or sharp tip



we now want to solve the $\nabla^2 \phi = 0$ with separation of variables, but now θ is restricted to range $0 \leq \theta \leq \beta$.

we still have azimuthal symmetry, but now, since we do not need solution to ϕ to be finite for all $\theta \in [0, \pi]$, but only $\theta \in (0, \beta)$, we have more solutions to the Θ equation, i.e. l does not have to be integer, - still need $l > 0$ to be finite at $\theta = 0$.

see Jackson sec. 3.4 for details.

Examples with azimuthal symmetry $m=0$

General solution to $\nabla^2 \phi = 0$ can be written in form

$$\phi(r, \theta) = \sum_{l=0}^{\infty} \left[A_l r^l + \frac{B_l}{r^{l+1}} \right] P_l(\cos \theta)$$

determine the A_l and B_l from the boundary conditions of the particular problem.

① Suppose one is given $\phi(R, \theta) = \phi_0(\theta)$ on surface of sphere of radius R .

To find solution of $\nabla^2 \phi = 0$ inside sphere

ϕ should not diverge at origin $\Rightarrow B_l = 0$ for all l

$$\phi(r, \theta) = \sum_{l=0}^{\infty} A_l r^l P_l(\cos \theta)$$

$$\Rightarrow \phi(R, \theta) = \phi_0(\theta) = \sum_{l=0}^{\infty} A_l R^l P_l(\cos \theta)$$

$$\begin{aligned} \Rightarrow \int_0^{\pi} d\theta \sin \theta \phi_0(\theta) P_m(\cos \theta) &= \sum_{l=0}^{\infty} A_l R^l \int_0^{\pi} d\theta \sin \theta P_l(\cos \theta) P_m(\cos \theta) \\ &= \sum_{l=0}^{\infty} A_l R^l \left(\frac{2}{2l+1} \right) \delta_{lm} \end{aligned}$$

$$= A_m R^m \frac{2}{2m+1}$$

$$A_m = \frac{2m+1}{2R^m} \int_0^{\pi} d\theta \sin \theta \phi_0(\theta) P_m(\cos \theta)$$

gives solution

To find solution of $\nabla^2 \phi = 0$ outside sphere

if require $\phi \rightarrow 0$ as $r \rightarrow \infty$, then $A_l = 0$ for all l

$$\phi(r, \theta) = \sum_{l=0}^{\infty} \frac{B_l}{r^{l+1}} P_l(\cos \theta)$$

$$\phi(R, \theta) = \phi_0(\theta) = \sum_{l=0}^{\infty} \frac{B_l}{R^{l+1}} P_l(\cos \theta)$$

gives
solution

$$B_m = \frac{2m+1}{2} R^{m+1} \int_0^\pi \sin \theta \phi_0(\theta) P_m(\cos \theta) d\theta$$

$$B_m = A_m R^{2m+1}$$

- (2) Suppose one is given surface charge density $\sigma(\theta)$ fixed on surface of sphere of radius R . What is ϕ inside and outside?

From previous example

$$\phi(r, \theta) = \begin{cases} \sum_{l=0}^{\infty} A_l r^l P_l(\cos \theta) & r < R \\ \sum_{l=0}^{\infty} \frac{B_l}{r^{l+1}} P_l(\cos \theta) & r > R \end{cases}$$

boundary conditions at $r = R$ on surface

(i) ϕ continuous

$$\Rightarrow \sum_{l=0}^{\infty} \left[A_l R^l - \frac{B_l}{R^{l+1}} \right] P_l(\cos \theta) = 0$$

If an expansion in Legendre polynomials vanishes for all θ , then each coefficient in the expansion must vanish

$$\Rightarrow A_l R^l = \frac{B_l}{R^{l+1}} \Rightarrow \boxed{B_l = A_l R^{2l+1}}$$

(ii) jump in electric field at σ

$$-\left. \frac{\partial \phi^{\text{out}}}{\partial r} \right|_{r=R} + \left. \frac{\partial \phi^{\text{in}}}{\partial r} \right|_{r=R} = 4\pi\sigma$$

$$\Rightarrow \sum_{l=0}^{\infty} \left[\frac{(l+1) B_l}{R^{l+2}} + l A_l R^{l-1} \right] P_l(\cos\theta) = 4\pi\sigma$$

$$\Rightarrow \sum_{l=0}^{\infty} \left[\frac{(l+1) A_l R^{2l+1}}{R^{l+2}} + l A_l R^{l-1} \right] P_l(\cos\theta)$$

$$\Rightarrow \sum_{l=0}^{\infty} (2l+1) R^{l-1} A_l P_l(\cos\theta) = 4\pi\sigma$$

$$(2m+1) R^{m-1} A_m \left(\frac{2}{2m+1} \right) = 4\pi \int_0^{\pi} d\theta \sin\theta \sigma(\theta) P_m(\cos\theta)$$

$$\boxed{A_m = \frac{4\pi}{2 R^{m-1}} \int_0^{\pi} d\theta \sin\theta \sigma(\theta) P_m(\cos\theta)}$$

Suppose $\sigma(\theta) = k \cos \theta$ what is ϕ ?

Note $\sigma(\theta) = k P_1(\cos \theta)$

hence only $A_1 \neq 0$ by orthogonality of $P_\ell(\cos \theta)$

$$\begin{aligned} A_1 &= \frac{4\pi k}{2} \int_0^\pi \sin \theta P_1(\cos \theta) P_1(\cos \theta) d\theta \\ &= \frac{4\pi k}{2} \left(\frac{2}{2+1} \right) = \frac{4\pi k}{3} \end{aligned}$$

$$\Rightarrow \phi(r, \theta) = \begin{cases} \frac{4\pi k}{3} r \cos \theta & r < R \\ \frac{4\pi k}{3} \frac{R^3}{r^2} \cos \theta & r > R \end{cases}$$

we will see that potential outside the sphere is that of an ideal dipole with dipole moment

$$p = \frac{4\pi R^3 k}{3}$$

Inside the sphere, the potential $\phi = \frac{4\pi k}{3} z$ where $z = r \cos \theta$. The electric field inside the sphere is therefore the constant


$$\vec{E} = -\vec{\nabla} \phi = -\frac{4\pi k}{3} \hat{z}$$

outside the sphere the field is

$$\vec{E} = -\vec{\nabla}\phi = -\frac{\partial\phi}{\partial r} \hat{r} - \frac{1}{r} \frac{\partial\phi}{\partial\theta} \hat{\theta}$$

$$= \frac{8\pi k R^3}{3} \cos\theta \hat{r} + \frac{4\pi k R^3}{3} \sin\theta \hat{\theta}$$

$$\vec{E} = \frac{4\pi R^3 k}{3} \frac{1}{r^3} \left[2\cos\theta \hat{r} + \sin\theta \hat{\theta} \right]$$



dipole field