

Another way to write homogeneous Maxwell Equations

Define  $\epsilon_{\mu\nu\lambda\sigma} = \begin{cases} +1 & \text{if } \mu\nu\lambda\sigma \text{ is even permutation} \\ & \text{of } 1234 \\ -1 & \text{if } \mu\nu\lambda\sigma \text{ is odd permutation} \\ & \text{of } 1234 \\ 0 & \text{otherwise} \end{cases}$

4-d Levi-Civita symbol

Define

$$\tilde{F}_{\mu\nu} = \frac{1}{2i} \epsilon_{\mu\nu\lambda\sigma} F_{\lambda\sigma} \quad \text{pseudo-tensor}$$

$$= \begin{pmatrix} 0 & -E_3 & E_2 & -iB_1 \\ E_3 & 0 & -E_1 & -iB_2 \\ -E_2 & E_1 & 0 & -iB_3 \\ iB_1 & iB_2 & iB_3 & 0 \end{pmatrix} \quad \begin{array}{l} \text{has wrong sign} \\ \text{under parity} \\ \text{transf} \end{array}$$

$$\frac{\partial \tilde{F}_{\mu\nu}}{\partial x^\nu} = 0 \quad \text{gives homogeneous Maxwell equations}$$

$$\left. \begin{aligned} \frac{1}{2} F_{\mu\nu} F_{\mu\nu} &= B^2 - E^2 \\ -\frac{1}{4} F_{\mu\nu} \tilde{F}_{\mu\nu} &= \vec{B} \cdot \vec{E} \end{aligned} \right\} \text{Lorentz invariant scalars}$$

If  $\vec{E} + \vec{B}$  ad  $|\vec{E}| = |\vec{B}|$  in one frame of reference, then it is so in all frames of reference.  
 $(\vec{E} \cdot \vec{B} = 0, |\vec{B}|^2 - |\vec{E}^2| = 0)$   
satisfied by EM waves in the vacuum

If  $\vec{E} - \vec{B} = 0$  in one frame, ad  $E^2 > B^2$ , then there exists a frame in which  $B' = 0$ . If in one frame  $\vec{E} \cdot \vec{B} = 0$  ad  $B^2 > E^2$ , then there exists a frame in which  $E' = 0$ .

From  $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$  we can get  
Linearity transf for  $E$  and  $B$

For a transformation from  $K$  to  $K'$  with  $K'$  moving  
with  $v$  along  $x$ , with respect to  $K$ ,

$$E'_1 = E_1$$

$$B'_1 = B_1$$

$$E'_2 = \gamma(E_2 - \frac{v}{c} B_3)$$

$$B'_2 = \gamma(B_2 + \frac{v}{c} E_3)$$

$$E'_3 = \gamma(E_3 + \frac{v}{c} B_2)$$

$$B'_3 = \gamma(B_3 - \frac{v}{c} E_2)$$

### Kinematics

"dot" is  $\frac{d}{ds}$



4-momentum  $p_\mu = m \dot{x}_\mu = m u_\mu = (m \gamma \vec{v}, \vec{mc}\gamma)$

$$p_\mu^2 = m^2 u_\mu^2 = -m^2 c^2$$

4-force  $K_\mu = (\vec{K}, i K_0)$  "Minkowski force"

Newton's 2nd law

$$m \frac{d^2 x_\mu}{ds^2} = K_\mu$$

$$\Rightarrow m \frac{d u_\mu}{ds} = \frac{d p_\mu}{ds} = K_\mu$$

$$p_\mu^2 = -m^2 c^2 \Rightarrow \frac{d}{ds} (p_\mu^2) = p_\mu \frac{d p_\mu}{ds} = p_\mu K_\mu = 0$$

$$\Rightarrow m \gamma \vec{v} \cdot \vec{K} - mc \gamma K_0 = 0 \quad \text{or}$$

$$K_0 = \frac{\vec{v}}{c} \cdot \vec{K}$$

Define the usual 3-force by

$$\frac{d\vec{p}}{dt} = \vec{F}$$

(we identify Newtonian momentum  $\vec{p}$  with the space components of  $P_\mu$ )

$$\frac{d\vec{p}}{ds} = \vec{K} \text{ and } \frac{d\vec{p}}{ds} = \gamma \frac{d\vec{p}}{dt} = \gamma \vec{F} \Rightarrow \vec{K} = \gamma \vec{F}$$

$$K_0 = \gamma \vec{v} \cdot \vec{F}$$

Consider <sup>4</sup>-component of Newton's eqn

$$\frac{m d u_4}{ds} = m \frac{d(\gamma c)}{ds} = i K_0 = i \gamma \frac{\vec{v}}{c} \cdot \vec{F}$$

$$d(mr) = \gamma \frac{\vec{v}}{c^2} \cdot \vec{F} ds = dt \frac{\vec{v} \cdot \vec{F}}{c^2} = d\vec{r} \cdot \vec{F}$$

Work-energy theorem:  $d(m\gamma c^2) = d\vec{r} \cdot \vec{F}$  = work done

$\Rightarrow d(m\gamma c^2)$  is change in <sup>kinetic</sup> energy

$E = m\gamma c^2$  is relativistic <sup>kinetic</sup> energy

$\vec{p}_\mu = (\vec{p}, \frac{iE}{c})$	$\vec{p} = m\gamma \vec{v}$
	$E = m\gamma c^2$

$$E = \frac{mc^2}{\sqrt{1 - \frac{v^2}{c^2}}} \approx mc^2 \left(1 + \frac{1}{2} \frac{v^2}{c^2}\right) = mc^2 + \frac{1}{2} mv^2$$

↑      ↑  
small  $v$       non-rel

$$\frac{d\vec{p}_\mu}{ds} = k_\mu \text{ is therefore}$$

relativistic analog of Newton's 3rd law  
as well as law of conservation of energy

restmass kinetic energy

## Lorentz force

$$\frac{dp_\mu}{ds} = K_\mu$$

what is the  $K_\mu$  that represents the Lorentz force  
and how can we write it in ~~relative~~ Lorentz  
covariant way?

$K_\mu$  should depend on the fields  $F_{\mu\nu}$   
and the particles trajectory  $x_\mu$

$$\text{as } \vec{v} \rightarrow 0 \quad \vec{K} = g \vec{E}$$

$K_\mu$  can't depend directly on  $x_\mu$  as should be  
indep of origin of coords. So can  
depend only on  $\overset{\circ}{x}_\mu, \overset{\circ}{x}_\mu, \text{etc.}$

as  $v \rightarrow 0$ ,  $K$  does not depend on the  
acceleration, so  $K$  does not depend on  $\overset{\circ}{x}_\mu$

$K_\mu$  only depends on  $F_{\mu\nu}$  and  $\overset{\circ}{x}_\mu$

We need to form a  $\eta$ -vector out of  
 $F_{\mu\nu}$  and  $\overset{\circ}{x}_\mu$  that is linear in the fields  $F_{\mu\nu}$   
and proportional to the charge  $g$ .

The only possibility is

$$g f(\overset{\circ}{x}_\mu^2) F_{\mu\nu} \overset{\circ}{x}_\nu$$

But  $\dot{x}_\mu^2 = -c^2$  is a constant. Choose  $f(x_\mu^2) = \frac{1}{2}$

$K\mu = \frac{g}{c} F_{\mu\nu} \dot{x}_\nu$  is only possibility

This gives force

$$\vec{F} = \frac{1}{g} \vec{K}$$

$$F_i = \frac{1}{g} K_i = \frac{g}{c} (F_{ij} \dot{x}_j + F_{i4} \dot{x}_4)$$

$$= \frac{g}{c} \left( \frac{\partial A_j}{\partial x_i} - \frac{\partial A_i}{\partial x_j} \right) \dot{x}_j + \frac{g}{c} (-iE_i)(ic\gamma)$$

$$= \frac{g}{c} [ \epsilon_{ijk} B_k \gamma v_j ] + \frac{g}{c} E_i c \gamma$$

$$= g E_i + g \epsilon_{ijk} \frac{v_j}{c} B_k$$

$$\vec{F} = g \vec{E} + g \frac{v}{c} \vec{v} \times \vec{B}$$

Lorentz force is the same form in all inertial frames.  
No relativistic modification is needed.

## Relativistic Larmor's formula

$$\text{non-relativistic } P = \frac{2}{3} \frac{\epsilon^2 [\text{alt o}]^2}{c^3}$$

Consider inertial frame in which charge is instantaneously at rest. Call this rest frame  $K$ .

$$\text{power radiated in } K \text{ is } \dot{P} = \frac{d\overset{\circ}{E}(t)}{dt}$$

where  $\overset{\circ}{E}$  is energy radiated. In  $K$ , the momentum density  $\overset{\circ}{T} = \frac{1}{4\pi c} \overset{\circ}{E} \times \overset{\circ}{B} \sim \overset{\circ}{F}$  is in outward radial direction. Integrating over all directions, the radiated momentum vanishes  $\overset{\circ}{P} = 0$

energy-momentum is a 4-vector  $(\overset{\circ}{P}, \frac{i}{c} \overset{\circ}{E})$

To get radiated energy in original frame  $K'$  we can use Lorentz transf

$$\frac{\overset{\circ}{E}}{c} = \gamma \left( \frac{\overset{\circ}{E}}{c} - \frac{\vec{v}}{c} \cdot \overset{\circ}{P} \right) \Rightarrow \overset{\circ}{E} = \gamma \overset{\circ}{E} \text{ as } \overset{\circ}{P} = 0$$

and  $dt = \gamma dt'$  is time interval in  $K'$   
 $(d\overset{\circ}{T} = 0 \text{ as charge stays at origin in } K')$

$$\text{So } \frac{d\overset{\circ}{E}}{dt} = \frac{\gamma d\overset{\circ}{E}}{\gamma dt'} = \frac{d\overset{\circ}{E}}{dt} \Rightarrow \dot{P} = \overset{\circ}{P}$$

radiated power is Lorentz invariant!

in  $\hat{K}$  we can use non-relativistic Larmor's formula since  $v=0$ . So

$$P = \frac{2}{3} \frac{\gamma^2 a^2}{c^3}$$

$\ddot{a}$  is acceleration in  $\hat{K}$

To write an expression with out explicitly making mention of frame  $\hat{K}$ , we need to find a Lorentz invariant scalar that reduces to  $a^2$  as  $v \rightarrow 0$ .

Only choice is  $\alpha_{\mu}^2$  the 4-acceleration  $\alpha_{\mu} = \frac{du_{\mu}}{ds}$ .

$$\alpha_{\mu} = \frac{du_{\mu}}{ds} = \gamma \frac{du_{\mu}}{dt} = \gamma \frac{d}{dt} (\gamma \vec{v}, cc\gamma)$$

$$\vec{\alpha} = \gamma^2 \frac{d\vec{v}}{dt} + \gamma \vec{v} \frac{d\gamma}{dt}$$

$$\alpha_4 = cc\gamma \frac{d\gamma}{dt}$$

$$\frac{d\gamma}{dt} = \frac{1}{dt} \left( \frac{1}{\sqrt{1-v^2/c^2}} \right) = \frac{\vec{v} \cdot d\vec{v}}{c^2 dt} = \frac{1}{c^2} \gamma^3 \vec{v} \cdot \vec{a}$$

$$\text{as } \vec{v} \rightarrow 0, \gamma \rightarrow 1, \frac{d\gamma}{dt} \rightarrow 0, \text{ so } \left\{ \begin{array}{l} \vec{\alpha} \rightarrow \frac{d\vec{v}}{dt} = \vec{a} \\ \alpha_4 \rightarrow 0 \end{array} \right.$$

$$\alpha_{\mu}^2 \rightarrow |\vec{a}|^2 \text{ as desired}$$

Relativistic Larmor's formula

$$P = \frac{2}{3} \frac{\gamma^2}{c^3} \alpha_{\mu}^2 = \frac{2}{3} \frac{\gamma^2}{c^3} (\ddot{u}_{\mu})^2$$

$$\alpha_\mu = \left( \gamma^2 \frac{d\vec{v}}{dt} + \gamma \vec{v} \frac{d\gamma}{dt} \rightarrow c\gamma \frac{d\gamma}{dt} \right)$$

$$\frac{d\gamma}{dt} = \frac{1}{c^2} \gamma^3 \vec{v} \cdot \vec{a}$$

$$\alpha_\mu = \left( \gamma^2 \vec{a} + \gamma^4 \frac{1}{c^2} (\vec{v} \cdot \vec{a}) \vec{v} \rightarrow \frac{\gamma^4 \vec{v} \cdot \vec{a}}{c^2} \right)$$

$$\alpha_\mu^2 = \gamma^4 a^2 + \gamma^8 \frac{(\vec{v} \cdot \vec{a})^2 v^2}{c^4} + \frac{2\gamma^6 (\vec{v} \cdot \vec{a})^2}{c^2} - \frac{\gamma^8 (\vec{v} \cdot \vec{a})^2}{c^2}$$

$$= \gamma^4 \left[ a^2 + \gamma^4 \frac{(\vec{v} \cdot \vec{a})^2}{c^2} \left( \frac{v^2}{c^2} - 1 \right) + \frac{2\gamma^2 (\vec{v} \cdot \vec{a})^2}{c^2} \right]$$

$$= \gamma^4 \left[ a^2 - \frac{\gamma^2 (\vec{v} \cdot \vec{a})^2}{c^2} + \frac{2\gamma^2 (\vec{v} \cdot \vec{a})^2}{c^2} \right]$$

$$\alpha_\mu^2 = \gamma^4 \left[ a^2 + \frac{\gamma^2 (\vec{v} \cdot \vec{a})^2}{c^2} \right]$$

$$\text{as } \vec{v} \rightarrow 0 \rightarrow \alpha_\mu^2 \rightarrow a^2$$

$\alpha_\mu^2 = \dot{a}^2$  Lorentz invariant

$\dot{a}$  = acceleration in instantaneous rest

For a charge accelerating in linear motion, frame

$$\alpha_\mu^2 = \gamma^2 \frac{d\vec{v}}{dt} \cdot (\vec{v} \cdot \vec{a}) = v^2 a^2$$

$$\alpha_\mu^2 = \gamma^4 a^2 \underbrace{\left( 1 + \frac{\gamma^2 v^2}{c^2} \right)}_{\gamma^2} = \gamma^6 a^2$$

$$P = \frac{2\gamma a^2}{3c^3} \gamma^6$$

For a charge in circular motion  $(\vec{v} \cdot \vec{a}) = 0$

$$\alpha_\mu^2 = \gamma^4 a^2$$

$$P = \frac{2\gamma a^2}{3c^3} \gamma^4$$