

Factorization of canonical partition function — the ideal gas

Consider a system of N noninteracting particles

$$\rightarrow \mathcal{H}[\vec{q}_i, \vec{p}_i] = \sum_{i=1}^N \mathcal{H}^{(1)}(\vec{q}_i, \vec{p}_i)$$

where $\mathcal{H}^{(1)}$ is the single particle Hamiltonian that depends only on the three coordinates $\vec{q}_i$ and three momenta $\vec{p}_i$ of particle i .

$$Q_N = \frac{1}{N! h^{3N}} \left(\prod_{i=1}^N \int d\vec{q}_i d\vec{p}_i \right) e^{-\beta \mathcal{H}}$$

$$= \frac{1}{N!} \left(\prod_{i=1}^N \int \frac{d\vec{q}_i d\vec{p}_i}{h^3} \right) e^{-\beta \sum_j \mathcal{H}^{(1)}(\vec{q}_j, \vec{p}_j)}$$

factor the exponential

$$= \frac{1}{N!} \prod_{i=1}^N \left(\int \frac{d\vec{q}_i d\vec{p}_i}{h^3} e^{-\beta \mathcal{H}^{(1)}(\vec{q}_i, \vec{p}_i)} \right)$$

↑
factor for particle i is
identical to factor for particle j

$$\rightarrow \boxed{Q_N = \frac{1}{N!} (Q_1)^N} \quad \text{for noninteracting particles}$$

where Q_1 is the one particle partition function

$$Q_1 = \int \frac{d\vec{q} d\vec{p}}{h^3} e^{-\beta H^{(1)}(\vec{q}, \vec{p})}$$

Apply to the ideal gas.

$$H^{(1)}(\vec{q}, \vec{p}) = \frac{p^2}{2m}$$

$$Q_1 = \int \frac{d\vec{q}}{h^3} \int d\vec{p} e^{-\beta \frac{p^2}{2m}}$$

$$\int d\vec{q} = V \quad \text{volume of system}$$

$$\int d\vec{p} e^{-\beta \frac{p^2}{2m}} = \left(\frac{2\pi m}{\beta} \right)^{3/2} \quad \text{3D Gaussian integral}$$

$$Q_1 = \frac{V}{h^3} (2\pi m k_B T)^{3/2}$$

$$\Rightarrow Q_N = \frac{1}{N!} \left(\frac{V}{h^3} \right)^N (2\pi m k_B T)^{3N/2}$$

$$A(T, V, N) = -k_B T \ln Q_N$$

$$\ln N! = N \ln N - N$$

using Stirling's formula

$$= -k_B T \left\{ N \ln \left[\frac{V}{h^3} (2\pi m k_B T)^{3/2} \right] - N \ln N + N \right\}$$

$$A(T, V, N) = -k_B T N - k_B T N \ln \left[\frac{V}{h^{3N}} (2\pi m k_B T)^{3/2} \right]$$

Compute average energy

$$\langle E \rangle = -\frac{\partial}{\partial \beta} (\ln Q_N) = -\frac{\partial}{\partial \beta} (-\beta A)$$

$$= -\frac{\partial}{\partial \beta} \left(N + N \ln \left[\frac{V}{h^{3N}} \left(\frac{2\pi m}{\beta} \right)^{3/2} \right] \right)$$

$$= -N \frac{\partial}{\partial \beta} (\ln \beta^{-3/2}) = \frac{3}{2} N \frac{\partial}{\partial \beta} \ln \beta = \frac{3}{2} N \frac{1}{\beta}$$

$$\langle E \rangle = \frac{3}{2} N k_B T \quad \text{as expected}$$

entropy

$$S = -\left(\frac{\partial A}{\partial T} \right)_{V,N} = k_B N + k_B N \ln \left[\frac{V}{h^{3N}} (2\pi m k_B T)^{3/2} \right]$$

$$+ k_B T N \frac{3}{2} \left(\frac{1}{T} \right)$$

↖ from derivative of log

$$S = \frac{5}{2} N k_B + N k_B \ln \left[\frac{V}{h^{3N}} (2\pi m k_B T)^{3/2} \right]$$

substitute in $k_B T = \frac{2}{3} \frac{E}{N}$ to get

$$\Rightarrow S(E, V, N) = \frac{5}{2} N k_B + N k_B \ln \left[\frac{V}{h^{3N}} \left(\frac{4\pi m E}{3N} \right)^{3/2} \right]$$

we have recovered the Sackur-Tetrode equation which we earlier derived from the microcanonical ensemble! Canonical and microcanonical approaches are equivalent.

Because in computing Q_N we sum over all states with any energy, as compared to computing Ω where we restrict the sum to states in a particular energy shell E , it is usually easier to compute Q_N , rather than Ω .

We introduced the canonical distribution as a means of describing a physical system in contact with a heat bath.

The canonical distrib gives the same result as the microcanonical because in the $N \rightarrow \infty$ (thermodynamic) limit, the canonical probability distribution

$$P(E) = \frac{\Omega(E) e^{-E/k_B T}}{\Delta Q_N(V, T)}$$

approaches a delta-function* at the most probable energy = average energy, as set by the temperature T .

We could alternatively introduced the canonical ensemble just as a mathematical trick for computing $\Omega(E)$, removing the constraint of constant energy E by means of a Lagrange multiplier.

* since $E \sim N$ increases as $N \rightarrow \infty$

and $\langle E^2 \rangle - \langle E \rangle^2$ increases as $\sqrt{N}$

it is not really $P(E)$ that approaches a well defined function as $N \rightarrow \infty$. Rather it is the distribution

$p(e \equiv E/N)$, the probability density to have an energy per particle e , that approaches a delta function as $N \rightarrow \infty$

Note:


$$Q_N(\beta) = \int \frac{dE}{\Delta} \Omega(E) e^{-\beta E}$$

$Q_N(\beta)$ is Laplace transform of $\frac{\Omega(E)}{\Delta}$

$\Rightarrow \frac{\Omega}{\Delta}$ is inverse Laplace transform of Q_N

$$\begin{aligned} \frac{\Omega(E)}{\Delta} &= \frac{1}{2\pi i} \int_{\beta' - i\infty}^{\beta' + i\infty} e^{\beta E} Q_N(\beta) d\beta \quad (\beta' > 0) \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{i(\beta' + i\beta'') E} Q_N(\beta' + i\beta'') d\beta'' \end{aligned}$$

where $\beta' = \text{Re}(\beta) = 0^+$



contour of integration lies to right of imaginary axis

entropy $S = k_B \ln \Omega$

Helmholtz $-\frac{A}{T} = k_B \ln Q_N$

$$-\frac{A}{T} = S - \frac{E}{T}$$

Helmholtz free energy
is Legendre transf of S with
respect to E

Thermodynamic potentials which are Legendre transforms of each other, have ensemble partition functions that are Laplace transforms of each other.

Maxwell velocity distribution revisited

We found that the canonical partition function for a gas of N particles is given by

$$Q_N(T, V) = \int \frac{dE}{\Delta} \Omega(E) e^{-\beta E} = \frac{1}{N!} \prod_{i=1}^N \int d^3 p_i d^3 q_i e^{-\beta H[\vec{q}_i, \vec{p}_i]} \frac{1}{h^{3N}}$$

from which we conclude that the probability density for the system to ^{have} total energy E is

$$P(E) = \frac{\frac{\Omega(E) e^{-\beta E}}{\Delta}}{\int \frac{dE}{\Delta} \Omega(E) e^{-\beta E}}$$

Since $\Omega(E)$ is the number of states with total energy E , and all states with energy E are equally likely, we can conclude that the probability density for the system to be in some particular state $\{\vec{q}_i, \vec{p}_i\}$ is

$$P(\{\vec{q}_i, \vec{p}_i\}) = \frac{e^{-\beta H[\vec{q}_i, \vec{p}_i]}}{\prod_{i=1}^N \int d^3 q_i d^3 p_i e^{-\beta H[\vec{q}_i, \vec{p}_i]}}$$

To get the probability density that one particular particle k has momentum $\vec{p}_k$ we integrate $P(\{\vec{q}_i, \vec{p}_i\})$ over all degrees of freedom except $\vec{p}_k$.

$$P(\vec{P}_k) = \frac{\prod_i' \int d^3 \vec{q}_i \int d^3 \vec{p}_i e^{-\beta H[\vec{q}_i, \vec{p}_i]}}{\substack{\uparrow \\ \text{all degrees of freedom} \\ \text{except } \vec{P}_k} \prod_i \int d^3 \vec{q}_i \int d^3 \vec{p}_i e^{-\beta H[\vec{q}_i, \vec{p}_i]}}$$

For a general Hamiltonian with interactions between the particles, the above integrations can be very hard to do. But for non-interacting particles, it is easy! when

$$H[\vec{q}_i, \vec{p}_i] = \sum_i H^{(1)}(\vec{q}_i, \vec{p}_i)$$

$\uparrow$
 single particle Hamiltonian

then one has

$$e^{-\beta H[\vec{q}, \vec{p}]} = e^{-\beta \sum_i H^{(1)}(\vec{q}_i, \vec{p}_i)} = \prod_i e^{-\beta H^{(1)}(\vec{q}_i, \vec{p}_i)}$$

and the probability distribution $P(\vec{P}_k)$ becomes

$$P(\vec{P}_k) = \frac{\int d^3 \vec{q}_k e^{-\beta H^{(1)}(\vec{q}_k, \vec{P}_k)} \prod_{i \neq k} \left(\int d^3 \vec{q}_i \int d^3 \vec{p}_i e^{-\beta H^{(1)}(\vec{q}_i, \vec{p}_i)} \right)}{\prod_i \left(\int d^3 \vec{q}_i \int d^3 \vec{p}_i e^{-\beta H^{(1)}(\vec{q}_i, \vec{p}_i)} \right)}$$

$$= \frac{\int d^3 \vec{q}_k e^{-\beta H^{(1)}(\vec{q}_k, \vec{P}_k)}}{\int d^3 \vec{q}_k \int d^3 \vec{p}_k e^{-\beta H^{(1)}(\vec{q}_k, \vec{p}_k)}}$$

where all other terms for particles $i \neq k$ cancel out in numerator and denominator.

For the ideal gas $H^{(1)}(\vec{p}, \vec{p}) = \frac{|\vec{p}|^2}{2m}$ is indep of $\vec{q}$. Hence the $\int d^3q$ integrals in numerator and denominator each give a factor of V and

$$P(\vec{p}) = \frac{e^{-\beta |\vec{p}|^2 / 2m}}{\int d^3p e^{-\beta |\vec{p}|^2 / 2m}} = \frac{e^{-|\vec{p}|^2 / 2mk_B T}}{(2\pi m k_B T)^{3/2}}$$

This is exactly the Maxwell velocity distribution that we derived earlier from kinetic theory.

NOTE:

1) Maxwell's probability distribution

$$E = \frac{p^2}{2m}$$

$P(E) \propto e^{-\beta E}$ is probability for a single particle to have energy E and holds only in the limit of non-interacting particles

2) The probability to be in a particular state i with total energy E_i in the canonical distribution is

$$P_i \propto e^{-\beta E_i}$$

and holds generally for any type of system. Here E_i is the total energy and " i " specifies a state of the entire system (NOT just one particle in the system)

Virial Theorem - Classical Systems Only

Consider $\langle x_i \frac{\partial H}{\partial x_j} \rangle = \frac{\int \prod_k dq_k dp_k x_i \frac{\partial H}{\partial x_j} e^{-\beta H}}{\int \prod_k dq_k dp_k e^{-\beta H}}$

where x_i and x_j are any of the $6N$ generalized coordinates
 $q, p \quad i=1, \dots, 3N$

$$\int \prod_k dq_k dp_k x_i \frac{\partial H}{\partial x_j} e^{-\beta H} = -\frac{1}{\beta} \int \prod_k dq_k dp_k x_i \frac{\partial}{\partial x_j} (e^{-\beta H})$$

integrate by parts with respect to x_j

to be assumed from now on,

$$= -\frac{1}{\beta} \int \prod_k dq_k dp_k x_i e^{-\beta H} \Big|_{x_j^{(1)}}^{x_j^{(2)}} + \frac{1}{\beta} \int \prod_k dq_k dp_k \left(\frac{\partial x_i}{\partial x_j} \right) e^{-\beta H}$$

↑ integral over all coordinates except x_j

↑ $x_j^{(1)}$ and $x_j^{(2)}$ are the extremal values of x_j

the boundary integral vanishes because H becomes infinite at the extremal values of any coordinate

- if x_j is a momentum p , then extremal values are $p = \pm \infty$ and $H \propto p^2/m \rightarrow \infty$.

- if x_j is a spatial coord q , then extremal values are at boundary of ~~system~~ system, where the potential energy confining the particle to the volume V becomes infinite.

$$\Rightarrow \int \prod_k dq_k dp_k x_i \frac{\partial H}{\partial x_j} e^{-\beta H} = \frac{1}{\beta} \int \prod_k dq_k dp_k \left(\frac{\partial x_i}{\partial x_j} \right) e^{-\beta H}$$

but $\frac{\partial x_i}{\partial x_j} = \delta_{ij}$

$$\Rightarrow \langle x_i \frac{\partial H}{\partial x_j} \rangle = \frac{1}{\beta} \delta_{ij} \frac{\int dx_k \int dp_k e^{-\beta H}}{\int dx_k \int dp_k e^{-\beta H}}$$

$$\boxed{\langle x_i \frac{\partial H}{\partial x_j} \rangle = k_B T \delta_{ij}} \Leftarrow \text{Virial Theorem}$$

If $x_i = x_j = p_i$ then

$$\langle p_i \frac{\partial H}{\partial p_i} \rangle = \langle p_i \dot{q}_i \rangle = k_B T$$

If $x_i = x_j = q_i$, then

$$\langle q_i \frac{\partial H}{\partial q_i} \rangle = -\langle q_i \dot{p}_i \rangle = k_B T$$

where we used Hamilton's eq's of motion

$$\partial H / \partial p_i = \dot{q}_i \text{ and } \partial H / \partial q_i = -\dot{p}_i$$

$$\Rightarrow \langle \sum_{i=1}^{3N} p_i \dot{q}_i \rangle = 3N k_B T$$

$$- \langle \sum_{i=1}^{3N} q_i \dot{p}_i \rangle = 3N k_B T \quad - \text{Virial Theorem} \\ \text{Clausius (1870)}$$