

Equation of state: low densities - virial expansion

$Z \ll 1$  "non-degenerate"

or help lowest terms in series expansion

$$\frac{P}{k_B T} = \frac{g_s}{\lambda^3} \left\{ \frac{f_{5/2}}{g_{5/2}} \right\} = \frac{g_s}{\lambda^3} \left( Z + \frac{Z^2}{2^{5/2}} + \dots \right) \quad - \text{FD}$$

$$+ \text{BE}$$

$$\frac{N}{V} = \frac{g_s}{\lambda^3} \left\{ \frac{f_{3/2}}{g_{3/2}} \right\} = \frac{g_s}{\lambda^3} \left( Z + \frac{Z^2}{2^{3/2}} + \dots \right)$$

$$\Rightarrow \frac{P}{k_B T} = \frac{N}{V} \frac{\left( Z + \frac{Z^2}{2^{5/2}} + \dots \right)}{\left( Z + \frac{Z^2}{2^{3/2}} + \dots \right)} = \frac{N}{V} \left( 1 + \frac{Z}{2^{5/2}} \right) \left( 1 \pm \frac{Z}{2^{3/2}} + \dots \right)$$

$$= \frac{N}{V} \left( 1 \pm \frac{Z}{2^{3/2}} \mp \frac{Z}{2^{5/2}} + \dots \right)$$

$$\frac{1}{2^{3/2}} - \frac{1}{2^{5/2}} = \frac{2}{2^{5/2}} - \frac{1}{2^{5/2}} = \frac{1}{2^{5/2}}$$

$$\phi V = N k_B T \left( 1 \pm \frac{Z}{2^{5/2}} + \dots \right)$$

↑ quantum correction to classical ideal gas law.

+ FD -  $\phi$  increases compared to classically

- effective repulsion due to Pauli exclusion

- BE -  $\phi$  decreases compared to classically

- effective attraction.

above is similar conclusion to what we saw from 2-particle density matrix.

for small  $Z$ , the leading term gives  $\frac{N}{V} = \frac{g_s}{\lambda^3} Z$

or  $Z = \left( \frac{N \lambda^3}{V g_s} \right) = \frac{N}{Q_1}$  - same result we had classically

$\Rightarrow$  small  $Z$  limit is the low density limit  $m \lambda^3 \ll 1$

$$\phi V = N k_B T \left( 1 \pm \frac{1}{2^{5/2} g_s} \frac{N \lambda^3}{V} + \dots \right) \quad \equiv \text{high T}$$

## Sommerfeld model of electrons in a conductor


Fermi gas - high density / low temperature limit  
 "degenerate" Fermi gas

Consider first  $T \rightarrow 0$

$$\langle n(\epsilon) \rangle = \frac{1}{e^{\beta(\epsilon-\mu)} + 1}$$

$$\text{as } T \rightarrow 0 \quad e^{\beta(\epsilon-\mu)} \rightarrow \begin{cases} \infty & \epsilon \geq \mu \\ 0 & \epsilon < \mu \end{cases}$$

$$\Rightarrow \langle n(\epsilon) \rangle \rightarrow \begin{cases} 0 & \epsilon \geq \mu \\ 1 & \epsilon < \mu \end{cases}$$



$\Rightarrow$  all states with  $\epsilon \geq \mu$  are filled, all states with  $\epsilon < \mu$  are empty. This is the  $T=0$  ground state of the Fermi gas.

We therefore see that  $\mu(T=0)$  is the energy of the highest energy single particle state that is occupied in the ground state. One calls this energy the Fermi-energy

$$\epsilon_F \equiv \mu(T=0)$$

$T=0$

$$N = g_s \sum_{\vec{k}} 1 \quad \text{count occupied states}$$

$\vec{k} \leftarrow \text{such that } \frac{\hbar^2 k^2}{2m} \leq \epsilon_F$

$$\frac{\hbar^2 k_F^2}{2m} \equiv \epsilon_F$$

$$\text{density } n = \frac{N}{V} = \frac{1}{V} g_s \sum_{\vec{k}} 1 = \int_0^{\epsilon_F} d\epsilon g(\epsilon)$$

$$= \frac{2g_s}{\sqrt{\pi}} \left( \frac{2\pi m}{\hbar^2} \right)^{3/2} \int_0^{\epsilon_F} d\epsilon \sqrt{\epsilon} = \frac{2g_s}{\sqrt{\pi}} \left( \frac{2\pi m}{\hbar^2} \right)^{3/2} \frac{2}{3} \epsilon_F^{3/2}$$


$$= \frac{2g_s}{\sqrt{\pi}} \left( \frac{2\pi m}{4\pi^2 \hbar^2} \right)^{3/2} \frac{2}{3} \epsilon_F^{3/2} = \frac{2g_s}{\sqrt{\pi}} \frac{2}{8\pi^{3/2}} \frac{2}{3} \left( \frac{2m}{\hbar^2} \right)^{3/2} \epsilon_F^{3/2}$$

$$= \frac{g_s}{6\pi^2} \left( \frac{2m \epsilon_F}{\hbar^2} \right)^{3/2}$$

$= k_F$

$$\boxed{\epsilon_F = \frac{\hbar^2}{2m} \left( \frac{6\pi^2 m}{g_s} \right)^{3/2}}$$

Now at finite T



region of energy where  $|m| >$  differs from ground state ( $T=0$ ) is a region of order  $k_B T$  about  $\mu$ .

So the T=0 approx is good when  $k_B T \ll \mu$   
~~since  $\mu(0) = \epsilon_F$~~  we have

Using  $\mu \approx \mu(0) = \epsilon_F$  we have

$$k_B T \ll \frac{\pi^2}{2m} \left( \frac{6\pi^2 m}{g_s} \right)^{2/3} \Rightarrow \frac{2\pi m k_B T}{\hbar^2} \ll \frac{1}{4\pi} \left( \frac{6\pi^2 m}{g_s} \right)^{2/3}$$

$$\Rightarrow \lambda^2 \gg 4\pi \left( \frac{g_s}{6\pi^2 m} \right)^{2/3}$$

$$\Rightarrow m \lambda^3 \gg \frac{(4\pi)^{2/3}}{6\pi^2} g_s = \frac{4}{3\sqrt{\pi}} g_s$$

so this is equivalent to a low T or a high density limit  
 $m \lambda^3 \gg 1$  - called the "degenerate" limit.

(just as the classical limit  $\lambda \approx m \lambda^3 \ll 1$  was a high T low density limit)

Fermi temperature  $T_F = \epsilon_F / k_B$ . Degenerate limit is  $T \ll T_F$

For electrons in a metal,  $T_F \approx 10000 \text{ K}$ .

So electrons in a metal are always in the degenerate limit.

$$g(\epsilon) = C\sqrt{\epsilon}$$

$$g(\epsilon_F) = C\sqrt{\epsilon_F}$$

$$n = \int_0^{\epsilon_F} d\epsilon g(\epsilon) = \int_0^{\epsilon_F} d\epsilon C \epsilon^{1/2} = \frac{2}{3} C \epsilon_F^{3/2}$$

$$\Rightarrow C = \frac{3}{2} \frac{n}{\epsilon_F^{3/2}}$$

$$\boxed{g(\epsilon_F) = \frac{3}{2} n \frac{\epsilon_F^{1/2}}{\epsilon_F^{3/2}} = \frac{3}{2} \frac{n}{\epsilon_F}}$$

$$g(\epsilon) = g(\epsilon_F) \frac{g(\epsilon)}{g(\epsilon_F)} = \frac{3}{2} \frac{n}{\epsilon_F} \sqrt{\frac{\epsilon}{\epsilon_F}}$$

Energy in the degenerate limit  $T=0$

$$\frac{E}{V} = \int_0^{E_F} d\epsilon g(\epsilon) \epsilon$$

$$g(\epsilon) = C \sqrt{\epsilon}$$

$$\text{with } C = \left(\frac{2\pi m}{h^2}\right)^{3/2} \frac{2g_s}{\pi}$$

$$n = \frac{N}{V} = \int_0^{E_F} d\epsilon g(\epsilon)$$

density of states

$$\Rightarrow \frac{E}{V} = C \int_0^{E_F} d\epsilon \epsilon^{3/2} = \frac{2}{5} C E_F^{5/2}$$

$$n = \frac{N}{V} = C \int_0^{E_F} d\epsilon \epsilon^{1/2} = \frac{2}{3} C E_F^{3/2}$$

$$\Rightarrow \frac{E}{V} = \frac{3}{5} \frac{N}{V} E_F$$

$$\frac{E}{V} = \frac{3}{5} n E_F$$

$$\boxed{\frac{E}{N} = \frac{3}{5} E_F}$$

↑ energy per volume

↑ energy per particle


Above gives  $T=0$  results. To get behavior at low  $T > 0$ , or to get quantities such as  $C_V = \left(\frac{\partial E}{\partial T}\right)_V$ , we need to get the next order terms in a low temperature expansion.

In general we need to do integrals of the form

$$\int d\epsilon \frac{\tilde{\phi}(\epsilon)}{z^{-1} e^{\beta\epsilon} + 1} = \int d\epsilon \tilde{\phi}(\epsilon) n(\epsilon), \quad \tilde{\phi}(\epsilon) \text{ some function}$$

ex: to compute  $n$ ,  $\tilde{\phi}(\epsilon) = g(\epsilon)$ ; to compute  $\frac{E}{V}$ ,  $\tilde{\phi}(\epsilon) = g(\epsilon) \epsilon$

## Single estimate of $C_V$



When we increase temperature to  $k_B T$ , the electrons near the Fermi energy  $\epsilon_F$  will increase their energy by an amount  $\sim k_B T$ . The number of such electrons ~~is roughly~~ per unit volume is roughly

$$g(\epsilon_F)(k_B T)$$

$\frac{1}{\text{density of states}}$  ↑ energy interval about  $\epsilon_F$  of states which ~~increase~~ get excited at  $\epsilon_F$

⇒ Increase in energy per unit volume is

$$\Delta U \sim (g(\epsilon_F) k_B T) (k_B T) \sim g(\epsilon_F) (k_B T)^2$$

$\frac{1}{\text{# electrons excited}}$

$\frac{1}{\text{excitation energy per excited electron}}$

for free electrons

$$\Rightarrow C_V = \frac{d\Delta U}{dT} \sim g(\epsilon_F) k_B^2 T = \frac{3}{2} \frac{m}{\epsilon_F} k_B^2 T = \frac{3}{2} m k_B \left(\frac{T}{T_F}\right)$$

The previous calculation gives the precise numerical coefficient