

Electrons in a crystal

When we discussed X-ray scattering we saw that the condition of constructive interference required that an incident wavevector $\vec{k}$ could only scatter into wavevectors $\vec{k}' = \vec{k} - \vec{K}$ where $\vec{K}$ was in the R.L.

For electrons in a crystal we expect similar behavior since our argument depended only on the wave nature of the scattering.

So a free electron plane wave state $e^{i\vec{k} \cdot \vec{r}}$ will get scattered into states $e^{i\vec{k}' \cdot \vec{r}}$ with $\vec{k}' = \vec{k} - \vec{K}$. We therefore expect the eigenstates of an electron in the periodic ionic potential of the crystal will be a mixture of these scattered states

$$\begin{aligned}\psi_{\vec{k}}(\vec{r}) &= \sum_{\vec{K}} e^{i(\vec{k} - \vec{K}) \cdot \vec{r}} c_{\vec{k} - \vec{K}} \\ &= e^{i\vec{k} \cdot \vec{r}} \sum_{\vec{K}} e^{-i\vec{K} \cdot \vec{r}} c_{\vec{k} - \vec{K}} \\ &= e^{i\vec{k} \cdot \vec{r}} u_{\vec{k}}(\vec{r}) \quad \text{where } u_{\vec{k}}(\vec{r}) = \sum_{\vec{K}} e^{-i\vec{K} \cdot \vec{r}} c_{\vec{k} - \vec{K}}\end{aligned}$$

$u_{\vec{k}}(\vec{r})$ is periodic on the Bravais lattice:

$$\begin{aligned}u_{\vec{k}}(\vec{r} + \vec{R}) &= \sum_{\vec{K}} e^{-i\vec{K} \cdot (\vec{r} + \vec{R})} c_{\vec{k} - \vec{K}} = \sum_{\vec{K}} e^{-i\vec{K} \cdot \vec{R}} e^{-i\vec{K} \cdot \vec{r}} c_{\vec{k} - \vec{K}} \\ &= \sum_{\vec{K}} e^{-i\vec{K} \cdot \vec{r}} c_{\vec{k} - \vec{K}} = u_{\vec{k}}(\vec{r}) \quad \text{since } e^{-i\vec{K} \cdot \vec{R}} = 1 \\ &\quad \text{for all } \vec{R} \text{ in R.L.}\end{aligned}$$

This then yields Bloch's Theorem which we prove more rigorously later.

The energy eigenstates of an electron in a potential with the periodicity of a Bravais lattice ~~can~~ can be written in the form

$$\psi_k(\vec{r}) = e^{i\vec{k} \cdot \vec{r}} u_k(\vec{r})$$

where $u_k(\vec{r} + \vec{R}) = u_k(\vec{r})$ is periodic on the B.L.

An equivalent alternative statement of Bloch's Theorem is

$$\psi_k(\vec{r} + \vec{R}) = e^{i\vec{k} \cdot \vec{R}} \psi_k(\vec{r})$$

If the wavevector $\vec{k}$ lies exactly on the Bragg plane bisected by R.L. vector $\vec{K}$, then the two free electron plane wave states $\vec{k}$ and $\vec{k}' = \vec{k} - \vec{K}$ are degenerate in energy and so we expect they must mix equally in forming the new eigenstates in the presence of the periodic potential. Since we start with two states $\vec{k}, \vec{k}'$ we should wind up with two new states. If they mix equally, their relative coefficients should just be a phase factor $e^{i\phi}$. Hence we expect the two eigenstates to have the form

$$\psi_1(\vec{r}) = e^{i\vec{k} \cdot \vec{r}} + e^{i\phi_1} e^{i\vec{k}' \cdot \vec{r}}$$

$$\psi_2(\vec{r}) = e^{i\vec{k} \cdot \vec{r}} + e^{i\phi_2} e^{i\vec{k}' \cdot \vec{r}}$$

The new eigenstates ψ_1 and ψ_2 should be orthogonal

$$0 = \langle \psi_1 | \psi_2 \rangle = \int d^3r \psi_1^*(\vec{r}) \psi_2(\vec{r})$$


$$= \int d^3r \left[e^{-i\vec{k}\cdot\vec{r}} + e^{-i\phi_1} e^{-i\vec{k}'\cdot\vec{r}} \right] \left[e^{i\vec{k}\cdot\vec{r}} + e^{i\phi_2} e^{i\vec{k}'\cdot\vec{r}} \right]$$

$$= \int d^3r \left[1 + e^{i(\phi_2 - \phi_1)} + e^{-i\phi_1} e^{i(\vec{k} - \vec{k}')\cdot\vec{r}} + e^{i\phi_2} e^{i(\vec{k}' - \vec{k})\cdot\vec{r}} \right]$$

these two terms integrate to zero - proof later

$$= V \left[1 + e^{i(\phi_2 - \phi_1)} \right]$$

$$= 0 \text{ only if } \phi_2 - \phi_1 = \pi$$



$$\vec{k}' = \vec{k} - \vec{K}$$

write $\vec{k} = \vec{k}_{||} + \vec{k}_{\perp}$
then $\vec{k}' = -\vec{k}_{||} + \vec{k}_{\perp}$

So

$$\psi_1(\vec{r}) = e^{i\phi_1/2} e^{i\vec{k}_{\perp}\cdot\vec{r}} \left\{ e^{-i\phi_1/2} e^{i\vec{k}_{||}\cdot\vec{r}} + e^{i\phi_1/2} e^{-i\vec{k}_{||}\cdot\vec{r}} \right\}$$

$$= e^{i\phi_1/2} e^{i\vec{k}_{\perp}\cdot\vec{r}} 2 \cos(\vec{k}_{||}\cdot\vec{r} - \phi_1/2)$$

Similarly

$$\psi_2(\vec{r}) = e^{i\phi_2/2} e^{i\vec{k}_{\perp}\cdot\vec{r}} 2 \cos(\vec{k}_{||}\cdot\vec{r} - \phi_2/2)$$

$$= e^{i\phi_2/2} e^{i\vec{k}_{\perp}\cdot\vec{r}} 2 \cos\left(\vec{k}_{||}\cdot\vec{r} - \frac{\phi_1}{2} - \frac{\pi}{2}\right)$$

$$= e^{i\phi_2/2} e^{i\vec{k}_{\perp}\cdot\vec{r}} 2 \sin\left(\vec{k}_{||}\cdot\vec{r} - \frac{\phi_1}{2}\right)$$

so

$$\left. \begin{aligned} |\psi_1(\vec{r})|^2 &\sim \cos^2(\vec{k}_{||} \cdot \vec{r} - \phi_1/2) \\ |\psi_2(\vec{r})|^2 &\sim \sin^2(\vec{k}_{||} \cdot \vec{r} - \phi_1/2) \end{aligned} \right\} \text{probability density}$$


reflection from Bragg plane gives standing waves in direction $\parallel$ to $\vec{k}$

Since $\vec{k}_{||} = \frac{\vec{k}}{2}$ we have $\vec{k}_{||} \cdot \vec{R} = \frac{\vec{k} \cdot \vec{R}}{2} = \pi n$ where n is an integer

$$\Rightarrow |\psi_1(\vec{r})|^2 \text{ and } |\psi_2(\vec{r})|^2 \text{ have the periodicity of the B.L.}$$

$$\propto |\psi_1(\vec{r} + \vec{R})|^2 \approx \cos^2(\vec{k}_{||} \cdot \vec{r} + \pi n - \phi_1/2) = \cos^2(\vec{k}_{||} \cdot \vec{r} - \phi_1/2) = |\psi_1(\vec{r})|^2$$

The only difference between ψ_1 and ψ_2 is their relative phase shift of $\frac{\pi}{2}$, which shifts the location of the peaks of $|\psi_1|^2$ and $|\psi_2|^2$ relative to the positions of the ions. By symmetry we expect one of these, say ψ_1 , will have its peaks positioned at the same sites as the ions, while the other, ψ_2 , will have its peaks positioned in between the sites of the ions. For example, if $\vec{k}$ is the smallest non-zero R.L. vector, $|\vec{k}| = \frac{2\pi}{a}$ with a = separation between ions, we expect,




The spatial periodicity of ψ_1 and ψ_2 is $\lambda = \frac{2\pi}{k_{11}} = \frac{2\pi}{(K/2)} = \frac{2\pi}{\pi/a} = 2a$. The spatial periodicity of $|\psi_1|^2$ and $|\psi_2|^2$ is therefore $\frac{1}{2}\lambda = a$. Since ψ_1 and ψ_2 are $\frac{\pi}{2}$ out of phase, then $|\psi_1|^2$ and $|\psi_2|^2$ are π out of phase, hence their relative peaks are separated by $a/2$ as in the diagram above.

Now when one turns on the attractive ionic potential, the state ψ_1 will have a lower energy than the state ψ_2 since ψ_1 has its peaks located at the ions, where the interaction is strongest, while ψ_2 has its peaks in between the ions, where the interaction is weakest.

Hence we expect that the two states ψ_1 and ψ_2 , which were degenerate in energy before turning on the ionic potential, now split in energy, with a finite energy gap between them, $E_1 < E_2$.

⇒ Ionic potential cause a gap to appear in the energy dispersion relation $E(\vec{k})$ whenever $\vec{k}$ crosses a Bragg plane. Since the Brillouin zones are formed by the intersections of Bragg planes, gaps in $E(\vec{k})$ open as $\vec{k}$ crosses the boundary of any Brillouin zone.


For a 1-dimensional B.L. of lattice constant a , We expect the dispersion relation to look as in the sketch below:



$R = ma$ m integers
 $K = nb$, $b = \frac{2\pi}{a}$, n integers
 gap open in $E(k)$
 every time k crosses
 the boundary of a
 Brillouin Zone

Bragg Planes are at
 $\frac{K}{2} = \left(\frac{2\pi n}{a}\right)\left(\frac{1}{2}\right) = \frac{\pi n}{a}$
 for n integers

For a 2-dimensional square B.L. of lattice constant a



Bragg Planes divide k -space
 into Brillouin Zones

1st BZ

2nd BZ

3rd BZ

whenever k crosses the
 surface of a BZ, there
 there is a discrete jump
 in the energy $E(k)$

Each BZ is a primitive cell
 of the R.L.

Each $\vec{k}$ in k -space can be written as

$\vec{k} = \vec{g} + \vec{K}$ with $\vec{K}$ a R.L. vector and $\vec{g}$ a vector in the 1st BZ. $\vec{g}$ is unique

$\Rightarrow$ each n^{th} BZ may be mapped onto the 1st BZ by translating its pieces by appropriate R.L. vectors $\vec{K}$

It is customary to label the eigenstates and eigenvalues by the $\vec{g}$ and by discrete index n . $\vec{g}$ is called the "crystal momentum" and n the "band index". The state $(\vec{g}, n)$ corresponds to the free electron state in the n^{th} BZ with wave vector $\vec{k} = \vec{g} + \vec{K}$ ($\vec{K}$ is the RL vector that translates $\vec{g}$ into the n^{th} BZ)

The wavefunctions $\psi_{\vec{g},n}$ and energies $E_n(\vec{g})$ are called the band structure

Born - von Karman boundary conditions and Fourier transforms for a Bravais Lattice

We generalize the idea of periodic (or Born-von Karman) boundary conditions to electron states on a Bravais lattice.

B.L. vectors $\vec{R} = n_1 \vec{a}_1 + n_2 \vec{a}_2 + n_3 \vec{a}_3$
 n_1, n_2, n_3 integers.

For a BL of finite size we have $0 \leq n_i < N_i$

$N = N_1 N_2 N_3$ is total number of points in the BL.

Total volume of this finite BL is

$$V = \underbrace{\vec{a}_1 \cdot (\vec{a}_2 \times \vec{a}_3)}_{\text{volume of primitive cell}} N$$

We want our electron wavefunctions to be periodic on such a finite BL

$$\psi(\vec{r} + N_i \vec{a}_i) = \psi(\vec{r})$$

As we saw earlier for free electrons, this imposes a constraint on the wave vectors $\vec{k}$ that can appear in the Fourier transform of $\psi(\vec{r})$.

Write $\psi(\vec{r})$ in terms of its Fourier transform

$$\psi(\vec{r}) = \sum_{\vec{k}} e^{i \vec{k} \cdot \vec{r}} c_{\vec{k}}$$

↑ Fourier coefficients for $\psi(\vec{r})$

Then

$$\begin{aligned}\psi(\vec{r} + N_i \vec{a}_i) &= \sum_{\vec{k}} e^{i\vec{k} \cdot (\vec{r} + N_i \vec{a}_i)} c_{\vec{k}} \\ &= e^{i N_i \vec{k} \cdot \vec{a}_i} \sum_{\vec{k}} e^{i\vec{k} \cdot \vec{r}} c_{\vec{k}} \\ &= e^{i N_i \vec{k} \cdot \vec{a}_i} \psi(\vec{r})\end{aligned}$$

$\Rightarrow$ must have $e^{i N_i \vec{k} \cdot \vec{a}_i} = 1$ for all $\vec{k}$
that appear in Fourier transform of $\psi(\vec{r})$

Write $\vec{k}$ in terms of the primitive vectors of the RL

$$\vec{k} = x_1 \vec{b}_1 + x_2 \vec{b}_2 + x_3 \vec{b}_3 \quad x_i \text{ are not in general integers}$$

$$\text{Then } e^{i N_i \vec{k} \cdot \vec{a}_i} = e^{i N_i 2\pi x_i} = 1$$

$$\Rightarrow x_i N_i = m_i \text{ integers}$$

$$x_i = \frac{m_i}{N_i}$$

So the allowed wavevectors are

$$\boxed{\vec{k} = \frac{m_1}{N_1} \vec{b}_1 + \frac{m_2}{N_2} \vec{b}_2 + \frac{m_3}{N_3} \vec{b}_3}$$

where m_1, m_2, m_3 are integers.

This is the Born - von Karman boundary conditions